



Beyond the dmft: The Human and Economic Cost of Early Childhood Caries

Paul S. Casamassimo, Sarat Thikkurissy, Burton L. Edelstein and Elyse Maiorini
J Am Dent Assoc 2009;140;650-657

The following resources related to this article are available online at jada.ada.org (this information is current as of March 8, 2011):

Updated information and services including high-resolution figures, can be found in the online version of this article at:

<http://jada.ada.org/cgi/content/full/140/6/650>

Information about obtaining **reprints** of this article or about permission to reproduce this article in whole or in part can be found at:

<http://www.ada.org/prof/resources/pubs/jada/permissions.asp>

Beyond the dmft

The human and economic cost of early childhood caries

Paul S. Casamassimo, DDS, MS; Sarat Thikkurissy, DDS, MS; Burton L. Edelstein, DDS, MPH; Elyse Maiorini, BS

The death of 12-year-old Deamonte Driver in 2007 as a result of untreated dental caries¹ gave lack of access to dental care a new and disturbing face for many Americans—that is, the potential for serious morbidity and even death resulting from dental caries. Behind the headlines and outrage surrounding that watershed event exists a shadow world of suffering, pain, diminished quality of life, and even death known to but a few on the front lines of the early childhood caries (ECC) epidemic. The educated, insured and employed, including many in the dental profession, have been shielded from the consequences of untreated ECC experienced by hundreds of thousands of the 4.5 million children who develop the condition annually. Death and serious morbidity resulting from ECC and its treatment are not new or confined to one disseminated infection. Death—resulting from local anesthetic overdose, sedation or general anesthesia mishap and even choking²—has in fact happened in attempts to treat this most common of all chronic childhood diseases. While these

ABSTRACT

Background. Early childhood caries (ECC) is the most common disease of childhood and often is accompanied by serious comorbidities affecting children, their families, the community and the health care system. This report describes morbidity and mortality associated with ECC and its treatment.

Methods. The authors reviewed the literature for descriptions and quantification of morbidity associated with ECC and organized a wide range of studies into a visual model—the morbidity and mortality pyramid—that begins to convey the breadth and depth of ECC's penetration.

Results. ECC exacts a toll on children, affecting their development, school performance and behavior, and on families and society as well. In extreme cases, ECC and its treatment can lead to serious disability and even death. In finding access to care and managing chronic pain and its consequences, families experience stress and, thus, a diminished quality of life. Communities devote resources to prevention and management of the condition. The health care system is confronted with management of the extreme consequences of ECC in hospital emergency departments and operating rooms.

Conclusions. Traditional epidemiologic measures such as the decayed-missing-filled teeth (dmft) index do not adequately portray the effects of ECC on children, families, society and the health care system.

Clinical Implications. The impact of prevention and management of ECC requires the attention of health care professionals and decision makers and extends well beyond the dental office to regulatory and child advocacy agencies as well as public health officials and legislators.

Key Words. Dental caries; quality of life; pediatric dentistry; caries susceptibility.

JADA 2009;140(6):650-657.

Dr. Casamassimo is a professor and the chair and chief of dentistry, Department of Dentistry, Nationwide Children's Hospital, 700 Children's Drive, Columbus, Ohio 43205, e-mail "casamasp@chi.osu.edu". Address reprint requests to Dr. Casamassimo.

Dr. Thikkurissy is an assistant professor, Division of Pediatric Dentistry, Department of Dentistry, The Ohio State University College of Dentistry, Columbus.

Dr. Edelstein is a professor of clinical dental medicine and clinical health policy and management and the chair, Section of Social and Behavioral Sciences, College of Dental Medicine, Columbia University, New York City.

Ms. Maiorini is a research assistant, Division of Community Health, College of Dental Medicine, Columbia University, New York City.

extreme events tend to come before the public, the daily consequences of ECC to families, the community and society often go unnoticed.

Traditional surveillance measures for ECC fall short in conveying ECC's impact and in connecting it to the full range of consequences experienced by affected children and their families. Dental epidemiologic measures such as the decayed, missing and filled surfaces (dmfs) and decayed, missing and filled teeth (dmft) indexes do not imply the full scope of the disease's impact on children, families, society and the health care system. According to Jacques,³ "The ultimate value of any classification system ... will be determined by its ability to provide the user with information that will assist in understanding or solving clinical problems."

Similarly, utilization measures such as treatment visits fail to impart a robust picture of the effects of ECC. Service utilization is an imperfect surrogate for morbidity,⁴ as the true impact of a disease often lies outside the realm of direct medical care delivery. While consequences of ECC—explored in this article—are far-ranging, supportive data often are dispersed, difficult to find or not collected at all. In this report, we review evidence from a variety of perspectives to argue that the consequences of symptomatic ECC are multiple and significant and that broader surveillance of the disease's impact is needed.

BEYOND THE BIOLOGICAL MODEL: A MORBIDITY PYRAMID

The presence of high levels of ECC, despite a reduction in permanent-dentition caries through fluoridation of water and use of fluoridated toothpastes, begs for a broader look at social and behavioral factors that correlate with this form of the disease. Fisher-Owens and colleagues⁵ proposed a model for pediatric oral disease that superimposes child, family and community factors over the classic Keyes biological model to indicate the influence of these nonbiological factors on caries initiation (Figure 1). Various additional research efforts support a relationship between ECC and these behavioral, environmental and social factors.⁶⁻¹⁰ Consistent with the "common determinants" approach to understanding disease occurrence in populations,¹¹ coincident morbidities are common in ECC.^{12,13}

A traditional method of describing disease impact is a morbidity and mortality (M&M)

pyramid, in which increasingly severe consequences are stacked one on another.¹⁴ ECC, with its low rate of associated fatality and high rate of dysfunction, takes on the classical shape, with a broad base and narrow apex (shown in Figure 2 [page 653] as Type A).¹⁵ A disease with a high degree of fatality, such as pancreatic cancer with its low five-year survival rate, may appear as an inverted pyramid¹⁶ (Type C), while one with an intermediate degree of fatality appears more as a rectangle¹⁷ (Type B). The M&M pyramid classification system provides a valuable means of quantifying disease impact through consequences ranging from mild dysfunction to death.

An M&M pyramid allows one to both observe a meaningful measure of consequence and relate that measure to other consequences. For example, for every death resulting from ECC or its treatment, one expects a certain number of hospital admissions, missed school days or episodes of pain-induced difficulty in eating or sleeping. Use of M&M pyramids can help clinicians relate the occurrence of comorbidities and, through intersecting tiers, identify groups at higher risk of experiencing illness or adverse effects.¹⁸ The study of M&M pyramids can highlight the magnitude of a disease's effect on society and its components.¹⁹ Through the examination of a disease's impact, health care professionals can better direct resource allocation and utilization of services to maximize prevention and treatment.²⁰ An M&M pyramid can represent dimensions of a disease in a range of patients, from those who are at low risk to those who experience the worst outcomes; therefore, it can provide insight into associated expenditures and loss of human capital.^{21,22}

Figure 3 (page 654) depicts a draft ECC M&M pyramid. Whereas clinicians know well that the impact of ECC involves many tiers, supportive data are insufficient to fully quantify these tiers. Some must be constructed with aggregate data, often without clarity or safeguards regarding accuracy. For example, an oft-cited statistic related to dental disease is the number of days of school attendance lost,²³ but this measure includes not only the days of school that children miss because of pain or dysfunction, but also elec-

ABBREVIATION KEY. **CT:** Computed tomography. **dmfs:** Decayed, missing and filled surfaces. **dmft:** Decayed, missing and filled teeth. **ECC:** Early childhood caries. **ED:** Emergency department. **M&M:** Morbidity and mortality.




Figure 1. A multifactorial model of early childhood caries depicting possible roles for the child, the family and the community beyond the classical biological infectious disease model. Reprinted with permission of the publisher from Fisher-Owens and colleagues.⁵

tive time away from school for routine dental visits. Constructing pyramids also is hampered in part by multiple consequences. For example, there is overlap within tiers, such as the total cost of a hospitalization for a child with ECC, which might include the costs of emergency department (ED) care as well as charges for a multiday stay and for treatment in the operating room. These significant events, even if measured carefully, still may not indicate the human costs to families arising from disruption of life, work and school because of the difficulty encountered in quantifying secondary impacts.

Mortality associated with dental caries and dental intervention. The case of Deamonte Driver probably is the best-known case of caries-related mortality. Lesser known are the cases of Alexander Callender, a 6-year-old Mississippi boy

who died of sepsis,²⁴ and Jackie Martinez, a 7-year-old California girl who choked on a crown during a dental visit.² Unknown to but a few involved in their care are the names of many other children who have died as a result of sedation mishaps or an overdose of local anesthetic during treatment for ECC. ECC-associated mortality secondary to infection and treatment likely never will be known owing to inadequate surveillance, lack of an ECC registry, issues of confidentiality, the terms of some legal settlements, missing or incorrect diagnoses, and even inconsistent diagnostic coding choices by hospitals and physicians. Among brain abscesses alone, 15 percent result from infections of unknown source, some or many of which may be of dental origin.²⁵ It is likely that mortality related to ECC and its treatment is underreported. Coté and col-

leagues,²⁶ in an attempt to identify pediatric deaths related to sedation during an almost 30-year period, commented that their study sample represented a gross underreporting while also stating that dental specialists were disproportionately represented among all pediatric health providers.

Hospital admissions, ED care and use of general anesthetics. In many hospitals' EDs, a leading pediatric admission symptom is dental pain. Families seek out ED dental care for a variety of reasons, including lack of a primary care dentist, inability to pay a dentist, a perception that their child is in serious danger or pain, and proximity or convenience.²⁷ ED dental intervention is, in most cases, limited to management of pain and infection, leaving the source untreated at significant cost to the patient, the hospital and society and impeding a system designed and staffed for emergent medical events.²⁸

Many ED admissions become prolonged hospitalizations for management of facial cellulitis. The length of stay averages five days but can be far longer, and the cost of care can be significant.²⁹ Ettlbrick and colleagues³⁰ reported in 2000 that the average cost of care across five children's hospitals for a single admission for odontogenic infection was \$3,223. As with ED management of ECC infection and pain, many such hospitalizations do not result in definitive care for either the offending tooth or other carious teeth. Worse, these interventions may have untoward consequences. In a recent study of pediatric patients with facial cellulitis, researchers found that ED physicians were more likely to order computed tomographic (CT) imaging than were pediatric dentists, with no difference in treatment outcome.³¹ This finding is most relevant because a growing body of literature suggests that head and neck CT imaging is responsible for an increase in thyroid cancer incidence in children.^{32,33}

Treatment under general anesthesia for extensive dental repair is another costly and potentially risky consequence of ECC. Tens of thousands of young children in the United States undergo restoration and extraction of teeth under general anesthesia annually. The absolute numbers are not known as, for example, dental office-based use of anesthetics may not be recorded and dental treatment performed in conjunction with physicians' surgical services may not be tracked for either medical or dental procedures in existing registries. Estimates available




Figure 2. Examples of the morbidity and mortality pyramid, the shape of which is based on the characteristics of an individual disease entity. Reprinted with permission of the publisher from Wadman and colleagues.¹⁵

are based largely on records of Medicaid-insured children treated in hospitals. White and colleagues³⁴ reported that more than 5,500 children received general anesthetics for dental services in North Carolina alone across a two-year period. Griffin and colleagues³⁵ reported that more than 2,100 Medicaid-covered children received dental treatment under general anesthesia in Louisiana in a one-year period, with 60 percent of these children being 3 years or younger. The costs of these services to families and the public are significant, with mean costs in this one study calculated at \$1,508 per admission.³⁵ Extrapolating these costs across tens of thousands of children who receive general anesthetic services annually in the United States exposes an expenditure of millions of dollars for treatment of a largely preventable disease. The human toll of treating children under general anesthesia also can be significant. Cravero and colleagues³⁶ included dental cases in their assessment of adverse events in sedation and general anesthesia, stating that of all patients who receive these services, the pediatric population is at highest risk and has the lowest tolerance for error.

ECC, child development and well-being.

Parents of children seeking emergency dental care reported that 19 percent of the children experienced interference with play, 32 percent with school, 50 percent with sleeping and 86 percent with eating.³⁷ As early as 1992, Acs and colleagues³⁸ reported a relationship between ECC



Figure 3. A proposed early childhood caries morbidity and mortality pyramid.

and failure to thrive—a condition of poor growth in young children—in a cohort of low-income children. Body measurements and blood test results indicative of malnourishment are significantly associated with severe ECC and suggest iron-deficiency anemia.³⁹ These reports and others stimulated interest in the effects of ECC and its treatment on child growth, looking at factors such as dysfunctions in eating, sleeping, mood and attention. Knowledge of the impact of ECC on the development of the child in physical, emotional and intellectual terms is fragmented. Episodic pain from dental caries is well-established as a constant finding, even from an early age, affecting up to 20 percent of preschoolers.^{37,39-41}

The effect of ECC-related pain on distraction from learning and school performance, while not generally measured, is significant. One cross-

sectional study reported that more than one in 10 schoolchildren experienced tooth pain,⁴² and another identified an association between poor systemic and oral health and poor school performance.^{43,44} Anecdotally, school systems nationwide, particularly those that serve a significant low-income population, report that dental problems contribute to learning difficulties. School nurses act as dental case managers on a daily basis.⁴⁵ A study in Michigan has documented loss of sleep, inability to concentrate in school and absences from school all caused by dental caries-related pain.⁴⁶

When the federal government last surveyed the impact of caries on the activity of children in 1996, a rough estimate of 9.3 percent of U.S. children younger than 5 years experienced “restricted activity days,” and an additional esti-

mated 3.7 percent of these preschoolers incurred “bed days” because of dental problems.⁴⁷ The study noted disparities by income and race that correlate closely with disparities in disease experience among subpopulations of children in the United States.

A growing concern in pediatric emergency medical care is acetaminophen toxicity caused by excessive administration of the drug by parents for management of pain.⁴⁸ Toxic doses of the medication can accumulate rapidly, causing liver damage in small children.⁴⁹ The use of acetaminophen for management of ECC-related pain is common, but its extent and effect remain unknown.

Consequences of oral infection for systemic health have been explored more completely in adults with regard to periodontal infections than in children with regard to odontogenic infections secondary to ECC. The biological rationale for oral-systemic associations, however, may have strong utility for anticipating the effect of chronic odontogenic infections on children’s overall well-being. Little has been studied about the effect of chronic pulpal infection, bacteremia and circulating inflammatory proteins on a young, healthy child’s function, growth and development, healing and disturbances in development of pain perception.⁵⁰ Similarly, little is yet known about the potential ECC-driven exacerbation of chronic pediatric diseases among children affected by such conditions as asthma and diabetes. Some research supports a role for ECC in both physical and emotional health, as ECC treatment ameliorates many ECC-related dysfunctions. For example, dental treatment of teeth with ECC results in pain elimination and has been associated with improved growth velocities⁵¹ and improved quality of life.⁵²⁻⁵⁴

The effects of ECC on family, community and health care systems. *Family.* At the level of family consequences, there is a troubling association between ECC and child maltreatment. Sheller and colleagues⁵⁵ concluded that a dysfunctional family or social situation can lead to a recurrence of ECC, often with emotional outbursts and the threat of or actual violence. The relationship between ECC and neglect is well-established, but only recently have child maltreatment experts included dental caries in their listing of health conditions that predispose children to maltreatment.^{56,57}

Community. The impact of ECC on communities is beginning to be realized. State officials

have begun to recognize that citizens are concerned about dental caries and access to care.^{58,59} Care of children with ECC consumes a disproportionate share of dental expenditures because of the typical extent of disease and concomitant costs of treatment under general anesthesia in the operating room. Recently adopted policies to address ECC (such as that of the American Academy of Pediatric Dentistry’s Clinical Affairs Committee⁵⁹) will bring children into dental offices far earlier in an attempt to prevent dental caries, but they will require an increase in or diversion of workforce and health expenditures. Whereas early interventions to prevent and manage ECC have been modeled as both cost effective⁶⁰ and cost saving,⁶¹ authors from the Centers for Disease Control and Prevention⁶² reported in 2005 that under current dental system capacity constraints, widespread implementation of early intervention may crowd out existing reparative care that is being provided to children with ECC.

Health care systems. ECC’s impact on health care systems may well be significant. Some have proposed significant changes in dental insurance coverage to reduce care for low-risk children and expand care for high-risk children.⁶³ Others propose reorganization of the dental workforce, either via creating an intermediate-level therapist capable of providing disease management, restorative care or both to affected children⁶⁴ or via expanding the engagement of medical providers.⁶² Both of these approaches will affect costs associated with reconfiguring the educational infrastructure, shifting practice patterns of the existing workforce and addressing the behavioral aspects of ECC management.

ECC has a tremendous, but often invisible, impact on society and the health care system. Recognition of its pervasiveness likely will drive oral health planning for the foreseeable future.

Growing evidence gathered with epidemiologic instruments supports a diminished quality of life for families with children affected by ECC,⁵²⁻⁵⁴ but the pressures families experience in a daily-life context are less clear. ECC is strongly associated with vulnerable subpopulations, including children of impoverished, minority, immigrant, migrant and homeless families whose social and economic capital is limited.⁶⁵ Loss of a job, loss of income for time spent taking a child to multiple dental appointments, the cost of transportation, taking time to find a willing dentist and financing

care are real and significant issues for these families, exacerbated in today's chaotic economy.

The downward spiral of ECC has become an all-too-familiar cycle. Children develop ECC that advances to a painful level. Parents delay care because of finances and access-to-care problems, and the child's condition continues to worsen until it becomes so acute as to demand intervention regardless of the effect on the family's resources. Often, when these families finally gain access to care, they encounter significant delays, because the available treatment resources are far outstripped by the need and demand for them. Finally, the cycle is perpetuated as ECC predisposes children to future caries in primary and permanent teeth.⁶⁶⁻⁶⁸

SUMMARY

Surveillance measures such as the percentage of children affected by ECC or disease extent as measured by dmfs and dmft indexes need to be supplemented by objective measures of disease consequences ranging from dysfunction to death. Meaningful assessments of the effect of ECC on child development, learning and family function and the economic burdens it places on families, communities and the health care system are needed to describe the importance of this preventable disease adequately. Until such a comprehensive assessment—including routine diagnostic coding—is developed, the epidemic of ECC likely will continue to put the health and lives of children at risk, because few public and professional policymakers will consider ECC of sufficient importance to take meaningful action. Only when the picture of an affected child's life replaces a technical dental statistic will society act to redress the most common of childhood's illnesses. ■

Disclosure. None of the authors reported any disclosures.

1. Bingaman J, Cardin BL. Preventing decay, preventing tragedy. *Washington Post* March 18, 2007:B08.

2. Hines L, Asbury J. Riverside girl, 7, dies days after choking on tooth. *The Press-Enterprise*. March 11, 2008. "www.pe.com/localnews/healthcare/stories/PE_News_Local_D_tooth12.1d27aa.html". Accessed April 15, 2000.

3. Jacques DP. Measuring morbidity. *Ann Surg* 2004;240(2):214-215.

4. Garrison HG, Rutherford G. Injury statistics. In: Fingerhut LA, ed. *Proceedings of the International Collaborative Effort on Injury Statistics Volume I*; 1995:28-1-28-5. Hyattsville, Md.: National Center for Health Statistics. Department of Health and Human Services publication PHS 95-1252.

5. Fisher-Owens SA, Gansky SA, Platt LJ, et al. Influences on children's oral health: a conceptual model. *Pediatrics* 2007;120(3):e510-e520.

6. Patrick DL, Lee RS, Nucci M, Grembowski D, Jolles CZ, Milgrom P. Reducing oral health disparities: a focus on social and cultural determinants. *BMC Oral Health* 2006;6(suppl 1):S4.

7. Finlayson TL, Siefert K, Ismail AI, Sohn W. Psychosocial factors and early childhood caries among low-income African-American children in Detroit. *Community Dent Oral Epidemiol* 2007;35(6):439-448.

8. Reisine S, Litt M, Tinanoff N. A biopsychosocial model to predict caries in preschool children. *Pediatr Dent* 1994;16(6):413-418.

9. Southward LH, Robertson A, Wells-Parker E, et al. Oral health status of Mississippi Delta 3- to 5-year-olds in child care: an exploratory study of dental health status and risk factors for dental disease and treatment needs. *J Public Health Dent* 2006;66(2):131-137.

10. Peres MA, de Oliveira Latorre Mdo R, Sheiham A, et al. Social and biological early life influences on severity of dental caries in children aged 6 years. *Community Dent Oral Epidemiol* 2005;33(1):53-63.

11. Sheiham A, Watt RG. The common risk factor approach: a rational basis for promoting oral health. *Community Dent Oral Epidemiol* 2000;28(6):399-406.

12. Larson K, Russ SA, Crall JJ, Halfon N. Influence of multiple social risk on children's health. *Pediatrics* 2008;121(2):337-344.

13. Mouradian WE, Huebner CE, Ramos-Gomez F, Slavkin HC. Beyond access: the role of family and community in children's oral health. *J Dent Educ* 2007;71(5):619-631.

14. Hayhurst ER. Review of industrial accident prevention: a scientific approach. *Am J Public Health Nations Health* 1932;22(1):119-120.

15. Wadman MC, Muelleman RL, Coto JA, Kellermann AL. The pyramid of injury: using ecodes to accurately describe the burden of injury. *Ann Emerg Med* 2003;42(4):468-478.

16. He AR, Lindenberg AP, Marshall JL. Biologic therapies for advanced pancreatic cancer. *Expert Rev Anticancer Ther* 2008;8(8):1331-1338.

17. March L, Lapsley M. What are the costs to society and the potential benefits from the effective management of early rheumatoid arthritis? *Best Pract Res Clin Rheumatol* 2001;15(1):171-185.

18. Australian Institute of Health and Welfare. *Australia's Health 2006*. Canberra, Australian Capital Territory, Australia: Australian Institute of Health and Welfare; 2006:37-59.

19. Rice DP. Cost of illness studies: what is good about them? *Inj Prev* 2000;6(3):177-179.

20. Hogan P, Dall T, Nikolov P; American Diabetes Association. Economic costs of diabetes in the US in 2002. *Diabetes Care* 2003;26(3):917-932.

21. Hodgson TA. Costs of illness in cost-effective analysis: a review of the methodology. *Pharmacoeconomics* 1994;6(6):536-552.

22. Rice DP, Kelman S, Miller LS, Dunmeyer S. *The Economic Costs of Alcohol and Drug Abuse and Mental Illness: 1985*. Rockville, Md.: National Institute on Drug Abuse; 1990.

23. Gift HC, Reisine ST, Larach DC. The social impact of dental problems and visits (published correction appears in *Am J Public Health* 1993;83(6):816). *Am J Public Health* 1992;82(12):1663-1668.

24. Palmer C. Maryland youth's death focuses attention on access to dental care. *ADA News*. March 19, 2007. "www.ada.org/prof/resources/pubs/adanews/adanewsarticle.asp?articleid=2403". Accessed April 15, 2009.

25. Mathisen GE, Johnson JP. Brain abscess. *Clin Infect Dis* 1997;25(4):763-779.

26. Coté CJ, Karl HW, Notterman DA, Weinberg JA, McCloskey C. Adverse sedation events in pediatrics: analysis of medications used for sedation. *Pediatrics* 2000;106(4):633-644.

27. Rowley ST, Sheller B, Williams BJ, Mancl L. Utilization of a hospital for treatment of pediatric dental emergencies. *Pediatr Dent* 2006;28(1):10-17.

28. Graham DB, Webb MD, Seale NS. Pediatric emergency room visits for nontraumatic dental disease. *Pediatr Dent* 2000;22(2):134-140.

29. Lin YT, Lu PW. Retrospective study of pediatric facial cellulitis of odontogenic origin. *Pediatr Infect Dis J* 2006;25(4):339-342.

30. Ettlbrick KL, Webb MD, Seale NS. Hospital charges for dental caries related emergency admissions. *Pediatr Dent* 2000;22(1):21-25.

31. Rawlins J. *Cost and Treatment of Dental Caries-Related Cellulitis* (master's thesis). Columbus, Ohio: The Ohio State University; 2008.

32. Mazonakis M, Tzedakis A, Damilakis J, Gourtsoyiannis N. Thyroid dose from common head and neck CT examinations in children: is there an excess risk for thyroid cancer induction? *Eur Radiol* 2007;17(5):1352-1357.

33. Jimenez RR, Deguzman MA, Shiran S, Karrellas A, Lorenzo RL. CT versus plain radiographs for evaluation of c-spine injury in young children: do benefits outweigh risks? *Pediatr Radiol* 2008;38(6):635-644.

34. White HR, Lee JY, Rozier RG. The effects of general anesthesia legislation on operating room visits by preschool children undergoing dental treatment. *Pediatr Dent* 2008;30(1):70-75.

35. Griffin SO, Gooch BF, Beltrán E, Sutherland JN, Barsley R. Dental services, costs, and factors associated with hospitalization for Medicaid-eligible children, Louisiana 1996-97. *J Public Health Dent* 2000;60(1):21-27.
36. Cravero JP, Blike GT, Beach M, et al. Incidence and nature of adverse events during pediatric sedation/anesthesia for procedures outside the operating room: report from the Pediatric Sedation Consortium. *Pediatrics* 2006;118(3):1087-1096.
37. Edelstein B, Vargas CM, Candelaria D, Vemuri M. Experience and policy implications of children presenting with dental emergencies to U.S. pediatric dentistry training programs. *Pediatr Dent* 2006;28(5):431-437.
38. Acs G, Lodolini G, Kaminsky S, Cisneros GJ. Effect of nursing caries on body weight in a pediatric population. *Pediatr Dent* 1992;14(5):302-305.
39. Clarke M, Locker D, Berall G, Pencharz P, Kenny DJ, Judd P. Malnourishment in a population of young children with severe early childhood caries. *Pediatr Dent* 2006;28(3):254-259.
40. Tickle M, Blinkhorn AS, Milsom KM. The occurrence of dental pain and extractions over a 3-year period in a cohort of children aged 3-6 years. *J Public Health Dent* 2008;68(2):63-69.
41. Siegal MD, Yeager MS, Davis AM. Oral health status and access to dental care for Ohio Head Start children. *Pediatr Dent* 2004;26(6):519-525.
42. Meadow D, Edelstein B. An evaluation of the management of dental emergencies by the school nurse. *Pediatr Dent* 1981;3(4):325-328.
43. Vargas CM, Macek MD, Goodman HS, Wagner ML. Dental pain in Maryland school children. *J Public Health Dent* 2005;65(1):3-6.
44. Blumenshine SL, Vann WF Jr, Gizlice Z, Lee JY. Children's school performance: impact of general and oral health. *J Public Health Dent* 2008;68(2):82-87.
45. Alameda County Public Health Department. More Than a Toothache: Untreated Dental Disease in Our School Children: The Alameda County Oral Health Needs Assessment of Kindergarten and 3rd Grade Children; 2006. Oakland, Calif.: Alameda County Public Health Department. "www.acphd.org/AXBYCZ/ADMIN/DataReports/morethanatoothacheformail2106.pdf". Accessed April 15, 2009.
46. Children's oral health: more vigilance needed—study shows effects on quality of life. *DentalUM* 2006;22(1):69-70. "www.dent.umich.edu/alumni/dentalum/springsummer2006/Childrens_Oral_Health.pdf". Accessed April 15, 2009.
47. Adams PF, Hendershot GE, Marano MA; Centers for Disease Control, National Center for Health Statistics. Current estimates from the National Health Interview Survey, 1996. *Vital Health Stat* 1999;10(200):1-203.
48. Squires RH Jr, Shneider BL, Bucuvalas J, et al. Acute liver failure in children: the first 348 patients in the pediatric acute liver failure study group. *J Pediatr* 2006;148(5):652-658.
49. Amar PJ, Schiff ER. Acetaminophen safety and hepatotoxicity: where do we go from here? *Expert Opin Drug Saf* 2007;6(4):341-355.
50. Schechter N. The impact of acute and chronic dental pain on child development. *J Southeast Soc Pediatr Dent* 2000;6(1):16-17.
51. Acs G, Shulman R, Ng MW, Chussid S. The effect of dental rehabilitation on the body weight of children with early childhood caries. *Pediatr Dent* 1999;21(2):109-113.
52. Filstrup SL, Briskie D, da Fonseca M, Lawrence L, Wandera A, Inglehart MR. Early childhood caries and quality of life: child and parent perspectives. *Pediatr Dent* 2003;25(5):431-440.
53. Low W, Tan S, Schwartz S. The effect of severe caries on the quality of life in young children. *Pediatr Dent* 1999;21(6):325-326.
54. Easton JA, Landgraf JM, Casamassimo PS, Wilson S, Ganzberg S. Evaluation of a generic quality of life instrument for early childhood caries-related pain. *Community Dent Oral Epidemiol* 2008;36(5):434-440.
55. Sheller B, Williams BJ, Hays K, Mancl L. Reasons for repeat treatment under general anesthesia for the healthy child. *Pediatr Dent* 2003;25(6):546-552.
56. Friedlaender EY, Rubin DM, Alpern ER, Mandell DS, Christian CW, Alessandrini EA. Patterns of health care use that may identify young children who are at risk for maltreatment. *Pediatrics* 2005;116(6):1303-1308.
57. Valencia-Rojas N, Lawrence HP, Goodman D. Prevalence of early childhood caries in a population of children with history of maltreatment. *J Public Health Dent* 2008;68(2):94-101.
58. Damiano PC, Momany ET, Willard JC, et al. First evaluation of the IowaCare Program: Executive Summary. Iowa City, Iowa: Public Policy Center, The University of Iowa; 2008. "http://ppc.uiowa.edu/download/IowaCareExecSummFINAL.pdf". Accessed April 15, 2009.
59. American Academy of Pediatric Dentistry Clinical Affairs Committee, Infant Oral Health Subcommittee; American Academy of Pediatric Dentistry Council on Clinical Affairs. Guideline on infant oral health care. *Pediatr Dent* 2008-2009;30(7 suppl):90-93.
60. Ramos-Gomez FJ, Shepard DS. Cost-effectiveness model for the prevention of early childhood caries. *J Calif Dent Assoc* 1999;27(7):539-544.
61. Zavras AI, Edelstein BL, Vamvakidis A. Health care savings from microbiological caries risk screening of toddlers: a cost estimation model. *J Public Health Dent* 2000;60(3):182-188.
62. Jones K, Tomar SL. Estimated impact of competing policy recommendations for age of first dental visit. *Pediatrics* 2005;115(4):258-263.
63. Edelstein BL, Schneider D, DiBiasi A. Dental benefits in the Medicaid/CHP+ Streamlining HIFA Waiver. Washington: Children's Dental Health Project; 2005.
64. Casamassimo PS. Commentary on "Developing a pediatric oral health therapist to help address oral health disparities among children" by David A. Nash. *J Dent Educ* 2004;68(1):20-21.
65. Edelstein BL. Dental care considerations for young children. *Spec Care Dentist* 2002;22(3 suppl):11S-25S.
66. Powell LV. Caries prediction: a review of the literature. *Community Dent Oral Epidemiol* 1998;26(6):361-371.
67. Wandera A, Bhakta S, Barker T. Caries prediction and indicators using a pediatric risk assessment teaching tool. *ASDC J Dent Child* 2000;67(6):408-412.
68. O'Sullivan DM, Tinanoff N. The association of early dental caries patterns with caries incidence in preschool children. *J Public Health Dent* 1996;56(2):81-83.