

Acquired Temporomandibular Disorders in Infants, Children, and Adolescents

Review Council

Council on Clinical Affairs

Latest Revision

2015

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that disorders of the temporomandibular joint (TMJ), masticatory muscles, and associated structures occasionally occur in infants, children, and adolescents. This guideline is intended to assist the practitioner in the recognition and diagnosis of temporomandibular disorders (TMD) and to identify possible treatment options. It is beyond the scope of this document to recommend the use of specific treatment modalities.

Methods

This guideline was originally developed by the Clinical Affairs Committee – Temporomandibular Joint Problems in Children Subcommittee and adopted in 1990. This document is a revision of the previous version, last revised in 2010. The update included an electronic search using the terms: temporomandibular disorder, TMJ dysfunction, TMD AND adolescents, TMD AND gender differences, TMD AND occlusion, TMD AND treatment; fields: all fields; limits: within the last 15 years, humans, English, clinical trials. The reviewers agreed upon the inclusion of 78 references to support this guideline. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

Background

Definition of TMD

TMD is a collective term for a group of musculoskeletal and neuromuscular conditions which includes several clinical signs and symptoms involving the muscles of mastication, the TMJ, and associated structures.¹ While TMD has been defined as “functional disturbances of the masticatory system”,² others include masticatory muscle disorders,³ degenerative and inflammatory TMJ disorders,⁴ and TMJ disk displacements⁵ under the umbrella of TMD.

In a recent update, the American Academy of Orofacial Pain divided TMD in two broad categories: TMJ disorders and masticatory muscle disorders.

Prevalence of TMD in children and adolescents

The reported prevalence of TMD in infants, children, and adolescents varies widely in the literature.⁶⁻¹⁰ This variation may be due to differences in populations studied, diagnostic criteria, examination methods, and inter- and/or intra-rated variations of examining practitioners.^{11,12} Prevalence of signs and symptoms increases with age. One study reported the TMD-related symptoms were rare in three- and five-year-olds whereas five to nine percent of 10- and 15-year olds reported more severe symptoms.¹³ Another found 4.2 percent of adolescents aged 12-19 years reported TMD pain.¹⁴ A study of the primary dentition reported 34 percent of patients with signs and/or symptoms of TMD.¹⁵ This could, in part, be due to inclusion of muscular signs versus symptoms. An epidemiological study of 4724 children aged five through 17 years reported 25 percent with symptoms. Clicking was seen in 2.7 percent of children in the primary dentition and 10.1 percent in late mixed dentition, and further increased to 16.6 percent in patients with permanent dentition.¹⁶ Headaches appear to be independently and highly associated with TMD in adolescents, with headaches most commonly occurring before the onset of TMD pain (odds ratio 9.4).¹⁷ Although TMD pain in children increases with age in both boys and girls, recent surveys have indicated a significantly higher prevalence of symptoms and greater need for treatment in girls than boys¹⁸ with the development of symptomatic TMD correlated with the onset of puberty in girls.^{19,20} For ages 16-19 years, 32.5 percent of girls compared to 9.7 percent of boys reported school absences and analgesic consumption due to TMD pain.¹⁸

Etiology of TMD

Temporomandibular disorders have multiple etiological factors.²¹ Research is insufficient to reliably predict which patients will or will not develop TMD as most published investigations evaluate static, morphologic variables rather than study the dynamic relationship between the joint and

ABBREVIATIONS

AAPD: American Academy Pediatric Dentistry. **CBCT:** Cone-beam computed tomography. **TMD:** Temporomandibular disorder. **TMJ:** Temporomandibular joint.

the teeth during function.²² There are predisposing (or risk) factors, precipitating (or initiating) factors, and perpetuating (or sustaining) factors that contribute to the development of TMD.²² Many studies show a poor correlation between any single etiological factor and resulting signs (i.e., findings identified by the dentist during the examination) and symptoms (i.e., findings reported by the child or parent). Alterations in any one or a combination of teeth, periodontal ligament, the TMJ, or the muscles of mastication may lead to TMD.²³ Furthermore, systemic and psychosocial factors may reduce the adaptive capacity of the masticatory system and contribute to TMD.¹

Etiologic factors suggested as contributing to the development of TMD are:

1. Macrotrauma: This would include impact injuries such as trauma to the chin. A common occurrence in childhood because of falling, chin trauma is reported to be a factor in the development of TMD in pediatric patients.²⁵⁻²⁷ Additional macrotraumatic injuries occur due to motor vehicle accidents, sports, physical abuse, forceful intubation, and third molar extraction.²⁸ Unilateral and bilateral intracapsular or subcondylar fractures are the most common mandibular fractures in children.²⁹ Closed reduction and prolonged immobilization can result in ankylosis.^{30,31} Improperly treated fractures may result in facial asymmetry.^{30,31}
2. Microtrauma from parafunctional habits: Bruxism, clenching, hyperextension, and other repetitive habitual behaviors are thought to contribute to the development of TMD by joint overloading that leads to cartilage breakdown, synovial fluid alterations, and other changes within the joint.³² Bruxism may occur while the patient is asleep or awake; sleep bruxism is a different entity from daytime bruxism. Sleep bruxism has been classified as a sleep-related movement disorder.³³ A study of 854 patients younger than 17 years old found the prevalence of bruxism to be 38 percent,³⁴ but studies do not distinguish between sleep or daytime bruxism. The literature on the association between parafunction and TMD in pediatric patients is contradictory.³⁵⁻³⁷ However, childhood parafunction was found to be a predictor of the same parafunction 20 years later.³⁸ Other studies found correlations between reported bruxism and TMD³⁹ with a 3.4 odds ratio.⁴⁰ Children who grind their teeth were found to complain more often of pain and muscle tenderness when eating.⁴¹ Other examples of microtrauma include repetitive strain such as playing a wind instrument, fingernail biting,²² or another activity in which the mouth is held open outside of rest position.
3. Anatomical factors (skeletal and occlusal): There is a relatively low association of occlusal factors and the development of TMD.^{42,43} It is reasonable that some occlusal factors may place greater adaptive demands on the masticatory system. Current literature does not support that the development of TMD is caused or improved by

orthodontic treatment,⁴⁴⁻⁴⁸ regardless of whether premolars were extracted prior to treatment.²⁸ Changes in free-way dimension of the rest position (normally two to four millimeters) may be impinged by occlusal changes, disease, muscle spasms, nervous tension, and/or restorative prosthetics.² While most children and adolescents may be able to compensate without problem, in others, failure of the masticatory system to adapt may lead to greater risk of dysfunction. In a study of 4,724 children aged five-17 years grouped by stage of dental development, the following malocclusions were found to be associated with TMD:¹⁶

- Skeletal anterior open bite.¹⁶
- Steep articular eminence of the temporal bone.¹
- Overjet greater than six to seven millimeters.^{16,49,50}
- Class III malocclusion.¹⁶
- Posterior crossbite.

Craniocervical posture has been suggested to be associated with occlusion and with dysfunction of the TMJ, including abnormalities of the mandibular fossa, condyle, ramus, and disc. Cervical pain and dysfunction can be a result of poor posture. Cervical pain is frequently referred to orofacial structures and can be misinterpreted as TMD.²⁴

4. Psychosocial factors: Psychosocial factors may play a part in the etiology of TMD.⁵¹ Emotional stress predisposes to clenching and bruxism which in turn contribute to orofacial pain. Results from a case-control study indicate that management of stress and anxiety can mitigate the signs and symptoms of TMD.⁵¹ Depression, anxiety, post-traumatic stress disorder, psychologic distress, and sleep dysfunction may influence TMD prognosis and symptoms.⁵² Behavioral factors such as somatization and depression influence TMD pain to a larger degree in girls than in boys. Higher pain intensity in the orofacial region correlated with greater impact on quality of life including difficulty with prolonged jaw opening, eating hard/soft foods, and sleeping.⁵³
5. Systemic factors: Systemic factors contributing to TMD include connective tissue diseases such as rheumatoid arthritis, systemic lupus erythematosus, juvenile idiopathic arthritis, and psoriatic arthritis.²² These systemic diseases occur as a result of imbalance of pro-inflammatory cytokines which causes oxidative stress, free radical formation, and ultimately joint damage.⁵⁴ Other systemic factors may include joint hypermobility, genetic susceptibility, and hormonal fluctuations. Generalized joint laxity or hypermobility (e.g., Ehler Danlos syndrome) has been cited but has a weak association with TMD.⁵⁵ There is little research in regard to the genetic susceptibility for development of TMD. Recently, study of catechol-O-methyl-transferase (COMT) haplotypes found that the presence of one low pain sensitivity haplotype decreased the risk of developing TMD.⁵⁶

The role of hormones in the etiology of TMD is debatable. Randomized controlled trials indicate that estrogen does not play a role in the etiology of TMD, whereas cohort and case-controlled studies show the opposite.¹ Although the biological basis for gender-based disparity in TMD is unclear, the time course of symptoms is of note in females. Additional studies have shown that TMJ pain and other symptoms vary in relation to phases of the menstrual cycle.⁵⁷ The suggestion of a hormonal influence in development of TMD is supported clinically by a study of 3,428 patients who sought treatment for TMD. This study revealed that 85.4 percent of patients seeking treatment were female and the peak age for treatment seeking was 33.8 years.⁵⁸ In a similar study of adolescents,²² 15.1 percent of all patients evaluated for TMD were less than 20 years of age and girls accounted for 89.9 percent of patients aged 15-19 seeking care and 75.5 percent of patient six-14 years of age.

Diagnosing TMD

All comprehensive dental examinations should include a screening evaluation of the TMJ and surrounding area.⁵⁹⁻⁶¹ Diagnosis of TMD is based upon a combination of historical information, clinical examination, and/or craniocervical and TMJ imaging.⁶² The findings are classified as symptoms and signs.⁵⁹ These symptoms may include pain, headache, TMJ sounds, TMJ locking, and ear pain. Certain medical conditions are reported to occasionally mimic TMD. Among these differential diagnoses are trigeminal neuralgia, central nervous system lesions, odontogenic pain, sinus pain, otological pain, developmental abnormalities, neoplasias, parotid diseases, vascular diseases, myofascial pain, cervical muscle dysfunction, and Eagle's syndrome.²⁴ Other common medical conditions (e.g., otitis media, allergies, airway congestion, rheumatoid arthritis) can cause symptoms similar to TMD.⁶³

Clinical and physical assessment of the patient may include history and determination of joint sounds, evaluation of mandibular range of motion, appraisal of pain, evaluation for signs of inflammation, and select radiographic examination.

A screening history, as part of the health history, may include questions such as:^{1,22}

- Do you have difficulty opening your mouth?
- Do you hear noises within your jaw joint?
- Do you have pain in or around your ears or your cheeks?
- Do you have pain when chewing, talking, or using your jaws?
- Do you have pain when opening your mouth wide or when yawning?
- Has your bite felt uncomfortable or unusual?
- Does your jaw ever lock or go out?
- Have you ever had an injury to your jaw, head, or neck? If so, when? How was it treated?
- Have you previously been treated for a temporomandibular disorder? If so, when? How was it treated?

Physical assessment should include the following:

1. Palpation of the muscles of mastication and cervical muscles for tenderness, pain, or pain referral patterns.¹
2. Palpation of the lateral capsule of the TMJs.²²
3. Mandibular function and provocation tests.^{1,22}
4. Palpation and auscultation for TMJ sounds.²²
5. Mandibular range of motion.²²

Evaluation of jaw movements including assessment of mandibular range of motion using a millimeter ruler (i.e., maximum unassisted opening, maximum assisted opening, maximum lateral excursion, maximum protrusive excursion) and mandibular opening pattern (i.e., symmetrical vs. asymmetrical). Both limited and excessive mandibular range of motion may be seen in TMD.²²

TMJ imaging is recommended when there is a recent history of trauma or developing facial asymmetry, or when hard-tissue grinding or crepitus is detected.²² Imaging should also be considered in patients that have failed to respond to conservative TMD treatment.³² Radiographic assessment may include:

- Panoramic or full mouth periapical films;
- Lateral cephalogram;
- TMJ tomography;
- Magnetic resonance imaging (both open and closed mouth to view disc position); or
- Cone-beam computed tomography (CBCT).

TMJ arthrography is not recommended as a routine diagnostic procedure.⁶³⁻⁶⁵ The readily available panoramic radiograph is reliable for evaluating condylar head morphology and angulation but does not permit evaluation of the joint space, soft tissues, or condylar motion.²² The panograph may indicate osseous changes, but negative findings do not rule out TMJ pathology.⁶⁶ The CBCT can be used to detect bony abnormalities and fractures and to assess asymmetry,⁶⁴⁻⁶⁶ but generates a much higher radiation burden than the panoramic image. Magnetic resonance imaging provides visualization of soft tissues, specifically the position and contours of the TMJ disc, and can be used to detect inflammation.^{22,65}

TMD has been divided into two broad categories, TMJ disorders and masticatory muscles disorders,¹ which are listed below.

- I. TMJ disorders:
 - a. Joint pain: arthralgia (synovitis, capsulitis, and retrodiscitis)
 - b. Joint disorders:
 1. Disc-condyle complex disorders (disc displacement with reduction, disc displacement without reduction).
 2. Hypomobility disorders (intra-articular fibrous adhesions, ankylosis).
 3. Hypermobility disorders (subluxation, luxation).

- c. Joint diseases:
 1. Osteoarthritis also known as degenerative joint disease, condylitis, osteochondritis dissecans, osteonecrosis.
 2. Systemic arthritides such as rheumatoid arthritis, idiopathic juvenile arthritis, spondyloarthropathies, psoriatic arthritis, infections arthritis, Reiter syndrome, and crystal induced disease.
 3. Neoplasms.
 4. Fractures (open and closed condylar and subcondylar).
- II. Masticatory muscle disorders:
 - a. Muscle pain limited to orofacial region (myalgia, myofascial pain, tendonitis, myositis, spasm).
 - b. Muscle pain due to systemic/central disorders (centrally mediated myalgia, fibromyalgia).
 - c. Movement disorders (dyskinesia, dystonia).
 - d. Other muscle disorders (contracture, hypertrophy, neoplasm).

Treatment of TMD

The goals of TMD treatment include restoral of function, decreased pain, and return of quality of life. Few studies document success or failure of specific treatment modalities for TMD in infants, children, and adolescents on a long-term basis. It has been suggested that simple, conservative, and reversible types of therapy are effective in reducing most TMD symptoms in children.⁶⁷ The focus of treatment should be to find a balance between active and passive treatment modalities. Active modalities include participation of the patient whereas passive modalities may include wearing a stabilization splint. In a randomized trial, adolescents undergoing occlusal appliance therapy combined with information attained a clinically significant improvement on the pain index.⁶⁸ Combined approaches may be more successful in treating TMD than single treatment modalities.⁶⁹

Treatment of TMD can be divided into reversible and irreversible treatment. Reversible therapies may include:

- Patient education (e.g., relaxation training, developing behavior coping strategies, modifying inadequate perceptions about TMD, patient awareness of clenching and bruxing habits, if present).⁶⁹
- Physical therapy [e.g., jaw exercises or transcutaneous electrical nerve stimulation (TENS), ultrasound, iontophoresis, massage, thermotherapy, coolant therapy)].^{32,70,71}
- Behavioral therapy (e.g., avoiding excessive chewing of hard foods or gum, voluntary avoidance of stressors, habit reversal; decreasing stress, anxiety, and/or depression obtaining adequate, uninterrupted sleep).^{32,71}
- Prescription medication (e.g., non-steroidal anti-inflammatory drugs, anxiolytic agents, muscle relaxers). While antidepressants have proved to be beneficial, they should be prescribed by a practitioner familiar with pain management.^{32,72}
- Occlusal splints. The goal of an occlusal appliance is to provide orthopedic stability to the TMJ. These alter

the patient's occlusion temporarily and may be used to decrease parafunctional activity.^{68,73,74} Occlusal splints may be made of hard or soft acrylic. The stabilization type of splint covers all of the teeth on either the maxillary or mandibular arch and is balanced so that all teeth are in occlusion when the patient is closed and the jaw is in a musculoskeletally stable position.^{24,32}

Irreversible therapies can include:

- Occlusal adjustment (i.e., permanently altering the occlusion or mandibular position by selective grinding or full mouth restorative dentistry).⁷⁵
- Orthodontics. This may include mandibular positioning devices designed to alter the growth or permanently reposition the mandible (e.g., headgear, functional appliances). There is little evidence that orthodontic treatment can prevent or relieve TMD.^{76,77}
- Botulinum toxin A injections. Although recently approved for use in adults to provide masticatory muscle relaxation, this modality has not been approved for use in children.³²

Controversy surrounds the significance of signs and symptoms in this age group, the value of certain diagnostic procedures, and what constitutes appropriate therapy. It is not clear whether these signs and symptoms constitute normal variation, preclinical features, or manifestations of a disease state. Whether these signs and symptoms warrant treatment as predictors of TMD in adulthood is questionable.³⁸

Referral should be made to other health care providers, including those with expertise in TMD, oral surgery, or pain management, when the diagnostic and/or treatment needs are beyond the treating dentist's scope of practice.

Recommendations

Every comprehensive dental history and examination should include a TMJ history and assessment. The history should include questions concerning the presence of head and neck pain and mandibular dysfunction, previous orofacial trauma, and history of present illness with an account of current symptoms. In the presence of a positive history and/or signs and symptoms of TMD, a more comprehensive examination (e.g., palpation of masticatory and associated muscles and the TMJ's, documentation of joint sounds, occlusal analysis, and assessment of range of mandibular movements including maximum opening, protrusion, and lateral excursions) should be performed. A referral may be considered. Joint imaging may be recommended by other specialists to investigate joint sounds in the absence of other TMD signs and symptoms.

Therapeutic modalities to prevent TMD in the pediatric population are yet to be supported by controlled studies. For children and adolescents with signs and symptoms of TMD, reversible therapies should be considered. Because of inadequate data regarding their usefulness, irreversible therapies should be avoided.^{68,75,77} Referral to a medical specialist may be indicated when primary headaches, otitis media, allergies,

abnormal posture, airway congestion, rheumatoid arthritis, connective tissue disease, psychiatric disorders, or other medical conditions are suspected.

References

1. American Academy of Orofacial Pain. Orofacial Pain: Guidelines for assessment, diagnosis, and management. deLeeuw R, Klasser GD eds. 5th edition. Chicago, IL: Quintessence Publishing; 2013:127-186.
2. Okeson J. Etiology of functional disturbances in the masticatory system. In: Management of Temporomandibular Disorders and Occlusion. 7th ed. St Louis, Mo: Elsevier Mosby Inc; 2013:102-69.
3. Stohler CS. Clinical perspectives on masticatory and related muscle disorders. In: Sessle BJ, Bryant PS, Dionne RA, eds. Temporomandibular Disorders and Related Pain Conditions. Vol 4. Seattle, Wash: IASP Press; 1995:3-30.
4. Kopp S. Degenerative and inflammatory temporomandibular joint disorders. In: Sessle BJ, Bryant PS, Dionne RA, eds. Temporomandibular Disorders and Related Pain Conditions. Vol 4. Seattle, Wash: IASP Press; 1995: 119-32.
5. Dolwich MF. Temporomandibular joint disk displacement. In: Sessle BJ, Bryant PS, Dionne RA, eds. Temporomandibular Disorders and Related Pain Conditions. Volume 4. Seattle, Wash: IASP Press; 1995:79-113.
6. Alamoudi N, Farsi N, Salako N, Feteih R. Temporomandibular disorders among school children. *J Clin Pediatr Dent* 1998;22(4):323-9.
7. List T, Wahlund K, Wenneberg B, Dworkin SF. TMD in children and adolescents: Prevalence of pain, gender differences, and perceived treatment need. *J Orofac Pain* 1999;13(1):9-20.
8. Stockstill JW, Bowley JF, Dunning D, Spalding P, Stafford K, Erickson L. Prevalence of temporomandibular disorders in children based on physical signs. *J Dent Child* 1998;65(6):459-67.
9. Paesani D, Salas E, Martinez A, Isberg A. Prevalence of temporomandibular joint disk displacement in infants and young children. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999;87(1):15-9.
10. National Institute of Dental and Craniofacial Research. Facial Pain. Available at: "<http://www.nidcr.nih.gov/DataStatistics/FindDataByTopic/FacialPain/>". Accessed August 20, 2015.
11. Manfredini D, Guarda-Nardini L, Winocur E, Piccotti F, Ahlberg J, Lobbezoo F. Research diagnostic criteria for temporomandibular disorders: A systematic review of axis I epidemiologic findings. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112(4):453-62.
12. Nydell A, Helkimo M, Koch G. Craniomandibular disorder in children: A critical review of the literature. *Swed Dent J* 1994;18:191-205.
13. Köhler AA, Helkimo AN, Magnusson T, Hugoson A. Prevalence of symptoms and signs indicative of temporomandibular disorders in children and adolescents. A cross-sectional epidemiological investigation covering two decades. *Euro Arch Paed Dent* 2009;10(1):16-25.
14. Nilsson IM. Reliability, validity, incidence and impact of temporomandibular pain disorders in adolescents. *Swed Dent J Suppl* 2007;(183):7-86.
15. Bonjardim LR, Gavião MB, Carmagnani FG, Pereira LF, Castelo PM. Signs and symptoms of temporomandibular joint dysfunction in children with primary dentition. *J Clin Pediatr Dent* 2003;28(1):53-8.
16. Thilander B, Rubio G, Pena L, De Mayorga C. Prevalence of temporomandibular dysfunction and its association with malocclusion in children and adolescents: An epidemiologic study related to specified stages of dental development. *Angle Orthod* 2002;72(2):146-54.
17. Nilsson IM, List T, Drangsholt M. Headache and comorbid pains associated with TMD pain in adolescents. *J Dent Res* 2013;92(9):802-7.
18. Nilsson IM, Drangsholt M, List T. Impact of temporomandibular disorder pain in adolescents: differences by age and gender. *J Orofac Pain* 2009;23(2):115-22.
19. LeResche L, Mancl LA, Drangsholt MT, Huang G, von Korff MV. Predictors of onset of facial pain and temporomandibular disorders in early adolescence. *Pain* 2007;129(3):269-78.
20. LeResche L, Mancl LA, Drangsholt MT, Saunders K, Von Korff M. Relationship of pain and symptoms to pubertal development in adolescents. *Pain* 2005;118(1-2):201-9.
21. Greene CS. The etiology of temporomandibular disorders: Implications for treatment. *J Orofac Pain* 2001; 15(2):93-105.
22. Howard JA. Temporomandibular joint disorders in children. *Dent Clin N Am* 2013;57(1):99-127.
23. Hiatt JL, Gartner LP. Temporomandibular joint. In: Textbook of Head and Neck Anatomy. 2nd ed. Baltimore, Md; Williams and Wilkins; 1987:223.
24. Okeson JP. Temporomandibular joint pains. In Bell's Oral and Facial Pain, 7th edition. Chicago, IL; Quintessence Publishing; 2014:327-69.
25. Greco CM, Rudy TE, Turk DC, Herlich A, Zaki HH. Traumatic onset of temporomandibular disorders: Positive effects of a standardized conservative treatment program. *Clin J Pain* 1997;13(4):337-47.
26. Fischer DJ, Mueller BA, Critchlow CW, LeResche L. The association of temporomandibular disorder pain with history of head and neck injury in adolescents. *J Orofac Pain* 2006;20(3):191-8.
27. Imahara SD, Hopper RA, Wang J, Rivara FP, Klein MB. Patterns and outcomes of pediatric facial fractures in the United States: A survey of the National Trauma Data Bank. *J Am Coll Surg* 2008;207(5):710-6.
28. Akhter R, Hassan NM, Ohkubo R, et al. The relationship between jaw injury, third molar removal, and orthodontic treatment and TMD symptoms in university students in Japan. *J Orofac Pain* 2008;22(1):50-6.

29. Leuin SC, Frydendall E, Gao D, Chan KH. Temporomandibular joint dysfunction after mandibular fracture in children: A 10 year review. *Arch Otolaryngol Head Neck Surg* 2011;137(1):10-14.
30. Kaban L. Acquired abnormalities of the temporomandibular joint. In: Kaban LB, Troulis MJ, eds. *Pediatric Oral and Maxillofacial Surgery*. Philadelphia, Pa; WB Saunders; 2004:340-76.
31. Güven O. A clinical study on temporomandibular joint ankylosis in children. *J Craniofac Surg* 2008;19(5):1263-9.
32. Dym H, Israel H. Diagnosis and treatment of temporomandibular disorders. *Dent Clin N Am* 2012;56:149-61.
33. American Academy of Sleep Medicine. Sleep related bruxism. In *International Classification of Sleep Disorders. Diagnosis and Coding Manual*, 3rd ed. Westchester, IL: American Academy of Sleep Medicine, 2014:182-5.
34. Cheifetz AT, Osganian SK, Allred EN, Nleedleman HL. Prevalence of bruxism and associated correlates in children as reported by parents. *J Dent Child (Chic)* 2005;72(2):67-73.
35. Barbosa Tde S, Miyakoda LS, Pocztaruk Rde L, Rocha CP, Gavião MBD. Temporomandibular disorders and bruxism in childhood and adolescence: Review of the literature. *Int J Pediatr Otorhinolaryngol* 2008;72(3):299-314.
36. Castelo PM, Gavião MB, Pereira LJ, Bonjardim LR, Gavião MBD. Relationship between oral parafunctional/nutritive sucking habits and temporomandibular joint dysfunction in primary dentition. *Int J Paediatr Dent* 2005;15(1):29-36.
37. Winocur E, Gavish A, Finkelshtein T, Halachmi M, Gazit E. Oral habits among adolescent girls and their association with symptoms of temporomandibular disorders. *J Oral Rehabil* 2001;28(7):624-9.
38. Carlsson GE, Egermark I, Magnusson T. Predictors of signs and symptoms of temporomandibular disorders: A 20-year follow-up study from childhood to adulthood. *Acta Odontol Scand* 2002;60(3):180-5.
39. Magnusson T, Egermark I, Carlsson GE. A prospective investigation over two decades on signs and symptoms of temporomandibular disorders and associated variables. A final summary. *Acta Odontol Scand* 2005;63(2):99-109.
40. Gesch D, Bernhardt O, Mack F, John U, Kocher T, Dietrich A. Association of malocclusion and functional occlusion with subjective symptoms of TMD in adults: Results of the Study of Health in Pomerania (SHIP). *Angle Orthod* 2005;75(2):183-90.
41. Alamoudi N. Correlation between oral parafunction and temporomandibular disorders and emotional status among Saudi children. *J Clin Pediatr Dent* 2001;26(1):71-80.
42. De Boever JA, Carlsson GE, Klineberg IJ. Need for occlusal therapy and prosthodontic treatment in the management of temporomandibular disorders. Part I. Occlusal interference and occlusal adjustment. *J Oral Rehabil* 2000;27(5):367-79.
43. Taskaya-Yilmaz N, Ögütçen-Toller M, Saraç YŞ. Relationship between the TMJ disc and condyle position on MRI and occlusal contacts on lateral excursions in TMD patients. *J Oral Rehab* 2004;31(8):754-8.
44. Henrikson T, Nilner M. Temporomandibular disorders, occlusion and orthodontic treatment. *J Orthod* 2003;30(2):129-37; discussion 127.
45. Egermark I, Carlsson GE, Magnusson T. A prospective long-term study of signs and symptoms of temporomandibular disorders in patients who received orthodontic treatment in childhood. *Angle Orthod* 2005;75(4):645-50.
46. Henrikson T, Nilner M, Kurol J. Symptoms and signs of temporomandibular disorders before, during and after orthodontic treatment. *Swed Dent J* 1999;23(5-6):193-207.
47. Henrikson T, Nilner M, Kurol J. Signs of temporomandibular disorders in girls receiving orthodontic treatment. A prospective and longitudinal comparison with untreated Class II malocclusions and normal occlusion subjects. *Eur J Orthod* 2000;22(3):271-81.
48. Kim MR, Graber TM, Viana MA. Orthodontics and temporomandibular disorder: A meta-analysis. *Am J Orthod Dentofac Orthop* 2002;121(5):438-46.
49. Phillips JT. What skeletal and dental characteristics do TMD patients have in common? *Funct Orthod* 2007;24(1):24-6, 28, 30.
50. Pahkala R, Qvarnström M. Can temporomandibular dysfunction signs be predicted by early morphological or functional variables? *Euro J Orthod* 2004;26(4):367-73.
51. List T, Wahlund K, Larsson B. Psychosocial functioning and dental factors in adolescents with temporomandibular disorders: a case-control study. *J Orofac Pain* 2001;15(3):218-27.
52. American Academy of Orofacial Pain. Axis II: Biobehavioral considerations. In: de Leeuw R, Klasser GD, eds. *Orofacial Pain: Guidelines for assessment, diagnosis, and management*. 5th ed. Chicago, IL: Quintessence Publishing; 2013:392-426.
53. Karibe H, Goddard G, Aoyagi K, et al. Comparison of subjective symptoms of temporomandibular disorders in young patients by age and gender. *Cranio* 2012;30(2):114-20.
54. Milam SB, Zardeneta G, Schmitz JP. Oxidative stress and degenerative temporomandibular joint disease: A proposed hypothesis. *J Oral Maxillofac Surg* 1998;56:214-33.
55. Buckinham RB, Braun T, Harinstein DA, et al. Temporomandibular joint dysfunction syndrome: A close association with systemic joint laxity (the hypermobile joint syndrome). *Oral Surg Oral Med Oral Pathol* 1991;72:514-9.
56. Fillingim R, Slade G, Diatchenko L, et al. Summary of findings from the OPPERA baseline case-control study: Implications and future directions. *J Pain* 2011;12(11 supplement):T102-T107.

57. LeResche L, Mancini L, Sherman JJ, Gandara B, Dworkin SF. Changes in temporomandibular pain and other symptoms across the menstrual cycle. *Pain* 2003;106(3):253-61.
58. LeResche L, Saunders K, Von Korff MR, Barlow W, Dworkin SF. Use of exogenous hormone and risk of temporomandibular pain. *Pain* 1997;69:153-60.
59. McDonald RE, Avery DR, Dean JA. Examination of the mouth and other relevant structures. In: Dean JA, Avery DR, McDonald RE, eds. *McDonald and Avery's Dentistry for the Child and Adolescent*. 9th ed. Maryland Heights Mo: Mosby Elsevier; 2011:1-18.
60. Casamassimo PS, Christensen JR, Fields HW Jr, Ganzberg S. Examination, diagnosis, and treatment planning for general and orthodontic problems. In: Casamassimo PS, Fields HW Jr, McTigue DJ, Nowak AJ, eds. *Pediatric Dentistry: Infancy through Adolescence*. 5th ed. St. Louis, Mo: Elsevier Saunders; 2013:580-5.
61. American Academy of Pediatric Dentistry. Guideline on record keeping. *Pediatr Dent* 2015;36(special issue):302-9.
62. American Academy of Orofacial Pain. General assessment of the orofacial pain patient. In: de Leeuw R, Klasser GD, eds. *Orofacial Pain: Guidelines for Assessment, Diagnosis, and Management*. Carol Stream, Ill: Quintessence Publishing Co Inc; 2013:25-47.
63. Loos PJ, Aaron GA. Standards for management of the pediatric patient with acute pain in the temporomandibular joint or muscles of mastication. *Pediatr Dent* 1989;11(4):331.
64. Brooks SL, Brand JW, Gibbs SJ, et al. Imaging of the temporomandibular joint: A position paper of the American Academy of Oral and Maxillofacial Radiology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1997;83(5):609-18.
65. DeSenna BR, dos Santos S, Franca JP, Marques LS, Pereira LJ. Imaging diagnosis of the temporomandibular joint: Critical review of indications and new perspectives. *Oral Radiol* 2009;25(2):86-98.
66. Hunter A, Kalathingal S. Diagnostic imaging for temporomandibular disorders and orofacial pain. *Dent Clin N Am* 2013;57:405-18.
67. Bodner L, Miller VJ. Temporomandibular joint dysfunction in children: Evaluation of treatment. *Int J Pediatr Otorhinolaryngol* 1998;44(2):133-7.
68. Wahlund K, List T, Larsson B. Treatment of temporomandibular disorders among adolescents: A comparison between occlusal appliance, relaxation training, and brief information. *Acta Odontol Scand* 2003;61(4):203-11.
69. Michelotti A, Steenks MH, Farella M, Parisini F, Cimino R, Martina R. The additional value of a home physical therapy regimen versus patient education only for the treatment of myofascial pain of the jaw muscles: Short-term results of a randomized clinical trial. *J Orofac Pain* 2004;18(2):114-25. [Erratum in *J Orofac Pain* 2006;20(2):106].
70. Medlicott, MS, Harris SR. A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder. *Phys Ther* 2006;86(7):955-73.
71. Crider AB, Glaros AG. A meta-analysis of EMG biofeedback treatment of temporomandibular disorders. *J Orofac Pain* 1999;13(1):29-37.
72. List T, Axelsson S, Leijon G. Pharmacologic interventions in the treatment of temporomandibular disorders, atypical facial pain, and burning mouth syndrome. A qualitative systematic review. *J Orofac Pain* 2003;17(4):301-10.
73. Simmons HC III, Gibbs SJ. Anterior repositioning appliance therapy for TMJ disorders: Specific symptoms relieved and relationship to disk status on MRI. *Cranio* 2005;23(2):89-99.
74. Fujii T, Torisu T, Nakamura S. A change of occlusal conditions after splint therapy for bruxers with and without pain in the masticatory muscles. *Cranio* 2005;23(2):113-8.
75. Koh H, Robinson PG. Occlusal adjustment for treating and preventing temporomandibular joint disorders. *Cochrane Database Syst Rev* 2003;(1):CD003812.
76. Luther F, Layton S, McDonald F. Orthodontics for treating temporomandibular joint (TMJ) disorders. *Cochrane Database Syst Rev* 2010(7):CD006541.
77. Rey D, Oberti G, Baccetti T. Evaluation of temporomandibular disorders in Class III patients treated with mandibular cervical headgear and fixed appliances. *Am J Orthod Dentofac Orthop* 2008;133(3):379-81.