

Policy on Third-party Reimbursement of Fees Related to Dental Sealants

Originating Committee

Clinical Affairs Committee

Review Council

Council on Clinical Affairs

Adopted

1999

Revised

2000, 2006

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that the placement of sealants and their continued maintenance are scientifically-sound and cost-effective techniques for prevention of pit and fissure caries.

Methods

This revision is based upon a review of current dental literature related to dental sealants. A MEDLINE search was conducted using the terms “dental sealants”, “indications”, and “insurance”.

Background

According to national estimates, by 17 years of age, 78% of children in the United States have experienced caries.¹ As much as 90% of all caries in school-aged children occurs in pits and fissures. The teeth at highest risk by far are permanent first and second molars where fluoride has its least preventive effect on the pits and fissures. Any tooth, including primary teeth and permanent teeth other than molars, may benefit from sealant application due to fissure anatomy and caries risk factors.^{2,3} Caries risk may increase due to changes in patient habits, oral microflora, or physical condition, and unsealed teeth subsequently might benefit from sealant application.^{2,3}

Current data also show that, although initial sealant retention rates are high, sealant loss does occur.² It is in the patient's interest to receive periodic evaluation of sealants for maintenance or replacement. Without recall and maintenance, sealant failure will compound over time, leaving previously sealed surfaces with a caries susceptibility equal to that of surfaces that never were sealed. With appropriate followup care, the success rate of sealants may be 80 to 90%, even after a decade.²

Although sealants are safe and effective, their use continues to be low.⁴ Sealants are particularly effective in preventing pit and fissure caries and providing cost savings if placed on patients during periods of greatest risk.^{5,6} However, initial insurance coverage for sealants often is denied, and insurance coverage for repair and/or replacement may be limited.⁷

Recommendations

1. The dentition should be evaluated periodically for developmental defects and deep pits and fissures that may contribute to caries risk. Dental sealants should be placed on susceptible teeth and should be evaluated for repair or replacement as part of a periodic dental examination.
2. Third party coverage for sealants should not be based upon a patient's age. Timing of the eruption of teeth can vary widely. Furthermore, caries risk may increase at any time during a patient's life.
3. The AAPD shall work with other dental organizations, the insurance industry, and consumer groups to make the advantages of dental sealants understood and to seek reimbursement for fees associated with their placement, maintenance, and repair.

References

1. US Dept of Health and Human Services. Oral Health in America: A Report of the Surgeon General. Rockville, Md: US Dept of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000.
2. Feigal RJ. The use of pit and fissure sealants. *Pediatr Dent* 2002;24(5):415-22.
3. American Academy of Pediatric Dentistry. Guideline on pediatric restorative dentistry. *Pediatr Dent* 2005;27(suppl):122-9.
4. US Dept of Health and Human Services. Healthy people 2010. Rockville, Md: US Dept of Health and Human Services, National Institutes of Health; 2000.
5. Weintraub JA. Pit and fissure sealants in high-caries risk individuals. *J Dent Educ* 2001;65(10):1084-90.
6. American Dental Association Council on Access, Prevention, and Interprofessional Relations, American Dental Association Council on Scientific Affairs. Dental sealants. *J Am Dent Assoc* 1997;128(4):485-8.
7. American Dental Association. Statement on preventive coverage in dental benefits plans. Chicago, Ill; 1992:602; 1994:656.