

Policy on Use of Xylitol in Pediatric Dentistry

Latest Revision

2024

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that there is considerable research on sugar substitutes, particularly xylitol, and their potential oral health benefits for infants, children, adolescents, and persons with special health care needs. The AAPD encourages practitioners to follow evolving literature on the effectiveness and limitations of sugar substitutes in preventing caries.

Methods

This policy was developed by the Council on Clinical Affairs, adopted in 2006¹, and last revised in 2020². For this revision, literature searches of the PubMed®/MEDLINE and Google Scholar databases were conducted using the terms: xylitol AND dental, systematic review; field: all fields; limits: within the last five years, humans, English, birth through 18. One hundred seventy-four articles matched these criteria; 17 systematic reviews and/or meta-analyses were reviewed for this revision. When data did not appear sufficient or were inconclusive, policy was based upon expert and/or consensus opinion by experienced researchers and clinicians.

Background

Xylitol is a five-carbon sugar alcohol derived from forest and agricultural materials.^{3,4} In 1963, the United States Food and Drug Administration (FDA) approved xylitol as a nutritional additive.^{4,6} Dental benefits of xylitol first were suggested from Finnish studies using animal models in 1970.⁷ The first xylitol studies in humans, the Turku Sugar Studies,^{8,9} demonstrated the relationship between dental plaque and xylitol, as well as the safety of xylitol for human consumption. Xylitol as well as other sugar alcohols are not readily metabolized by oral bacteria and, thus, are considered noncariogenic sugar substitutes.⁶ It is hypothesized that xylitol reduces Mutans streptococci (MS) counts and lactic acid production by bacteria in plaque and saliva with minimal effect on normal flora.^{4,10,11} Some evidence shows a 30 to 80 percent decrease in caries incidence with consumption of xylitol (five to 10 grams three times per day).¹²⁻¹⁴ However, frequencies less than three times a day (less than 3.44 grams per day [g/day]) yielded no protective effect.^{4,15} Literature, systematic reviews, and meta-analyses show mixed results.^{4,16-18}

Evidence regarding the frequency of use, duration of exposure, and age for introduction of xylitol in preventing early childhood caries is inconsistent.¹⁷ The impact of xylitol on preventing caries by interrupting the transfer of cariogenic

How to Cite: American Academy of Pediatric Dentistry. Policy on use of xylitol in pediatric dentistry. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2024:114-6.

microorganisms from mother to child is also inconclusive.^{8,9,19-27} These studies have been performed with xylitol intake ranging from four to 15 g/day divided into three to seven consumption periods²⁰⁻²², with use at higher daily doses yielding significantly better results.^{4,15}

Xylitol is available in many nutraceutical products including gums, gummies, candies, mints, chewable tablets, lozenges, toothpastes, mouthwashes, cough mixtures, and oral wipes.⁴ Some studies report the chewing process stimulates saliva production which enhances the caries inhibitory effect.^{15,17} This can be a significant confounding variable when evaluating the efficacy of xylitol gum as a primary prevention agent.^{15,19-25}

The safety profile of xylitol is well documented.^{4,28} Studies show few adverse events with xylitol when administered at a recommended dose of six g/day.^{4,28} Abdominal distress and osmotic diarrhea are common side effects with higher doses.^{15,20-27,29,30} Xylitol also can cause life-threatening toxicoses in dogs.³¹

Xylitol products are branded as a preventive measure for dental caries.^{4,15} However, the evidence supporting xylitol's role in the primary prevention of dental caries is mixed, suggesting insufficient evidence to show xylitol products reduce caries rates on their own.^{15,20-27,29,32} Design flaws and/or bias (e.g., insufficient sample size, inconsistent results, randomization, blinding, conflict of interest) limit the significance and applicability of some xylitol studies.^{15,20-27,29,30} In addition, optimal results for primary prevention require consistent use of 100 percent xylitol, chewed or consumed three to five times per day after meals with a total dose of five to 10 g/day, which may be unrealistic in clinical practice.^{4,15,23,24}

In light of current evidence, the true value of xylitol products lies as sugar substitutes, as they are noncariogenic natural alternatives.³³⁻³⁵ In 2015, the World Health Organization recommended an ideal reduction of the intake of free sugars to below five percent of total energy intake to reduce the burden of overweight, obesity, and related diseases like dental caries.^{36,37} In 2023, the FDI World Dental Federation (FDI) published a position paper on free sugars that called for policies and guidelines to reduce global sugar consumption, delivering the human resources for health, promoting industry

ABBREVIATIONS

AAPD: American Academy of Pediatric Dentistry. **G/day:** grams per day. **MS:** Mutans streptococci.

accountability, and promoting healthy schools, hospitals, and workplaces.³⁸ It specifically suggested that dental associations should advocate for and support integrated strategies to reduce free sugars consumption addressing equitable access to appropriate oral healthcare, oral health literacy, health promotion, policy implementation, health surveillance, and monitoring.³⁸ Judicious use of natural sugar substitutes could be a feasible strategy for reducing sugar consumption as part of a comprehensive caries management plan.^{38,39}

Policy statement

The AAPD:

- supports the use of xylitol and other sugar alcohols as noncariogenic sugar substitutes.
- acknowledges the current lack of consistent evidence supporting xylitol as a primary caries-preventive agent, specifically in demonstrating significant reductions in MS among children.
- recognizes that the large dose and high frequency of xylitol used in clinical trials may be unrealistic in clinical practice.
- supports further research to clarify the impact of xylitol delivery vehicles, the frequency of exposure, and the optimal dosage to reduce caries and improve the oral health of children.

References

1. American Academy of Pediatric Dentistry. Use of xylitol in caries prevention. *Pediatr Dent* 2006;28(suppl):31-2.
2. American Academy of Pediatric Dentistry. Use of xylitol in pediatric dentistry. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill.: American Academy of Pediatric Dentistry; 2020:113-4.
3. Mäkinen KK. Biochemical principles of the use of xylitol in medicine and nutrition with special consideration of dental aspects. *Experientia Suppl* 1978;30:1-160.
4. AlHumaid J, Bamashmous M. Meta-analysis on the effectiveness of xylitol in caries prevention. *J Int Soc Prev Community Dent* 2022;12(2):133-8.
5. Drugs.com. Xylitol. Last updated April 1, 2022. Available at: "<https://www.drugs.com/npp/xylitol.html>". Accessed March 11, 2024.
6. Milgrom P, Rothen M, Milgrom L. Developing public health interventions with xylitol for the US and US-associated territories and states. *Suom Hammaslaakarilehti* 2006;13(10-11):2-11.
7. Muhlemann HR, Regolati B, Marthaler TM. The effect on rat fissure caries of xylitol and sorbitol. *Helv Odontol Acta* 1970;14(1):48-50.
8. Scheinin A, Mäkinen KK, Tammissalo E, Rekola M. Turku sugar studies. XVIII. Incidence of dental caries in relation to 1-year consumption of xylitol chewing gum. *Acta Odontol Scand* 1975;33(5):269-78.
9. Scheinin A, Mäkinen KK, Ylitalo K. Turku sugar studies. V. Final report on the effect of sucrose, fructose and xylitol diets on caries incidence in man. *Acta Odontol Scand* 1976;34(4):179-216.
10. Bastos JR. Utilization of xylitol as a preventive substance in dentistry. *Braz J Oral Sci* 2005;4:891-3.
11. Stecksén-Blicks C, Holgerson PL, Twetman S. Effect of xylitol and xylitol-fluoride lozenges on approximal caries development in high-caries-risk children. *Int J Paediatr Dent* 2008;18(3):170-7.
12. Hujoel PP, Mäkinen KK, Bennett CA, et al. The optimum time to initiate habitual xylitol gum-chewing for obtaining long-term caries prevention. *J Dent Res* 1999;78(3):797-803.
13. Mäkinen KK, Bennett CA, Hujoel PP, et al. Xylitol chewing gums and caries rates: A 40-month cohort study. *J Dent Res* 1995;74(12):1904-13.
14. Söderling E, Isokangas P, Pienihäkkinen K, Tenovuo J. Influence of maternal xylitol consumption on acquisition of mutans streptococci by infants. *J Dent Res* 2000;79(3):882-7.
15. Marghalani AA, Guinto E, Phan M, Dhar V, Tinanoff N. Effectiveness of xylitol in reducing dental caries in children. *Pediatr Dent* 2017;39(2):103-10.
16. Maguire A, Rugg-Gunn AJ. Xylitol and caries prevention — Is it a magic bullet? *Br Dent J* 2003;194(8):429-36. Available at: "<https://www.nature.com/articles/4810022>". Accessed February 27, 2024.
17. Söderling E, Pienihäkkinen K. Effects of xylitol chewing gum and candies on the accumulation of dental plaque: A systematic review. *Clin Oral Investig* 2022;26(1):119-29.
18. van Loveren C. Sugar alcohols: What is the evidence for caries-preventive and caries-therapeutic effects? *Caries Res* 2004;38(3):286-93. Available at: "<https://doi.org/10.1159/000077768>". Accessed February 27, 2024.
19. Nakai Y, Shinga-Ishihara C, Kaji M, Moriya K, Murakami-Yamanaka K, Takimura M. Xylitol gum and maternal transmission of mutans streptococci. *J Dent Res* 2010;89(1):56-60.
20. Antonio AG, Pierro VS, Maia LC. Caries preventive effects of xylitol-based candies and lozenges: A systematic review. *J Public Health Dent* 2011;71(2):117-24.
21. Lin HK, Fang CE, Huang MS, et al. Effect of maternal use of chewing gums containing xylitol on transmission of mutans streptococci in children: A meta-analysis of randomized controlled trials. *Int J Paediatr Dent* 2016;26(1):35-44.
22. Riley P, Moore D, Ahmed F, Sharif MO, Worthington H. Xylitol-containing products for preventing dental caries in children and adults. *Cochrane Database Syst Rev* 2015;(3):CD010743.

References continued on the next page.

23. Chou R, Pappas M, Dana T, et al. Screening and Interventions to Prevent Dental Caries in Children Younger Than Age Five Years: A Systematic Review for the U.S. Preventive Services Task Force [Internet]. Rockville (Md): Agency for Healthcare Research and Quality (US); December 2021. (Evidence Synthesis, No. 210.) Available at: "<https://www.ncbi.nlm.nih.gov/books/NBK575915/>". March 10, 2024.
24. Janakiram C, Deepan Kumar CV, Joseph J. Xylitol in preventing dental caries: A systematic review and meta-analyses. *J Nat Sci Biol Med* 2017;8(1):16-21.
25. Mickenautsch S, Yengopal V. Anticariogenic effect of xylitol versus fluoride – A quantitative systematic review of clinical trials. *Int Dent J* 2012;62(1):6-20.
26. Fontana M. Enhancing fluoride: Clinical human studies of alternatives or boosters for caries management. *Caries Res* 2016;50(Suppl 1):22-37.
27. Muthu MS, Ankita S, Renugalakshmi A, Richard K. Impact of pharmacological interventions in expectant mothers resulting in altered mutans streptococci levels in their children. *Pediatr Dent* 2015;37(5):422-8.
28. Mäkinen KK. Xylitol and oral health. *Adv Food Res* 1979;25:137-58.
29. Wang Y, Li J, Sun W, Li H, Cannon R, Mei L. Effect of non-fluoride agents on the prevention of dental caries in primary dentition: A systematic review. *PLoS One* 2017;12(8):e0182221.
30. Xiao J, Alkhers N, Kopycka-Kedzierawskial DT, et al. Prenatal oral health care and early childhood caries prevention: A systematic review and meta-analysis. *Caries Res* 2019;53(4):411-21.
31. Piscitelli CM, Dunayer EK, Aumann M. Xylitol toxicity in dogs. *Compend Contin Educ Vet* 2010;32(2):E1-4.
32. He S, Choong EKM, Duangthip D, Chu CH, Lo ECM. Clinical interventions with various agents to prevent early childhood caries: A systematic review with network meta-analysis. *Int J Paediatr Dent* 2023;33(5):507-20.
33. Lei P, Chen H, Ma J, et al. Research progress on extraction technology and biomedical function of natural sugar substitutes. *Front Nutr* 2022;9:952147.
34. Wölnerhanssen BK, Meyer-Gerspach AC, Beglinger C, Islam MS. Metabolic effects of the natural sweeteners xylitol and erythritol: A comprehensive review. *Crit Rev Food Sci Nutr* 2020;60(12):1986-98.
35. Zhan L. Rebalancing the caries microbiome dysbiosis: Targeted treatment and sugar alcohols. *Adv Dent Res* 2018;29(1):110-6.
36. Breda J, Jewell J, Keller A. The importance of the World Health Organization sugar guidelines for dental health and obesity prevention. *Caries Res* 2019;53(2):149-52. Available at: "<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6425811/>". Accessed February 27, 2024.
37. World Health Organization. Guideline: Sugar intake for adults and children. Geneva: World Health Organization; 2015. Available at: "<https://www.who.int/publications/i/item/9789241549028>". Accessed June 11, 2024.
38. FDI World Dental Federation. FDI Position on Free Sugars. Sydney, Australia. 2023. Available at: "https://www.fdiworlddental.org/sites/default/files/2023-10/EN_FDI_Position%20statement%20on%20free%20sugars.pdf". Accessed February 27, 2024.
39. Featherstone JDB, Crystal YO, Alston P, et al. Evidence-based caries management for all ages-practical guidelines. *Front Oral Health* 2021;2:657518.