

Guideline on Acquired Temporomandibular Disorders in Infants, Children, and Adolescents

Originating Committee

Clinical Affairs Committee – Temporomandibular Joint Problems in Children Subcommittee

Review Council

Council on Clinical Affairs

Adopted

1990

Revised

1999, 2002, 2006, 2010

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that disorders of the temporomandibular joint (TMJ) occasionally occur in infants, children, and adolescents. This guideline is intended to assist the practitioner in the recognition and diagnosis of temporomandibular disorders (TMD) and to identify possible treatment options. It is beyond the scope of this document to recommend the use of specific treatment modalities.

Methods

This document is an update of the previous document, revised in 2006. The update included an electronic search using the following parameters: Terms: “temporomandibular disorder”, “TMJ dysfunction”, “TMD AND adolescents”, “TMD AND gender differences”, “TMD AND occlusion”, “TMD AND treatment”; Fields: all fields; Limits: within the last 15 years; humans; English; clinical trials. The reviewers agreed upon the inclusion of 69 references to support this guideline. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

Background

Development of the TMJ

Function influences form as development and growth of the TMJ proceeds. The TMJ is comprised of 3 major components: the mandibular condyle, the mandibular fossa, and the associated connective tissue (including the articular disk).¹ The first evidence of development of the TMJ in humans is seen 8 weeks after conception.² During the first decade of life, the mandibular condyle becomes less vascularized and most of the major morphological changes are completed. During the second decade of life, there is continued but progressive slowing of growth. The shape of the mandibular condyle may change significantly during growth with approximately 5% of condyles undergoing radiographic changes in shape between

12 and 16 years of age.³ From adolescence to adulthood, the condyle changes to a form that is greater in width than length. Although the TMJ experiences active growth in the first 2 decades, it undergoes adaptive remodeling changes throughout life.

Definition of TMD

Temporomandibular disorder is a term adopted by the American Dental Association in 1983 to facilitate coordination of research and communication.⁴ While TMD has been defined as “functional disturbances of the masticatory system”⁵, others include masticatory muscle disorders⁶, degenerative and inflammatory TMJ disorders⁷, and TMJ disk displacements⁸ under the umbrella of TMD.

Certain medical conditions are reported to occasionally mimic TMD. Among them are trigeminal neuralgia, central nervous system lesions, odontogenic pain, sinus pain, otological pain, developmental abnormalities, neoplasias, parotid diseases, vascular diseases, myofascial pain, cervical muscle dysfunction, and Eagle’s syndrome. Other common medical conditions (eg, otitis media, allergies, airway congestion, rheumatoid arthritis) can cause symptoms similar to TMD.⁹

Etiology of TMD

Temporomandibular disorders have multiple etiological factors.¹⁰ Many studies show a poor correlation between any single etiological factor and resulting signs (findings identified by the dentist during the examination) and symptoms (findings reported by the child or parent).¹⁰ In fact, the TMJ and masticatory system is complex and, thus, requires a thorough understanding of the anatomy and physiology of the structural, vascular, and neurological components in order to manage TMD. Alterations in any one or a combination of teeth, periodontal ligament, the TMJ, or the muscles of mastication eventually can lead to TMD.¹¹ Research is insufficient to predict reliably which patient will or will not develop TMD. Etiologic factors suggested as contributing to the development of TMD are:

1. Trauma: This would include impact injuries such as trauma to the chin. A common occurrence in childhood because of falling, chin trauma is reported to be a factor in the development of TMD in pediatric patients.¹²⁻¹⁴ Unilateral and bilateral intracapsular or subcondylar fractures are the most common mandibular fractures in children.¹⁵ Closed reduction and prolonged immobilization can result in ankylosis.^{16,17}
2. Occlusal factors: There is a relatively low association of occlusal factors and the development of temporomandibular disorders.^{18,19} However, several features characterize malocclusions associated with TMD:
 - skeletal anterior open bite;²⁰
 - overjet greater than 6 to 7 mm;²⁰⁻²³
 - retrocuspal position (centric relation) to intercuspal position (centric occlusion) slides greater than 4 mm;²⁴
 - unilateral lingual cross bite;²⁰⁻²⁵
 - 5 or more missing posterior teeth;^{26,27}
 - Class III malocclusion.²⁸
3. Parafunctional habits (eg, bruxism, clenching, hyperextension, other repetitive habitual behavior): Bruxism is thought to contribute to the development of TMD by joint overloading that leads to cartilage breakdown, synovial fluid alterations, and other changes within the joint. These parafunctional habits may occur while the patient is asleep or awake. A study of 854 patients younger than 17 years old found the prevalence of bruxism to be 38%.²⁹ The literature on the association between parafunction and TMD in pediatric patients is contradictory.³⁰⁻³² However, childhood parafunction was found to be a predictor of the same parafunction 20 years later.³³ Other studies found correlations between reported bruxism and TMD³⁴ with a 3.4 odds ratio.³⁵ Children who grind their teeth were found to complain more often of pain and muscle tenderness when eating.³⁶
4. Posture: Craniocervical posture has been associated with occlusion and with dysfunction of the TMJ, including abnormalities of the mandibular fossa, condyle, ramus, and disc.³⁷⁻³⁹
5. Changes in “free-way” dimension of the rest position: Normally 2-4 mm, this may be impinged by occlusal changes, disease, muscle spasms, nervous tension, and/or restorative prosthetics.¹¹
6. Orthodontic treatment: Current literature does not support that the development of TMD is caused by orthodontic treatment,^{21,40-43} regardless of whether premolars were extracted prior to treatment.⁴⁴

Prevalence of TMD in children and adolescents

The reported prevalence of TMD in infants, children, and adolescents varies widely in the literature.⁴⁴⁻⁴⁷ Prevalence of signs and symptoms increases with age. One study of the primary dentition reported 34% of patients with signs and/or symptoms of TMD.⁴⁸ An epidemiological study of 4724 children aged 5-17 years reported 25% with symptoms. Clicking was

seen in 2.7% of children in the primary dentition and 10.1% in late mixed dentition and further increased to 16.6% in patients with permanent dentition.²⁰ A similar study in preschool children found TMJ sounds and clicking in 16.6% of patients.⁴⁹ A study of 217 adolescents found that over 20% had signs and/or symptoms of dysfunction, with TMJ sounds and tenderness in the lateral pterygoid muscle as the most common findings.⁵⁰ Clicking is seen more frequently than either locking or luxation and affects girls more than boys. In general, the prevalence of signs and symptoms of TMD is lower in children compared to adults and is even less the younger the child but increases with increasing age.²⁵ Recent surveys have indicated a significantly higher prevalence of symptoms and greater need for treatment in girls than boys⁴⁵ with the development of symptomatic TMD correlated with the onset of puberty in girls.^{51,52}

Controversy surrounds the significance of signs and symptoms in this age group, the value of certain diagnostic procedures, and what constitutes appropriate therapy. It is not clear whether these signs and symptoms constitute normal variation, preclinical features, or manifestations of a disease state. Whether these signs and symptoms warrant treatment as predictors of TMD in adulthood is questionable.³³

Diagnosing TMD

All comprehensive dental examinations should include a screening evaluation of the TMJ and surrounding area.⁵³⁻⁵⁵ Diagnosis of TMD is based upon a combination of historical information, clinical examination, and/or craniocervical and TMJ imaging.^{56,57} The findings are classified as symptoms and signs.⁵³

For a diagnosis of TMD, patients must have a history of facial pain combined with physical findings, supplemented by radiographic or imaging data when indicated.⁵⁸ A screening history, as part of the health history, may include questions such as:⁵⁹

- Do you have difficulty opening your mouth?
- Do you hear noises within your jaw joint?
- Do you have pain in or around your ears or your cheeks?
- Do you have pain when chewing?
- Do you have pain when opening your mouth wide or when yawning?
- Has your “bite” felt uncomfortable or unusual?
- Does your jaw ever “lock” or “go out”?
- Have you ever had an injury to your jaw, head, or neck? If so, when? How was it treated?
- Have you previously been treated for a temporomandibular disorder? If so, when? How was it treated?

Clinical and physical assessment of the TMJ may include:²³

- Manual palpation of the muscles and TMJ to evaluate for tenderness of intraoral and extraoral jaw muscles, neck muscles, and TMJ capsule;
- Evaluation of jaw movements including assessment of mandibular range of motion using a millimeter ruler (ie, maximum unassisted opening, maximum assisted

opening, maximum lateral excursion, maximum protrusive excursion) and mandibular opening pattern (ie, is it symmetrical?). Restricted mandibular opening with or without pain on mandibular movement may be interpreted as signs of TMJ internal derangement.⁶⁰

- Determination of TMJ sounds by palpation and auscultation with a stethoscope;
- Radiographs (panoramic, full mouth periapicals, lateral cephalometric), TMJ tomography, and magnetic resonance imaging to examine for TMJ pathology and/or dental pathology. TMJ arthrography is not recommended as a routine diagnostic procedure.^{4,59}

As some mental disorders can greatly influence a patient's pain experience, psychosocial factors related to temporomandibular symptoms should be considered; this would include mood disorders, anxiety disorders, musculoskeletal problems, migraine headaches, tension headaches, emotional factors, ulcers, colitis, occupational factors, and developmental/acquired craniofacial anomalies.¹¹

There is a need for improved classification of TMDs; however, they largely can be grouped into 3 classes:

1. Disorders of the muscles of mastication (including protective muscle splinting, muscle spasm, and muscle inflammation);
2. Disorders of the TMJ (including internal disk derangement, disk displacement with reduction accompanied by clicking, and anterior disk displacement without reduction seen as mechanical restriction or closed lock); and
3. Disorders in other related areas that may mimic TMD (eg, chronic mandibular hypomobility, inflammatory joint disorders such as juvenile rheumatoid arthritis, degenerative joint disease, extrinsic trauma such as fracture).⁴

Treatment of TMD

Few studies document success or failure of specific treatment modalities for TMD in infants, children, and adolescents on a long-term basis. These suggest that simple, conservative, and reversible types of therapy are effective in reducing most TMD symptoms in children.⁶¹ The focus of treatment should be to find a balance between active and passive treatment modalities. Active modalities include participation of the patient whereas passive modalities may include wearing a stabilization splint. The most common form of treatment of TMD in children was information combined with occlusal appliance therapy.⁶² It has been shown that combined approaches are more successful in treating TMD than single treatment modalities.^{62,63} Treatment of TMD can be divided into reversible and irreversible treatment.

Reversible therapies may include:

- Patient education (eg, relaxation training, developing behavior coping strategies, modifying inadequate perceptions about TMD, patient awareness of clenching and bruxing habits, if present).⁵⁹
- Physical therapy [eg, jaw exercises or transcutaneous elec-

trical nerve stimulation (TENS), ultrasound, iontophoresis, massage, thermotherapy, coolant therapy)].^{5,64,65}

- Behavioral therapy (eg, avoiding excessive chewing of hard foods or gum, voluntary avoidance of stressors, habit reversal, decreasing stress, anxiety, and/or depression).⁶⁶
- Prescription medication (eg, non-steroidal anti-inflammatory drugs, anxiolytic agents, muscle relaxers). While antidepressants have proved to be beneficial, they should be prescribed by a physician.⁶⁷
- Occlusal splints. The goal of an occlusal appliance is to provide orthopedic stability to the TMJ. These alter the patient's occlusion temporarily and may be used to decrease parafunctional activity.^{62,68,69}

Irreversible therapies can include:

- Occlusal adjustment (ie, permanently altering the occlusion or mandibular position by selective grinding or full mouth restorative dentistry);
- Mandibular repositioning [designed to alter the growth or permanently reposition the mandible (eg, headgear, functional appliances)];
- Orthodontics.

Referral should be made to other health care providers, including those with expertise in TMD, oral surgery, or pain management, when the diagnostic and/or treatment needs are beyond the treating dentist's scope of practice.

Recommendations

Every comprehensive dental history and examination should include a TMJ history and assessment. The history should include questions concerning the presence of head and neck pain and mandibular dysfunction, previous orofacial trauma, and history of present illness with an account of current symptoms. In the presence of a positive history and/or signs and symptoms of TMD, additional information is suggested and a referral may be considered. A more comprehensive examination should be performed and include palpation of masticatory and associated muscles and the TMJ's, documentation of joint sounds, occlusal analysis, and assessment of range of mandibular movements including maximum opening, protrusion, and lateral excursions.

Joint imaging may be recommended by other specialists to investigate joint sounds in the absence of other TMD signs and symptoms. For example, the presence of crepitus may indicate degenerative change that is not yet painful.

Therapeutic modalities to prevent TMD in the pediatric population are yet to be supported by controlled studies. For children and adolescents with signs and symptoms of TMD, reversible therapies should be considered. Because of inadequate data regarding their usefulness, irreversible therapies should be avoided.^{61,69} Referral to a medical specialist may be indicated when otitis media, allergies, abnormal posture, airway congestion, rheumatoid arthritis, or other medical conditions are suspected.⁵⁹

References

1. Carlson DS. Growth of the temporomandibular joint. In: Zarb GAC, Carlsson GE, Sessle BJ, Mohl ND, eds. *Temporomandibular Joint and Masticatory Muscle Disorders*. 2nd ed. Copenhagen: Munksgaard; 1994:128-50.
2. Dixon AD. Formation of the cranial base. In: Dixon AD, Howyte DAN, Rønning O, eds. *Fundamentals of Craniofacial Growth*. Boca Raton, FL: CRC press; 1997:100-29.
3. Dibbets JM, Van der Weele LT. Prevalence of structural bony change in the mandibular condyle. *J Craniomand Dis Fac Oral Pain* 1992;6(4):254-9.
4. Griffiths R. Report of the President's conference on the examination, diagnosis, and management of temporomandibular disorders and occlusion. *J Am Dent Assoc* 1983;106(1):75-7.
5. Okeson J. Etiology of functional disturbances in the masticatory system. In: *Management of Temporomandibular Disorders and Occlusion*. 6th ed. St Louis, Mo: Mosby Year Book Publication, Inc; 2008:130-63.
6. Stohler CS. Clinical perspectives on masticatory and related muscle disorders. In: Sessle BJ, Bryant PS, Dionne RA, eds. *Temporomandibular Disorders and Related Pain Conditions*. Vol 4. Seattle, Wash: IASP Press; 1995:3-30.
7. Kopp S. Degenerative and inflammatory temporomandibular joint disorders. In: Sessle BJ, Bryant PS, Dionne RA, eds. *Temporomandibular Disorders and Related Pain Conditions*. Vol 4. Seattle, Wash: IASP Press; 1995: 119-32.
8. Dolwich MF. Temporomandibular joint disk displacement. In: Sessle BJ, Bryant PS, Dionne RA, eds. *Temporomandibular Disorders and Related Pain Conditions*. Volume 4. Seattle, Wash: IASP Press; 1995:79-113.
9. Loos PJ, Aaron GA. Standards for management of the pediatric patient with acute pain in the temporomandibular joint or muscles of mastication. *Pediatr Dent* 1989; 11(4):331.
10. Greene CS. The etiology of temporomandibular disorders: Implications for treatment. *J Orofac Pain* 2001;15(2):93-105.
11. Hiatt JL, Gartner LP. Temporomandibular joint. In: *Textbook of Head and Neck Anatomy*. 2nd ed. Baltimore, Md; Williams and Wilkins; 1987:223.
12. Greco CM, Rudy TE, Turk DC, Herlich A, Zaki HH. Traumatic onset of temporomandibular disorders: Positive effects of a standardized conservative treatment program. *Clin J Pain* 1997;13(4):337-47.
13. Fischer DJ, Mueller BA, Critchlow CW, LeResche L. The association of temporomandibular disorder pain with history of head and neck injury in adolescents. *J Orofac Pain* 2006;20(3):191-8.
14. Imahara SD, Hopper RA, Wang J, Rivara FP, Klein MB. Patterns and outcomes of pediatric facial fractures in the United States: A survey of the National Trauma Data Bank. *J Am Col of Surg* 2008;207(5):710-6.
15. Posnick JC, Wells M, Pron GE. Pediatric facial fractures: Evolving patterns of treatment. *J Oral Maxillofac Surg* 1993;51(5):836-44; discussion 844-5.
16. Kaban L. Acquired abnormalities of the temporomandibular joint. In: Kaban LB, Troulis MJ, eds. *Pediatric Oral and Maxillofacial Surgery*. Philadelphia, Pa; WB Saunders; 2004:340-76.
17. Güven O. A clinical study on temporomandibular joint ankylosis in children. *J Craniofac Surg* 2008;19(5): 1263-9.
18. De Boever JA, Carlsson GE, Klineberg IJ. Need for occlusal therapy and prosthodontic treatment in the management of temporomandibular disorders. Part I. Occlusal interference and occlusal adjustment. *J Oral Rehabil* 2000;27(5):367-79.
19. Taskaya-Yilmaz N, Ögütçen-Toller M, Saraç YŞ. Relationship between the TMJ disc and condyle position on MRI and occlusal contacts on lateral excursions in TMD patients. *J Oral Rehab* 2004;31(8):754-8.
20. Thilander B, Rubio G, Pena L, De Mayorga C. Prevalence of temporomandibular dysfunction and its association with malocclusion in children and adolescents: An epidemiologic study related to specified stages of dental development. *Angle Orthod* 2002;72(2):146-54.
21. Henrikson T, Nilner M. Temporomandibular disorders, occlusion and orthodontic treatment. *J Orthod* 2003;30 (2):129-37; discussion 27.
22. Phillips JT. What skeletal and dental characteristics do TMD patients have in common? *Funct Orthod* 2007;24 (1):24-6,28,30.
23. Pahlkala R, Qvarnström M. Can temporomandibular dysfunction signs be predicted by early morphological or functional variables? *Euro J Orthod* 2004;26(4):367-73.
24. Pullinger AG, Seligman DA. Quantification and validation of predictive values of occlusal variables in temporomandibular disorders using a multifactorial analysis. *J Prosthet Dent* 2000;83(1):66-75.
25. Seligman DA, Pullinger AG. Analysis of occlusal variables, dental attrition, and age for distinguishing healthy controls from female patients with intracapsular temporomandibular disorders. *J Prosthet Dent* 2000;83(1): 76-82.
26. Ciancaglini R, Gherlone EF, Radaelli G. Association between loss of occlusal support and symptoms of functional disturbances of the masticatory system. *J Oral Rehabil* 1999;26(3):248-53.
27. Tallents RH, Macher DJ, Kyrkanides S, Katzberg RW, Moss ME. Prevalence of missing posterior teeth and intraarticular temporomandibular disorders. *J Prosthet Dent* 2002;87(1):45-50.
28. Rey D, Oberti G, Baccetti T. Evaluation of temporomandibular disorders in Class III patients treated with mandibular cervical headgear and fixed appliances. *Am J Orthod Dentofac Orthop* 2008;133(3):379-81.

29. Cheifetz AT, Osganian SK, Allred EN, Needleman HL. Prevalence of bruxism and associated correlates in children as reported by parents. *J Dent Child* 2005;72(2):67-73.
30. Barbosa Tde S, Miyakoda LS, Pocztaruk Rde L, Rocha CP, Gavião MBD. Temporomandibular disorders and bruxism in childhood and adolescence: Review of the literature. *Int J Pediatr Otorhinolaryngol* 2008;72(3):299-314.
31. Castelo PM, Gavião MB, Pereira LJ, Bonjardim LR, Gavião MBD. Relationship between oral parafunctional/nutritive sucking habits and temporomandibular joint dysfunction in primary dentition. *Int J Paediatr Dent* 2005;15(1):29-36.
32. Winocur E, Gavish A, Finkelshtein T, Halachmi M, Gazit E. Oral habits among adolescent girls and their association with symptoms of temporomandibular disorders. *J Oral Rehabil* 2001;28(7):624-9.
33. Carlsson GE, Egermark I, Magnusson T. Predictors of signs and symptoms of temporomandibular disorders: A 20-year follow-up study from childhood to adulthood. *Acta Odontol Scand* 2002;60(3):180-5.
34. Magnusson T, Egermark I, Carlsson GE, Magnusson T, Egermark I, Carlsson GE. A prospective investigation over two decades on signs and symptoms of temporomandibular disorders and associated variables. A final summary. *Acta Odontol Scand* 2005;63(2):99-109.
35. Gesch D, Bernhardt O, Mack F, John U, Kocher T, Dietrich A. Association of malocclusion and functional occlusion with subjective symptoms of TMD in adults: Results of the Study of Health in Pomerania (SHIP). *Angle Orthod* 2005;75(2):183-90.
36. Alamoudi N. Correlation between oral parafunction and temporomandibular disorders and emotional status among Saudi children. *J Clin Pediatr Dent* 2001;26(1):71-80.
37. Sonnesen L, Bakke B, Solow B. Temporomandibular disorder in relation to craniofacial dimensions, head posture and bite force in children selected for orthodontic treatment. *Eur J Orthod* 2001;20(2):179-92.
38. Kondo E, Nakahara R, Ono M, et al. Cervical spine problems in patients with temporomandibular disorder symptoms: An investigation of the orthodontic treatment effects for growing and nongrowing patients. *World J Orthod* 2002;3(4):295-312.
39. Motoyoshi M, Shimazaki T, Namura S. Biomechanical influences of head posture on occlusion: An experimental study using finite element analysis. *Eur J Orthod* 2002;24(4):319-26.
40. Egermark I, Carlsson GE, Magnusson T. A prospective long-term study of signs and symptoms of temporomandibular disorders in patients who received orthodontic treatment in childhood. *Angle Orthod* 2005;75(4):645-50.
41. Henrikson T, Nilner M, Kurol J. Symptoms and signs of temporomandibular disorders before, during and after orthodontic treatment. *Swed Dent J* 1999;23(5-6):193-207.
42. Henrikson T, Nilner M, Kurol J. Signs of temporomandibular disorders in girls receiving orthodontic treatment. A prospective and longitudinal comparison with untreated Class II malocclusions and normal occlusion subjects. *Eur J Orthod* 2000;22(3):271-81.
43. Kim MR, Graber TM, Viana MA. Orthodontics and temporomandibular disorder: A meta-analysis. *Am J Orthod Dentofac Orthop* 2002;121(5):438-46.
44. Alamoudi N, Farsi N, Salako N, Feteih R. Temporomandibular disorders among school children. *J Clin Pediatr Dent* 1998;22(4):323-9.
45. List T, Wahlund K, Wenneberg B, Dworkin SF. TMD in children and adolescents: Prevalence of pain, gender differences, and perceived treatment need. *J Orofac Pain* 1999;13(1):9-20.
46. Stockstill JW, Bowley JF, Dunning D, Spalding P, Stafford K, Erickson L. Prevalence of temporomandibular disorders in children based on physical signs. *J Dent Child* 1998;65(6):459-67, 438.
47. Paesani D, Salas E, Martinez A, Isberg A. Prevalence of temporomandibular joint disk displacement in infants and young children. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999;87(1):15-9.
48. Bonjardim LR, Bavião MB, Carmagnani FG, Pereira LF, Castelo PM. Signs and symptoms of temporomandibular joint dysfunction in children with primary dentition. *J Clin Pediatr Dent* 2003;28(1):53-8.
49. Widmalm SE, Christiansen RL, Gunn SM. Oral parafunctions as temporomandibular disorder risk factors in children. *Cranio* 1995;13(4):242-6.
50. Bonjardim LR, Gavião MB, Pereira LJ, Castelo PM, Garcia RC. Signs and symptoms of temporomandibular disorders in adolescents. *Braz Oral Res* 2005;19(2):93-8.
51. LeResche L, Mancl LA, Drangsholt MT, Saunders K, Von Korff M. Relationship of pain and symptoms to pubertal development in adolescents. *Pain* 2005;118(1-2):201-9.
52. LeResche L, Mancl LA, Drangsholt MT, Huang G, von Korff MV. Predictors of onset of facial pain and temporomandibular disorders in early adolescence. *Pain* 2007;129(3):269-78.
53. McDonald RE, Avery DR, Dean JA. Examination of the mouth and other relevant structures. In: Dean JA, Avery DR, McDonald RE, eds. *McDonald and Avery's Dentistry for the Child and Adolescent*. 9th ed. Maryland Heights Mo: Mosby Elsevier; 2011:1-18.
54. Casamassimo PS, Christensen JR, Fields HW Jr, Ganzberg S. Examination, diagnosis, and treatment planning for general and orthodontic problems. In: Pinkham JR, Casamassimo PS, McTigue DJ, Fields HW Jr, Nowak AJ, eds. *Pediatric Dentistry: Infancy through Adolescence*. 4th ed. St. Louis, Mo; Elsevier Saunders 2005:661-89.

55. American Academy of Pediatric Dentistry. Guideline on recordkeeping. *Pediatr Dent* 2009;31(special issue): 239-46.
56. American Academy of Orofacial Pain. General assessment of the orofacial pain patient. In: de Leeuw R, ed. *Orofacial Pain: Guidelines for Assessment, Diagnosis, and Management*. Carol Stream, Ill: Quintessence Publishing Co Inc; 2008:25-47.
57. Wahlund K, List T, Dworkin SF. Temporomandibular disorders in children and adolescents: Reliability of a questionnaire, clinical examination, and diagnosis. *J Orofac Pain* 1998;12(1):42-51.
58. De Boever JA, Nilner M, Orthlieb JD, Steenks MH, Educational Committee of the European Academy of Craniomandibular Disorders. Recommendations by the EACD for examination, diagnosis, and management of patients with temporomandibular disorders and orofacial pain by the general dental practitioner. *J Orofac Pain* 2008;22(3):268-78.
59. Brooks, SL, Brand JW, Gibbs SJ, et al. Imaging of the tempromandibular joint: A position paper of the American Academy of Oral and Maxillofacial Radiology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1997;83(5):609-18.
60. Hu YS, Schneiderman ED, Harper RP. The temporomandibular joint in juvenile rheumatoid arthritis: Part II. Relationship between computed tomographic and clinical findings. *Pediatr Dent* 1996;18(4):312-9.
61. Bodner L, Miller VJ. Temporomandibular joint dysfunction in children: Evaluation of treatment. *Int J Pediatr Otorhinolaryngol* 1998;44(2):133-7.
62. Wahlund K, List T, Larsson B. Treatment of temporomandibular disorders among adolescents: A comparison between occlusal appliance, relaxation training, and brief information. *Acta Odontol Scand* 2003;61(4):203-11.
63. Kurita H, Kurashina K, Kotani A. Clinical effect of full coverage occlusal splint therapy for specific temporomandibular disorder conditions and symptoms. *J Prosthet Dent* 1997;78(5):506-10.
64. Michelotti A, Steenks MH, Farella M, Parisini F, Cimino R, Martina R. The additional value of a home physical therapy regimen versus patient education only for the treatment of myofascial pain of the jaw muscles: Short-term results of a randomized clinical trial. *J Orofac Pain* 2004;18(2):114-25. [Erratum in *J Orofac Pain* 2006;20(2):106].
65. Medlicott, MS, Harris SR. A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder. *Phys Ther* 2006;86(7): 955-73.
66. Crider, AB, Glaros AG. A meta-analysis of EMG biofeedback treatment of temporomandibular disorders. *J Orofac Pain* 1999;13(1):29-37.
67. List T, Axelsson S, Leijon G. Pharmacologic interventions in the treatment of temporomandibular disorders, atypical facial pain, and burning mouth syndrome. A qualitative systematic review. *J Orofac Pain* 2003;17(4):301-10.
68. Fujii T, Torisu T, Nakamura S. A change of occlusal conditions after splint therapy for bruxers with and without pain in the masticatory muscles. *Cranio* 2005;23(2):113-8.
69. Koh H, Robinson PG. Occlusal adjustment for treating and preventing temporomandibular joint disorders. *The Cochrane Database of Systematic Reviews* 2003;1:Art. No. CD003812. DOP: 10.1002/146751858.CD003812.