

Use of Antibiotic Therapy for Pediatric Dental Patients

Latest Revision

2022

How to Cite: American Academy of Pediatric Dentistry. Use of antibiotic therapy for pediatric dental patients. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2023:537-41.

Abstract

This best practice provides clinicians with guidance in the appropriate use of antibiotics to treat oral infections in children. When correctly prescribed and administered, antibiotics can be effective in the treatment of oral bacterial infections. Antibiotic stewardship is important given the rise in antibiotic-resistant microorganisms and potential for adverse drug reactions. This document addresses the following clinical conditions: oral wounds, pulpitis/apical periodontitis/drainage sinus tract/localized intraoral swelling, acute facial swelling of dental origin, dental trauma, periodontal diseases, and salivary gland infections and offers guidance on the judicious use of antibiotics in their management. Antibiotics are not indicated in the management of conditions of viral origin. Potential interactions between antibiotics and oral contraceptives are addressed. Health care providers must be prudent in their prescribing practices to maximize effectiveness and minimize bacterial resistance and adverse reactions.

This document was developed through a collaborative effort of the American Academy of Pediatric Dentistry Councils on Clinical Affairs and Scientific Affairs to offer updated information and guidance on the use of antibiotic therapy for pediatric dental patients.

KEYWORDS: ANTIBIOTICS; ANTIMICROBIAL RESISTANCE; DENTAL INFECTION CONTROL, BACTERIAL INFECTIONS

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes the increasing prevalence of antibiotic-resistant microorganisms and potential for adverse drug reactions and interactions. These recommendations are intended to provide guidance in the proper and judicious use of antibiotic therapy in the treatment of oral conditions. The use of antibiotic prophylaxis for dental patients at risk for infection is addressed in a separate best practices document.¹ Information regarding commonly prescribed antibiotics can be found in AAPD's *Useful Medications for Oral Conditions*.²

Methods

Recommendations on the use of antibiotic therapy were developed by the Council on Clinical Affairs, adopted in 2001³, and last revised in 2019.⁴ This revision was based upon a new literature search of the PubMed®/MEDLINE database using the terms: pediatric dental antibiotic therapy AND antibacterial agents, antimicrobial agents, dental trauma, oral wound management, orofacial infections, periodontal disease, viral disease, and oral contraception; fields: all; limits: within the last 10 years, humans, English, clinical trials, birth through age 18. Four hundred seventy-eight articles matched these criteria. Papers for review were chosen from this search and from hand searching. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

Background

Antibiotics are beneficial to patient care when prescribed and administered correctly for bacterial infections. However, the widespread use of antibiotics has permitted common bacteria to develop resistance to drugs that once controlled them.^{5,6} Drug resistance is prevalent throughout the world.^{5,6} Each year in the United States, nearly three million antibiotic-resistant infections occur and result in more than 35,000 deaths.⁵ Some microorganisms may develop resistance to a single antimicrobial agent, while others develop multi-drug-resistant strains.⁶ To diminish the rate at which resistance is increasing, health care providers must be prudent in the use of antibiotics.^{5,7} A study showed 80 percent of prescriptions of antibiotics before dental procedures were unnecessary as risk-factors were not present.⁸ This highlights a concern on the appropriateness for prescribed antibiotic prophylaxis for dental procedures.⁸ While use of antibiotic prophylaxis is indicated for certain patients undergoing invasive dental procedures, overall emphasis should focus on establishment of a dental home, the prevention of disease, establishment and maintenance of good oral health care habits, and regular dental care.^{9,10} Conservative use of antibiotics is indicated to minimize the risk of developing resistance to current antibiotic regimens.^{5,6,9}

ABBREVIATIONS

AAPD: American Academy of Pediatric Dentistry. **JRP:** Juvenile recurrent parotitis.

Adverse events such as allergic reactions, development of *Clostridioides difficile* infection, or drug interactions and side effects can occur.^{5,11} Antibiotic adverse drug events are a common cause of emergency department visits for adverse drug events in children under the age of 18 years, with amoxicillin as the most commonly implicated drug in children less than nine years and sulfamethoxazole-trimethoprim in children aged 10-19.¹²

Amoxicillin, considered the first drug of choice for dental infections in non-allergic children¹³, is effective against a wide variety of gram-positive bacteria and offers greater gram-negative coverage than penicillin.¹⁴ It has been shown to be effective against oral flora¹⁵, be well absorbed from the gastrointestinal tract⁹, provide high, sustained serum concentrations⁹, and have a low incidence of adverse effects¹⁵. The American Heart Association no longer recommends clindamycin for prophylaxis against infective endocarditis due to frequent and severe reactions.⁹ Clindamycin has been associated with significant adverse drug reactions related to community-acquired *C. difficile* infections.⁹ Up to 15 percent of community-acquired *C. difficile* infection has been attributed to antibiotics prescribed for dental procedures.⁹ Doxycycline is recommended as an alternative to penicillin, cephalosporin, and macrolide allergy.⁹ Short-term use (less than 21 days) of doxycycline had not been associated with tooth discoloration in children under eight years of age.¹⁶⁻¹⁸ Azithromycin is one of the safest antibiotics for patients allergic to penicillins, but there are risks of cardiac complications including cardiotoxicity.¹⁹ The small, heightened risk appears to be related to pre-existing cardiovascular risk factors including prior myocardial infarction, diabetes, age, and gender.²⁰ Cardiac risk in pediatric patients seems to be due to an increased risk of QT prolongation associated with higher dosage levels.¹⁹

Recommendations

Practitioners should adhere to the following general principles for antibiotic usage for the pediatric dental patient.^{9,21}

- Prevention of dental diseases should be emphasized in order to reduce the need for antibiotic intervention.
- Antibiotics should be prescribed only when truly needed for a bacterial infection and only as an adjunct to, not an alternative for, other interventions (e.g., pulp therapy, extraction, scaling and root planing) implemented to control the infection source.
- Antibiotics should be selected based on properties of the agent (e.g., spectrum of coverage, safety), previous antibiotic use, and patient considerations (e.g., medical history, drug allergies, current medication use, ease of use) and then prescribed at an adequate pediatric dose.
- The most effective route of drug administration (intravenous versus intramuscular versus oral) must be considered. If the patient is receiving parenteral antimicrobial therapy for treatment of existing infections, the same antibiotic can be continued.⁹ Consultation

with an infectious disease physician is recommended if there is concern for resistant infections.

- The traditional minimal duration of drug regimen is five days beyond the point of substantial improvement (e.g., improved healing of wound, reduction of erythema or swelling, reduction of signs and symptoms). Usually, this is a five- to seven-day course of treatment, dependent upon the specific drug selected.^{22,23}
- However, in light of the growing problem of drug resistance, discontinuation of antibiotics should be considered following determination of either ineffectiveness or cure prior to completion of a full course of therapy.^{24,25}
- If an infection is not responsive to the initial drug selection, a culture and sensitivity testing from the infection site or, in some cases, a blood microbiology and culture and sensitivity may be indicated.^{5,25}
- Prescriptions should be documented in the patient's dental record.²⁶
- Individuals suspected to have an allergy to antibiotics should receive testing to confirm or refute the presence of a true allergy.

Additional considerations for specific clinical circumstances are discussed below.

Oral wounds

Factors related to host risk (e.g., age, systemic illness, comorbidities, malnutrition) and type of wound (e.g., laceration, puncture) must be evaluated when determining the risk for infection and subsequent need for antibiotics. Wounds can be classified as clean, potentially contaminated, or contaminated/dirty. Facial lacerations and puncture wounds may require topical antibiotic agents.²⁷ Intraoral puncture wounds and lacerations that appear to have been contaminated by extrinsic bacteria, debris (e.g., dirt, soil, gravel), foreign body, open fractures, and joint injury have an increased risk of infection and should be managed by systemic antibiotics.²⁷ If antibiotics are deemed beneficial to the healing process, the timing of their administration is critical to supplement the natural host resistance in bacterial killing. The drug should be administered as soon as possible for the best result.⁵

Pulpitis/apical periodontitis/draining sinus tract/localized intraoral swelling

Bacteria can gain access to the pulpal tissue through caries, exposed pulp or dentinal tubules, cracks into the dentin, and defective restorations. If a child presents with acute symptoms of pulpitis, treatment (i.e., pulpotomy, pulpectomy, or extraction) should be rendered. Antibiotic therapy is not indicated nor effective if the dental infection is contained within the pulpal tissue or the immediate surrounding tissue. In this case, the child will have no systemic signs of an infection (i.e., no fever, no facial swelling).²⁸

Consideration for use of antibiotics should be given in cases of advanced non-odontogenic bacterial infections such as staphylococcal mucositis, tuberculosis, gonococcal stomatitis, and oral syphilis. If suspected, referral for microbiology, culture and sensitivity testing, biopsy, or other laboratory tests for documentation and definitive treatment is indicated.

Acute facial swelling of dental origin

A child presenting with a facial swelling or facial cellulitis secondary to an odontogenic infection should receive prompt dental attention. The clinician should consider age, cooperation, the ability to obtain adequate anesthesia (local versus general), the severity of the infection, the medical status, and any social issues of the child.^{29,30} For odontogenic infections with nonlocalized and progressive swelling and systemic manifestations (e.g., fever, difficulty breathing or swallowing), immediate surgical intervention and medical management with intravenous antibiotic therapy contribute to a more rapid cure.³⁰⁻³³ Signs of systemic involvement and septicemia (e.g., fever, malaise, asymmetry, facial swelling, lymphadenopathy, trismus, tachycardia, dysphagia, airway compromise, respiratory distress) warrant emergency treatment.^{31,32} Additional imaging (e.g., radiographs, ultrasound, computed tomography scan) and testing (e.g., complete blood examination, c-reactive protein, bacterial culture and sensitivity testing) can aid in assessment and diagnosis.^{29,30} Penicillin derivatives remain the empirical choice for odontogenic infections; however, consideration of additional adjunctive antimicrobial therapy such as metronidazole can be given for anaerobic bacterial involvement.^{24,35} Cephalosporins could be considered as an alternative choice for management of odontogenic infections, especially when a child has had previous course(s) of penicillin/amoxicillin or if the child has a penicillin allergy.³⁵

Avulsions

Systemic antibiotics have been recommended as adjunctive therapy for avulsed permanent incisors with an open or closed apex.¹⁵ Amoxicillin or penicillin is the drug of choice due to effectiveness against oral flora and low incidence of adverse effects.¹⁵ Doxycycline is recommended as an alternative to penicillin.¹⁵ Doxycycline exhibits antimicrobial, anti-inflammatory, and antiresorptive properties which make its use appropriate for dental trauma.^{15,36} Using topical antibiotics (minocycline or doxycycline) to enhance pulpal revascularization and periodontal healing in immature nonvital traumatized teeth has shown potential in animal studies, but usage has not been proven effective in human studies, remains controversial, and has not been recommended by the International Association of Dental Traumatology.¹⁵ Further randomized clinical trials are needed.¹⁵ Antibiotics can be warranted in cases of concomitant soft tissue injuries (see **Oral wounds**) and when dictated by the patient's medical status.

Pediatric periodontal diseases

Three distinct forms of periodontal disease have been defined as: (1) periodontitis (grouping the two forms formerly recognized as aggressive or chronic); (2) necrotizing periodontitis; and (3) periodontitis as a manifestation of systemic disease.³⁷ Patients diagnosed with what formerly was known as aggressive periodontal disease may require adjunctive antimicrobial therapy in conjunction with localized treatment.³⁸⁻⁴⁰ In pediatric periodontal diseases associated with systemic conditions (e.g., severe congenital neutropenia, Papillon Lefèvre syndrome, leukocyte adhesion deficiency), the immune system is unable to control the growth of periodontal pathogens and, in some cases, treatment may involve antibiotic therapy or antibiotic prophylaxis.^{10,38} Culture and susceptibility testing of isolates from the involved sites are helpful in guiding the drug selection.⁴¹ In severe and refractory cases, extraction is indicated.⁴¹

Viral diseases

Conditions of viral origin such as acute primary herpetic gingivostomatitis should not be treated with antibiotic therapy.¹¹

Salivary gland infections

For acute salivary gland swellings of bacterial nature, antibiotic therapy is indicated.⁴² If the patient does not improve in 24-48 hours on antibiotics alone, incision and drainage may be warranted.⁵ Amoxicillin/clavulanate is used as empirical therapy to cover both staphylococcal and streptococcal species as most bacterial infections of the salivary glands originate from oral flora.⁴²

The most common inflammatory salivary gland disorder in the United States is juvenile recurrent parotitis (JRP), with first onset of symptoms between the ages of three and six, continuing to puberty.⁴³ Although JRP is self-limiting, administration of β -lactam antibiotics may shorten symptom duration.⁴³ For both acute bacterial submandibular sialadenitis and chronic recurrent submandibular sialadenitis, antibiotic therapy is included as part of the treatment.⁴⁴

Oral contraceptive use

Although caution previously was advised with the concomitant use of antibiotics and oral contraceptives,^{45,46} a 2018 systematic review of drug interactions between non-rifamycin antibiotics and hormonal contraception found that most women can expect no reduction in hormonal contraceptive effect with the concurrent use of non-rifamycin antibiotics.⁴⁷ The World Health Organization also reported in 2015 that most broad-spectrum antibiotics do not affect the contraceptive effectiveness of combined oral contraceptives, combined contraceptive patch, or the combined contraceptive vaginal ring.⁴⁸ In addition, no differences in ovulation were found when oral contraceptives were combined with ampicillin, doxycycline, temafloxacin, ofloxacin, ciprofloxacin, clarithromycin, roxithromycin, dirithromycin, or metronidazole.⁴⁷ Women should be encouraged to take oral contraceptives correctly and consistently at all times, including during periods of illness.⁴⁷ Rifamycin antibiotics,

such as rifampin or rifabutin, induce hepatic enzymes that are required for hormonal contraceptive metabolism, which could compromise the contraceptive or antibiotic effect.^{47,48} Consultation with the medical practitioner regarding use of other contraceptives is recommended with long-term use of these medications.⁴⁸

References

1. American Academy of Pediatric Dentistry. Antibiotic prophylaxis for dental patients at risk for infection. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill.: American Academy of Pediatric Dentistry; 2022: 500-6.
2. American Academy of Pediatric Dentistry. Useful medications for oral conditions. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill.: American Academy of Pediatric Dentistry; 2022:628-35.
3. American Academy of Pediatric Dentistry. Appropriate use of antibiotic therapy. *Pediatr Dent* 2001;23(special issue):71-3.
4. American Academy of Pediatric Dentistry. Use of antibiotic therapy for pediatric dental patients. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill.: American Academy of Pediatric Dentistry; 2019:443-6.
5. Centers for Disease Control and Prevention. Antibiotic/Antimicrobial Resistance Threats in the United States, 2019. Available at: "<https://www.cdc.gov/drugresistance/pdf/threats-report/2019-ar-threats-report-508.pdf>". Accessed January 24, 2022.
6. Costelloe C, Metcalfe C, Lovering A, et al. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-analysis. *BMJ* 2010;340:c2096.
7. Aidasani B, Solankis M, Khetarpal S, Ravi Pratap S. Antibiotics: Their use and misuse in paediatric dentistry. A systematic review. *Eur J Paediatr Dent* 2019;20(2):133-8.
8. Suda KJ, Calip GS, Zhou J, et al. Assessment of the appropriateness of antibiotic prescriptions for infection prophylaxis before dental procedures. *JAMA Netw Open* 2019;2(5):e193909.
9. Wilson WR, Gewitz M, Lockhart PB, et al. Prevention of viridans group streptococcal infective endocarditis: A scientific statement from the American Heart Association. *Circulation* 2021;143(20):e963-e978. Available at: "<https://www.ahajournals.org/doi/pdf/10.1161/CIR.00000000000969>". Accessed March 26, 2022. Erratum in: *Circulation* 2021;144(9):e192.
10. Squire JD, Gardner PJ, Moutsopoulos NM, Leiding JW. Antibiotic prophylaxis for dental treatment in patients with immunodeficiency. *J Allergy Clin Immunol Pract* 2019;7(3):819-23.
11. Centers for Disease Control and Prevention. Antibiotic Prescribing and Use. Antibiotic Use in Outpatient Settings, 2017. Available at: "<https://www.cdc.gov/antibiotic-use/stewardship-report/pdf/stewardship-report.pdf>". Accessed June 30, 2022.
12. Lovegrove MC, Geller A, Fleming-Dutra KE, Shehab N, Sapiano MRP, Budnitz DS. US emergency department visits for adverse drug events from antibiotics in children, 2011-2015. *J Pediatric Infect Dis Soc* 2019;8(5): 384-91.
13. Ahmadi H, Ebrahimi A, Ahmadi F. Antibiotic therapy in dentistry. *Int J Dent* 2021;2021:6667624.
14. Akhavan BJ, Khanna NR, Vijhani P. Amoxicillin. [Updated 2021 Aug 17]. In: StatPearls [Internet]. Treasure Island, Fla.: StatPearls Publishing; 2022 Jan. Available at: "<https://www.ncbi.nlm.nih.gov/books/NBK482250/>". Accessed June 30, 2022.
15. Fouad AF, Abbott PV, Tsilingaridis G, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 2. Avulsion of permanent teeth. *Dental Traumatology* 2020;36(4): 331-42.
16. Todd SR, Dahlgren FS, Traeger MS, et al. No visible dental staining in children treated with doxycycline for suspected Rocky Mountain Spotted Fever. *J Pediatr* 2015;166(5):1246-51. Available at: "<https://doi.org/10.1016/j.jpeds.2015.02.015>". Accessed January 24, 2022.
17. American Academy of Pediatrics. Tetracyclines. In: Kimberlin DW, Barnett ED, Lynfield R, Sawyer MH, eds. *Red Book: 2021-2024 Report of the Committee on Infectious Diseases*. Elk Grove Village, Ill.: American Academy of Pediatrics; 2021:905-6.
18. Stultz JS, Eiland LS. Doxycycline and tooth discoloration in children: Changing of recommendations based on evidence of safety. *Ann Pharmacother* 2019;53(11): 1162-6. Available at: "<https://doi.org/10.1177/1060028019863796>". Accessed September 26, 2021.
19. Zeng L, Xu P, Choonara I, et al. Safety of azithromycin in pediatrics: A systematic review and meta-analysis. *Eur J Clin Pharmacol* 2020;76(12):1709-21.
20. Bartold PM, du Bois AH, Gannon S, Haynes DR, Hirsch RS. Antibacterial and immunomodulatory properties of azithromycin treatment implications for periodontitis. *Inflammopharmacology* 2013;21(4):321-38.
21. Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases. Division of Healthcare Quality Promotion. Checklist for antibiotic prescribing in dentistry. 2019. Available at: "<https://www.cdc.gov/antibiotic-use/community/downloads/dental-fact-sheet-FINAL.pdf>". Accessed March 11, 2022.
22. Kuriyama T, Karasawa T, Nakagawa K, Saiki Y, Yamamoto E, Nakamura S. Bacteriological features and antimicrobial susceptibility in isolates from orofacial odontogenic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000;90(5):600-8.
23. Prieto-Prieto J, Calvo A. Microbiological basis of oral infections and sensitivity to antibiotics. *Med Oral Patol Oral Cir Bucal* 2004;9(suppl S):11-8.

24. American Academy of Pediatrics. Group A Streptococcal Infections. In: Kimberlin DW, Barnett ED, Lynfield R, Sawyer MH, eds. *Red Book: 2021-2024 Report of the Committee on Infectious Diseases*. Elk Grove Village, Ill.: American Academy of Pediatrics; 2021:694-915.
25. Engleberg NC. Strategies to combat bacterial infections. In: Engleberg NC, DiRita VJ, Imperiale MJ, eds. *Schaechter's Mechanisms of Microbial Disease*. 6th ed. Philadelphia, Pa.: Wolters-Kluwer; 2022:436.
26. American Academy of Pediatric Dentistry. Record-keeping. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill., USA: American Academy of Pediatric Dentistry; 2022:521-8.
27. Nakamura Y, Daya M. Use of appropriate antimicrobials in wound management. *Emerg Med Clin North Am* 2007;25(1):159-76.
28. Fluent MT, Jacobsen PL, Hicks LA. Considerations for responsible antibiotic use in dentistry. *J Am Dent Assoc* 2016;147(8):683-6.
29. Johri A, Piecuch JF. Should teeth be extracted immediately in the presence of acute infection? *Oral Maxillofac Surg Clin North Am* 2011;23(4):507-11.
30. Thikkurissy S, Rawlins JT, Kumar A, Evans E, Casamassimo PS. Rapid treatment reduces hospitalization for pediatric patients with odontogenic-based cellulitis. *Am J Emerg Med* 2010;28(6):668-72.
31. Adewumi AO. Oral surgery in children. In: Nowak AJ, Christensen JR, Mabry TR, Townsend JA, Wells MH, eds. *Pediatric Dentistry Infancy through Adolescence*. 6th ed. St Louis, Mo.: Elsevier; 2019:399-409.
32. Baker S, Parico L. Pathologic paediatric conditions associated with a compromised airway. *Int J Paediatr Dent*. 2010;20(2):102-11.
33. Solankis M, Khetarpal S, Ravi Pratap S. Antibiotics: Their use and misuse in paediatric dentistry. A systematic review. *Eur J Paediatr Dent* 2019;20(2):133-8.
34. Zirk M, Buller J, Goeddertz P, et al. Empiric systemic antibiotics for hospitalized patients with severe odontogenic infections. *J Craniomaxillofac Surg* 2016;44(8):1081-8.
35. Andreason JO, Storgaard Jensen S, Sae-Lim V. The role of antibiotics in preventing healing complications after traumatic dental injuries: A literature review. *Endod Topics* 2006;14:80-92.
36. Caton JC, Armitage G, Berglundh T, et al. A new classification scheme for periodontal and peri-implant diseases and conditions: Introduction and key changes from the 1999 classification. *J Periodontol* 2018;89(Suppl 1):S1-S8.
37. American Academy of Pediatric Dentistry. Classification of periodontal diseases in infants, children, adolescents, and individuals with special health care needs. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill., USA: American Academy of Pediatric Dentistry; 2022:451-65.
38. Keestra JAJ, Grosjean I, Coucke W, Quirynen M, Teughels W. Non-surgical periodontal therapy with systemic antibiotics in patients with untreated aggressive periodontitis: A systematic review and meta-analysis. *J Periodont Res* 2015;50(6):689-706.
39. Rabelo CC, Feres M, Gocalves C, et al. Systematic antibiotics in the treatment of aggressive periodontitis. A systematic review and a Bayesian Network meta-analysis. *J Clin Periodontol* 2015;42(7):647-57.
40. Schmidt JC, Walter C, Rischewski JR, Weiger R. Treatment of periodontitis as a manifestation of neutropenia with or without systemic antibiotics: A systematic review. *Pediatr Dent* 2013;35(2):E54-E63.
41. Patel A, Karlis V. Diagnosis and management of pediatric salivary gland infections. *Oral Maxillofac Surg Clin North Am* 2009;21:345-52
42. Garavello W, Redaelli M, Galluzzi, F. Juvenile recurrent parotitis: A systematic review of treatment studies. *Int J Pediatr Otorhinolaryngol* 2018;112:151-7.
43. Carlson ER. Diagnosis and management of salivary gland infections. *Oral Maxillofac Surg Clin North Am* 2009;21(3):293-312.
44. DeRossi SS, Hersh EV. Antibiotics and oral contraceptives. *Pediatr Clin North Am* 2002;46(4):653-64.
45. Becker DE. Adverse drug interactions. *Anesth Prog* 2011; 58(1):31-41.
46. Simmons K, Haddad L, Nanda K, Curtis, K. Drug interactions between non-rifamycin antibiotics and hormonal contraception: A systematic review. *Am J Obstet Gynecol* 2018;218(1):88-97.
47. World Health Organization. *Medical Eligibility Criteria for Contraceptive Use*. 5th ed. Geneva, Switzerland: World Health Organization; 2015. Available at: "[https:// www.who.int/reproductivehealth/publications/family _planning/ MEC-5/en/](https://www.who.int/reproductivehealth/publications/family_planning/MEC-5/en/)". Accessed January 24, 2022.