

# Periodontal Conditions in Pediatric Dental Patients

## Revised

2024

**How to Cite:** American Academy of Pediatric Dentistry. Periodontal conditions in pediatric dental patients. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2024:505-22.

## Abstract

*This best practice serves as a resource for clinicians to guide periodontal diagnosis, risk assessment, and management in pediatric dental patients, including those with special health care needs. Recommendations on the management of contributing factors and conditions that increase the risk for periodontal disease and pathologies, as well as treatment considerations on the use of adjunctive antibiotics and surgical therapies, are reviewed. Special attention is focused on care coordination, collaborations, and referral of care to specialists. In cases where the published data regarding periodontal diseases and pathologies among children and adolescents was limited, recommendations were extrapolated from evidenced-based literature among adult patients, as well as on the consensus opinions of the working group.*

*This document was developed through a collaborative effort of the American Academy of Pediatric Dentistry Councils on Clinical Affairs and Scientific Affairs to offer information and guidance regarding risk assessment and management of periodontal conditions in pediatric dental patients.*

**KEYWORDS:** PERIODONTAL DISEASES; PERIODONTITIS; RISK ASSESSMENT; BLEEDING ON PROBING; GINGIVITIS; GINGIVAL DISEASES; CHILD; ADOLESCENT

## Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes the importance of periodontal health and its effect on the well-being of pediatric patients, including those with special health care needs (SHCN). Periodontal-risk assessment (PRA) and management protocols are essential elements of contemporary clinical care for these patients. These recommendations are intended to assist practitioners in assessing risk for, diagnosing/classifying, and managing periodontal conditions in pediatric dental patients.

## Methods

Recommendations on risk assessment and management of periodontal diseases, developed by the Council on Clinical Affairs utilizing a consultant in periodontics, were adopted in 2022.<sup>1</sup> Utilizing a similar process and merging AAPD's *Classification of Periodontal Diseases in Infants, Children, Adolescents, and Individuals with Special Health Care Needs*<sup>2</sup>, this revision is based on searches of PubMed®/MEDLINE and Google Scholar databases using the terms: periodontal health AND children, periodontal health AND adolescents, gingival disease AND children, gingival disease AND adolescents, periodontal disease AND children, periodontal disease AND adolescents, gingivitis AND prevalence, periodontitis AND prevalence, gingival disease AND prevalence, periodontal disease AND prevalence, dental plaque AND children, dental plaque AND adolescents; periodontitis as a manifestation of systemic diseases, necrotizing periodontitis, aggressive periodontitis, localized periodontitis; fields: all; limits: within the last 10 years, human, English, clinical study, clinical trial, comparative study, multicenter study, observational study, randomized clinical trial, meta-analysis, and systematic reviews. In addition, papers from proceedings of the 2017 World Workshop on the Classification

of Periodontal and Peri-implant Diseases and Conditions<sup>3</sup> were reviewed. The articles were evaluated by title and/or abstract and relevance to dental care for children and adolescents. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

## Background

Studies show that gingivitis occurs in half of the population by age of four or five years and peaks nearly to 100 percent at puberty.<sup>4</sup> A periodontal examination and risk assessment are important parts of a comprehensive oral evaluation and periodic examination of pediatric dental patients. Morphological changes in the primary, mixed, and permanent dentitions reflect typical patterns of oral growth and development. The ability to distinguish normal physiological changes from gingival and periodontal diseases enables accurate diagnoses and prevents unnecessary treatment. Maintenance and restoration of gingival and periodontal health during childhood and adolescence will facilitate healthy gingival and periodontal health at older ages. Early diagnosis ensures the greatest opportunity for successful treatment, primarily by reducing etiological factors, establishing appropriate therapeutic measures, and developing an effective periodic maintenance protocol.<sup>5</sup>

## ABBREVIATIONS

**AAPD:** American Academy of Pediatric Dentistry. **BoP:** Bleeding on probing. **CAL:** Clinical attachment loss. **CEJ:** Cementoenamel junction. **CHX:** Chlorhexidine. **mm:** millimeter. **NSAIDs:** Non-steroidal anti-inflammatory drugs. **PDL:** Periodontal ligament. **PPD:** Periodontal pocket depth. **PRA:** Periodontal-risk assessment. **SHCN:** Special health care needs. **SIB:** Self-injury behavior. **SRP:** Scaling and root planning.

## Recommendations

### Diagnostic phase

The diagnostic criteria for gingivitis are based on clinical features, taking into consideration the presence of plaque biofilm and that the inflammatory response to plaque biofilm is an age-dependent phenomenon. Three distinct forms of periodontal disease have been defined as: (1) periodontitis (single category grouping the two forms of the disease formerly recognized as aggressive or chronic); (2) necrotizing periodontitis; and (3) periodontitis as a manifestation of systemic conditions.<sup>3</sup> Diagnosis may include genetic, microbiological, and biochemical tests or gingival biopsy. Diagnosis involves staging and grading of the periodontal disease.<sup>6</sup> Staging considers the severity and extent of disease while grading assesses the future risk of periodontitis progression and anticipated treatment outcomes. Grading also allows the clinician to incorporate individual patient risk factors into the diagnosis (Table 1).

### Periodontal-risk assessment (PRA)

Risk factors for periodontal disease are complex and may be biological, environmental (social), and behavioral.<sup>7</sup> PRA identifies risk factors that place individuals at an increased risk of developing gingival and periodontal diseases and pathologies, as well as factors that influence the progression of the disease. PRA can improve clinical decision making and allows the implementation of individualized treatment planning and proactive targeted interventions.<sup>8</sup> Evidenced-based PRA tools have been developed based on studies conducted among adult patients.<sup>9</sup> Due to the limited literature regarding PRA among children and adolescents, factors associated with elevated risk were extrapolated from evidence from adult patients<sup>10-15</sup> (Tables 2 and 3).

### Prognosis and treatment planning

Determination of the prognosis follows the diagnostic phase and is a dynamic process to be re-evaluated at all therapeutic phases (i.e., systemic, behavioral, nonsurgical, surgical, maintenance). Prognosis, based on the probability of disease progression and clinical parameters, can be categorized as favorable, questionable, unfavorable, and hopeless.<sup>16</sup>

The treatment plan is formulated after completing a comprehensive oral evaluation, establishing a diagnosis, determining the prognosis, and identifying the individual needs and desires of the patient and parent. It addresses immediate, intermediate, and long-term goals to arrest or slow periodontal disease progression. Important considerations include emergency treatment for pain or infections, need for exodontia, and esthetic demands.<sup>17</sup>

### General considerations

- A periodontal assessment includes a discussion of the chief complaint, detailed medical, dental, and social history reviews, extra- and intraoral examinations, radiographs, and periodontal probing as indicated. Further investigations (e.g., genetic, microbiological, gingival biopsy, and biochemical

tests) may be needed on an individual basis to differentiate types of periodontal diseases.

- Bleeding on probing (BoP) in primary teeth during early childhood, even at a low number of sites, is indicative of high susceptibility to periodontal diseases, due to the age-dependent reactivity of the gingival tissues to plaque.<sup>18,19</sup>
- Probing assessments may be initiated after the eruption of the first permanent molars and incisors and only if tolerated by the child. Pseudopockets (greater than three millimeters [mm]) may be present around partially and newly erupted teeth.<sup>20</sup> Probing assessment on primary teeth is required before the eruption of the first permanent molars and incisors when clinical and radiographic findings indicate the presence of periodontal diseases.
- Assessing for generalized (i.e., involving 30 or more percent of the teeth) gingivitis may be performed for patients unable to undergo probing due to age, anxiety, or SHCN.<sup>21</sup>
- Alveolar bone loss in the primary dentition indicates increased susceptibility to periodontal disease.<sup>22-24</sup>
- Good quality bitewing radiographs are necessary for diagnosing alveolar bone loss.<sup>24-26</sup> While bitewing radiographs are useful when assessing abnormal molar mobility,<sup>23-25,27</sup> periapical radiographs may help rule out any other associated pathology (e.g., root resorption). For abnormal anterior tooth mobility, periapical radiographs are the most appropriate images.<sup>28</sup>
- $1 \pm 0.5$  mm distance from the most coronal portion of the alveolar bone crest to the cemento-enamel junction (CEJ) is considered a normal alveolar bone height in the primary dentition,<sup>23,24,29</sup> while a distance of more than two mm is considered to represent bone loss.<sup>22</sup> A distance of more than two mm may be considered normal when the bone is adjacent to exfoliating primary teeth or erupting permanent teeth.<sup>30</sup>
- Two-mm distance (on average, varying between  $1.0 \pm 3.0$  mm) from the most coronal portion of the alveolar bone crest to the CEJ is considered a normal alveolar bone height in the permanent dentition.<sup>25</sup>

### Recommendations

- For patients in the primary dentition, a visual assessment of the gingiva should be part of every comprehensive oral evaluation and periodic examination. All dental radiographs should be examined for evidence of caries, alveolar bone loss, developmental anomalies, and other pathologies.
- A simplified basic periodontal examination is recommended for individuals aged seven to 17 years.<sup>20</sup> After the eruption of the first permanent molars and incisors, six index teeth (the first permanent molars, the permanent maxillary right central incisor, and the permanent mandibular left central incisor) are assessed for: (1) BoP; (2) presence of calculus; (3) plaque retention factors; (4) periodontal pocket depth (PPD); (5) furcation involvement; and (6) recession.

**Table 1. Periodontitis Staging and Grading (Adapted from Tonetti et al.<sup>6</sup>)**

Framework for periodontitis staging and grading		Disease Severity and Complexity of Management			
		Stage I: Initial periodontitis	Stage II: Moderate periodontitis	Stage III: Severe periodontitis with potential for additional tooth loss	Stage IV: Advanced periodontitis with extensive tooth loss and potential for loss of dentition
Evidence or risk of rapid progression, anticipated treatment response, and effects on systemic health	Grade A	Individual Stage and Grade Assignment			
	Grade B				
	Grade C				
Periodontitis stage		Stage I	Stage II	Stage III	Stage IV
Severity	Interdental CAL at site of greatest loss	1 to 2 mm	3 to 4 mm	≥5 mm	≥5 mm
	Radiographic bone loss	Coronal third (<15%)	Coronal third (<15% to 33%)	Extending to mid-third of root and beyond	Extending to mid-third of root and beyond
	Tooth loss	No tooth loss due to periodontitis		Tooth loss due to periodontitis of ≤4 teeth	Tooth loss due to periodontitis of ≥5 teeth
Complexity	Local	Maximum probing depth ≤4 mm Mostly horizontal bone loss	Maximum probing depth ≤5 mm Mostly horizontal bone loss	In addition to stage II complexity: – Probing depth ≥6 mm – Vertical bone loss ≤3 mm – Furcation involvement Class II or III – Moderate ridge defect	In addition to stage III complexity: Need for complex rehabilitation due to: – Masticatory dysfunction – Secondary occlusal trauma (tooth mobility degree ≥2) – Severe ridge defect – Bite collapse, drifting, flaring – Less than 20 remaining teeth (10 opposing pairs)
Extent and distribution	Add to stage as descriptor	For each stage, describe extent as localized (<30% of teeth involved), generalized, or molar/incisor pattern			
Periodontitis grade		Grade A: Slow rate of progression		Grade B: Moderate rate of progression	Grade C: Rapid rate of progression
Primary criteria	Direct evidence of progression	Longitudinal data (RBL or CAL)	Evidence of no loss over 5 years	<2 mm over 5 years	≥2 mm over 5 years
	Indirect evidence of progression	% bone loss/age	<0.25	0.25 to 1.0	>1.0
		Case phenotype	Heavy biofilm deposits with low levels of destruction	Destruction commensurate with biofilm deposits	Destruction exceeds expectation given biofilm deposits; specific clinical patterns suggestive of periods of rapid progression and/or early onset disease (e.g., molar/incisor pattern; lack of expected response to standard bacterial control therapies)
Grade modifiers	Risk factors	Smoking	Non-smoker	Smoker <10 cigarettes/day	Smoker ≥10 cigarettes/day
		Diabetes	Normoglycemic/no diagnosis of diabetes	HbA1c <7.0% in patients with diabetes	HbA1c ≥7.0% in patients with diabetes

- PRA, based on a child's age and biological, social/behavioral, and clinical/radiographic factors, should be a routine component of initial and periodic oral examinations.
- Practitioners may use the estimated risk level to establish a periodicity and intensity of diagnostic, counseling, and therapeutic interventions (Table 4).
- The treatment plan should be used to establish the methods and sequence of delivering periodontal treatment and include:
  - periodontal procedures to be performed;
  - medical consultation or referral for treatment when indicated;
  - monitoring during the course of periodontal therapy;
  - consideration of diagnostic testing that may include genetic, microbiological, gingival biopsy, or

biochemical tests or monitoring during the course of periodontal therapy;

- consideration of adjunctive restorative, prosthetic, orthodontic, and/or endodontic consultation or treatment;
- consideration of chemotherapeutic and antibiotic agents for adjunctive treatment;
- provision for re-evaluation during and after periodontal or dental implant therapy; and
- periodontal maintenance program.

### Behavioral phase

The success of both prevention and treatment of periodontal diseases and conditions relies significantly on the ability of the patient/caregivers to comply with requested oral hygiene (e.g., brushing, flossing) and nutrition (e.g., vitamin C intake)

**Table 2. Factors Associated with the Development and Progression of Periodontal Diseases and Pathologies for Patient <13 Years Old**

Factors	High risk	Moderate risk	Low risk
<b>Biological factors</b>			
Systemic conditions/genetic susceptibility (e.g., family history of aggressive periodontitis) and syndromes <sup>α</sup>	Yes		
Immunosuppressive or radiation therapy		Yes	
Medication(s) known to affect the periodontal tissues		Yes	
History of traumatic injury to the periodontal apparatus (e.g., avulsion, luxation)		Yes	
Traumatic gingival/oral mucosal lesions		Yes	
Nutritional deficiencies		Yes	
<b>Social and behavioral factors</b>			
Socioeconomic stability (e.g., adequate health literacy, regular dental care)			Yes
Adequate daily at-home oral hygiene either performed or supervised by caregiver			Yes
Tobacco or marijuana smoking/smokeless tobacco use	Yes		
<b>Clinical and radiographic factors</b>			
Adequate attached gingiva and normal frenum attachments			Yes
Tooth-related factors contributing to plaque retention		Yes	
Physical barriers for proper oral hygiene		Yes	
Generalized gingivitis (≥30% of teeth affected)		Yes	
Disproportional gingival inflammation in relation to age, amount of plaque accumulation, or oral and systemic developmental changes	Yes		
Presence of calculus	Subgingival	Supragingival	None
Bleeding on probing	Yes		
Periodontal probing depths >3 millimeter	Yes		
Chronic pericoronitis		Yes	
Abnormal tooth mobility	Yes		
Furcation involvement	Yes		
Radiographic alveolar bone loss	Yes		
Tooth loss due to periodontitis	Yes		

*Circling those conditions that apply to a specific patient helps the practitioner and caregiver understand the factors that contribute to the development and progression of periodontal diseases and pathologies. Clinical judgment may justify the use of one or more factors in determining the overall risk.*

Overall assessment of the patient's risk: High ☐ Moderate ☐ Low ☐

<sup>α</sup> Most common examples include, but are not limited to, agranulocytosis, Chédiak-Higashi syndrome, cyclic neutropenia, diabetes, Ehlers-Danlos syndrome, human immunodeficiency virus infection, hypophosphatasia, idiopathic immune disorders, Langerhans cell histiocytosis, leukemia, leukocyte adherence deficiency, osteoporosis, neutropenia, trisomy 21, Papillon Lefèvre syndrome, plasminogen deficiency, and respiratory diseases.

practices and to change behaviors regarding harmful risk factors (e.g., smoking, drug use). Psychological models and theories of motivation (e.g., health belief model, motivational interviewing, self-determination theory) may be used to help patients adopt healthier behaviors.<sup>31,32</sup>

### Recommendations

Dental professionals should utilize psychological theories of motivation to help patients adopt healthier behaviors and counsel their pediatric patients and parents on:

- the role of diet in the development and progression of periodontal conditions;
- the harms of all tobacco products to help prevent or cease tobacco use; and
- the serious health consequences of drug misuse, as well as refer to an appropriate provider for cessation when such a habit is identified.

**Table 3. Factors Associated with the Development and Progression of Periodontal Diseases and Pathologies for Patient ≥13 Years Old**

Factors	High risk	Moderate risk	Low risk
<b>Biological factors</b>			
Systemic conditions/genetic susceptibility (e.g., family history of aggressive periodontitis) and syndromes <sup>α</sup>	Yes		
Immunosuppressive or radiation therapy		Yes	
Medication(s) known to affect the periodontal tissues		Yes	
History of traumatic injury to the periodontal apparatus (e.g., avulsion, luxation)		Yes	
Traumatic gingival/oral mucosal lesions		Yes	
Nutritional deficiencies		Yes	
Mental health disorders (e.g., stress, depression)		Yes	
Pregnancy		Yes	
<b>Social and behavioral factors</b>			
Socioeconomic stability (e.g., adequate health literacy, regular dental care)			Yes
Adequate daily at-home oral hygiene			Yes
Tobacco or marijuana smoking/smokeless tobacco use	Yes		
Drug abuse (e.g., crack cocaine, methamphetamine)	Yes		
Intraoral/perioral piercing and oral jewelry/accessories		Yes	
Individuals with special health care needs living in supported community (group) homes		Yes	
<b>Clinical and radiographic factors</b>			
Adequate attached gingiva and normal frenum attachments			Yes
Adequate plaque biofilm control			Yes
Tooth-related factors contributing to plaque retention		Yes	
Physical barriers for proper oral hygiene		Yes	
Generalized gingivitis (≥30% of teeth affected)		Yes	
Disproportional gingival inflammation in relation to age, amount of plaque accumulation, or oral and systemic developmental changes	Yes		
Presence of calculus	Subgingival	Supragingival	None
Bleeding on probing (% of sites)	>25	10 to 25	0 to 9
Periodontal probing depths (mm)	>5	3.5 to 5	<3.5
Chronic pericoronitis		Yes	
Abnormal tooth mobility	Yes		
Furcation involvement	Yes		
Radiographic alveolar bone loss over 25% of sites	Yes		
Tooth loss due to periodontitis	Yes		

*Circling those conditions that apply to a specific patient helps the practitioner and caregiver understand the factors that contribute to the development and progression of periodontal diseases and pathologies. Clinical judgment may justify the use of one or more factors in determining the overall risk.*

Overall assessment of the patient's risk: High ☐ Moderate ☐ Low ☐

<sup>α</sup> Most common examples include, but are not limited to, agranulocytosis, Chédiak-Higashi syndrome, cyclic neutropenia, diabetes, Ehlers-Danlos syndrome, human immunodeficiency virus infection, hypophosphatasia, idiopathic immune disorders, Langerhans cell histiocytosis, leukemia, leukocyte adherence deficiency, osteoporosis, neutropenia, trisomy 21, Papillon Lefèvre syndrome, plasminogen deficiency, and respiratory diseases.

Table 4. Example of Management Pathways for Periodontal Diseases and Pathologies

Risk category	Diagnostics	Counseling						Nonsurgical therapy						Surgical therapy
		Twice daily brushing and daily flossing	Healthy diet and nutrition	Injury <sup>α</sup> prevention	Tobacco use and drug misuse <sup>β</sup>	Use of oral hygiene adjuncts <sup>γ</sup>	Compliance with medical care and/or periodontal treatment or maintenance	Oral prophylaxis: supragingival plaque and calculus removal	Debridement, scaling and root planing	Systemic antibiotics and/or use of adjunctive topical anti-microbials	Management of plaque retentive factors <sup>α</sup>	Monitor previous traumatic injuries to the periodontal apparatus	Management of oral conditions and side effects from therapies, medications, infections, gingival injuries, etc.	
Low risk	<ul style="list-style-type: none"> <li>– Recall every six to 12 months</li> <li>– Radiographs every 12 to 24 months</li> </ul>	Yes	Yes	Yes	Prevention			Every six to 12 months						
Moderate risk	<ul style="list-style-type: none"> <li>– Recall every six months</li> <li>– Radiographs every six to 12 months</li> <li>– Monitoring of systemic conditions by laboratory analysis and consultation with medical specialists, if indicated</li> </ul>	Yes	Yes	Yes	Prevention or cessation	Yes	Yes	Every six months	Every six months	Yes	Yes	Yes	Yes	Yes
High risk	<ul style="list-style-type: none"> <li>– Recall every three months</li> <li>– Radiographs every six months</li> </ul>	Yes	Yes	Yes	Prevention or cessation	Yes	Yes	Every two-four months depending on disease severity and disease response to treatment	Every two-four months depending on disease severity and disease response to treatment	Yes	Yes	Yes	Yes	Yes

<sup>α</sup> Plaque retentive factors include, but are not limited to, caries lesions, enamel defects, dental anatomical anomalies, malposed teeth, defective restorations, inadequate contoured crowns, orthodontic appliances, dental prostheses.

<sup>β</sup> Prevention of injuries resultant of accidents, piercings, habits.

<sup>γ</sup> Oral hygiene adjuncts include, but are not limited to, powered toothbrushes, interdental brushes, or oral irrigation; chemical antiplaque and anticalculus agents.



### Nonsurgical periodontal therapy (Phase I)

The primary goal of phase I therapy is to control the factors responsible for periodontal inflammation; this includes patient education focused on the role of adequate home care in facilitating the removal of bacterial plaque biofilm. In addition to plaque biofilm and calculus, other local factors can contribute to plaque biofilm retention and physical barriers for proper oral hygiene execution, thereby increasing periodontal disease risk and pathology initiation.<sup>33-38(pp560,561)</sup> As such, Phase I therapy includes oral hygiene instruction, oral prophylaxis or scaling and root planing (SRP), and other therapies such as caries control, replacement of defective restorations, occlusal therapy, orthodontic tooth movement, and cessation of confounding habits such as tobacco use.<sup>38(pp560,561)</sup>

#### *Management of bacterial plaque biofilm and calculus*

Oral prophylaxis and SRP are the basis of professional mechanical plaque biofilm control.<sup>23,37,38(pg561),39,40</sup> Supra- and subgingival instrumentation is an important component of initial and recall dental appointments. Oral prophylaxis removes supragingival plaque biofilm and calculus via hand or powered instruments. Subgingival instrumentation is divided into three treatment goals: (1) debridement (removal of subgingival plaque biofilm); (2) scaling (removal of supra- and subgingival plaque biofilm, calculus, and stains); and (3) root planing (removal of cementum or surface dentin that is rough, impregnated with calculus, or contaminated with toxins or microorganisms).<sup>38(pp560-563)</sup> When comparing subgingival instrumentation modes, hand instruments (e.g., curettes) remove a significantly greater amount of calculus and leave a smoother root surface than ultrasonic scalers.<sup>39</sup> On the other hand, ultrasonic devices cause less soft-tissue trauma, require a shorter treatment time, and are less technique/operator sensitive.<sup>39</sup> However, systematic reviews have found that treatment results are similar between hand instruments and ultrasonic devices, with many clinicians using a combination.<sup>41</sup> Therefore, clinician and patient preference along with anatomic characteristics of teeth, ease of access, and time required to perform treatment are factors to be considered.

#### Recommendations:

- Dental professionals should provide oral self-care instructions that are individualized and include appropriate adjuncts to facilitate efficiency and effectiveness of home care techniques.
- For adolescents and individuals with SHCN who exhibit poor oral hygiene, clinicians should consider the use of chemical antiplaque agents in mouthrinses or incorporated into fluoridated toothpastes to control plaque biofilm accumulation and gingival inflammation, along with instituting more frequent recall appointments.
- Because plaque biofilm and calculus serve as physical barriers for proper home oral hygiene execution, a the indicated nonsurgical therapy (i.e., dental prophylaxis, SRP) should be performed at both initial and recall dental appointments.

- Use of ultrasonic devices and mouthrinses may be contraindicated for patients who are unable to expectorate and at risk for aspiration.

#### *Management of local factors for periodontal disease and pathologies*

Local factors that may increase periodontal disease risk include caries lesions, defective restorations, malocclusion, orthodontic appliances, and dental enamel defects as well as other dental anomalies.

#### Recommendations:

Dental professionals should utilize psychological theories of motivation to help patients adopt healthier behaviors and counsel their pediatric patients and parents on:

- Clinicians should consider restoring open, arrested cavitated lesions when food impaction causes gingival inflammation, bleeding, or patient discomfort.
- Defective or failing restorations should be corrected by smoothing rough surfaces, removing overhangs with burs and/or hand instruments, or replacement.<sup>37,38(pg561)</sup>
- When placing preformed crowns, well-adapted restorations (i.e., contoured, well-fitted, and crimped) are recommended to maintain the health of the periodontium.
- Because orthodontic appliances often hinder brushing and flossing, clinicians should:
  - consider more frequent recall appointments and prophylaxis depending on home oral hygiene compliance and degree of periodontal inflammation, and
  - consider suspension of orthodontic treatment if the patient is not able to maintain proper oral hygiene.
- In cases of sensitivity associated with dental defects including dentinogenesis and amelogenesis imperfecta or molar-incisor hypomineralization, the use of desensitizing toothpastes, fluoride varnishes, toothbrushes with soft bristles, and sealants for the affected enamel of the teeth should be considered as decreased sensitivity may improve home oral hygiene.

#### *Topical antimicrobial adjuncts and systemic antibiotics*

Topical (local) agents, available as fibers, gels, chips, microspheres, and solutions, are delivered directly inside the periodontal pocket and present fewer side effects than systemic agents.<sup>39,42-44</sup>

Compared to systemic agents, they utilize a smaller total dosage and provide higher localized concentration of the drug, but lack the capability to reach different oral surfaces and saliva.<sup>39,42-44</sup> Although systematic reviews have reported that adjunctive local antibiotics improve PPD and clinical attachment loss (CAL) in short-term studies and PPD in long-term studies, their use is controversial due to high cost and small magnitude of clinically-relevant benefits.<sup>43,44</sup> Local antibiotic therapies have been used more commonly during the maintenance phase to treat remaining and isolated recurrent pockets.<sup>43</sup>

Systemic antibiotics are indicated when patients exhibit moderate periodontitis with three to four mm of CAL and PPD of less than five mm.<sup>45</sup> Younger patients with periodontitis

characterized by rapid attachment and bone loss<sup>39,44,46-48</sup>, patients with necrotizing periodontitis<sup>47,48</sup>, and those with periodontitis as a manifestation of systemic conditions<sup>39,48-51</sup> may benefit significantly from adjunctive antibiotic therapies in combination with SRP. Disadvantages of systemic administration include adverse drug effects (e.g., gastrointestinal symptoms, allergic reaction), poor patient compliance, and, very importantly, development of bacterial resistance due to indiscriminate use.<sup>44,46</sup> Several factors (e.g., patient's clinical parameters, health history, dental history, drug allergy, medication compliance, personal/parental preferences, adverse effects, bacterial resistance, treatment response in primary versus permanent dentitions) influence the decision to use topical or systemic antibiotic adjuncts to SRP.<sup>42,52-55</sup>

Several combinations of systemic antibiotics have been suggested in the literature.<sup>44,56</sup> When compared to SRP alone, the combination of amoxicillin and metronidazole (and, to a lesser degree, azithromycin and metronidazole) as an adjunctive therapy has shown to reduce the number of major periodontopathogenic bacteria, significantly improve CAL gain, and promote higher percentage of pocket closure, as well as reduce BoP, PPD, and frequency of pockets of greater than four mm.<sup>43-46,52,54,57-60</sup> Regimen durations of one to two weeks have been cited in the literature with respective advantages and disadvantages.<sup>39,52</sup> For patients allergic to penicillin, antibiotic regimen using metronidazole alone is an alternative treatment.<sup>58</sup> Additionally, azithromycin is effective against periodontal pathogens with positive immunomodulatory properties and has been proven effective in treating aggressive periodontitis in young patients<sup>61</sup> as well as adults<sup>62</sup>. Azithromycin is one of the safest antibiotics for patients allergic to penicillins, but there are risks of cardiac complications including cardiotoxicity.<sup>63,64</sup> Cardiac risk in pediatric patients seems to be due to an increased risk of QT prolongation associated with higher dosage levels,<sup>65</sup> and caution should be exercised in patients with cardiac risk factors. The recommended antibiotic dosage for periodontal treatment differs from that used for odontogenic infections or endocarditis prophylaxis.<sup>66</sup>

#### Recommendations

- Stand-alone antibiotic therapy is not recommended in the treatment of periodontal disease.
- Adjunctive antibiotic therapy to SRP should be considered for patients with advanced or aggressive periodontal disease.
- When adjunctive antibiotic therapy to SRP is indicated, the decision to use topical or systemic antibiotics should be carefully evaluated and based on patient's general health status, periodontal disease severity, compliance, and response to SRP.

#### *Re-evaluation (determining success or lack of success of nonsurgical therapy)*

After procedures of phase I (e.g., debridement, scaling, root planing, caries control, correction of defective restorations) are completed, the periodontal tissues will go through a process

of healing that may take four or more weeks to occur.<sup>38(pg562)</sup> Transient tissue sensitivity is often observed during the healing process and usually diminished with good home plaque biofilm control.<sup>38(pg562)</sup> Re-evaluation findings help determine the need for any further nonsurgical therapy procedure or periodontal surgery.<sup>38(pg563)</sup>

#### Recommendations:

- Components of re-evaluation appointments should include probing the periodontal tissues, examining all related anatomic structures, reinforcing home care regimens, and discussing existing harmful habits with a goal of cessation.
- The frequency of supportive periodontal therapy must be individualized and based on the patient's symptoms, clinical and radiographic findings, risk factors, initial severity of the disease, as well as residual diseased sites at the end of the active periodontal treatment in relation to the patient's age, treatment outcome, caries risk, and plaque or biofilm control.

### **Periodontitis as a manifestation of systemic disease (systemic phase)**

#### *General considerations*

Multiple systemic diseases including diabetes, connective tissue disorders, metabolic and endocrine disorders, and immunological disorders as well as medications can have an adverse effect on the periodontium.

#### Recommendations:

- Clinicians should consider systemic diseases and medications that can affect the periodontal attachment apparatus or the course of periodontal diseases in order to achieve accurate diagnoses and plan treatment.<sup>49,67</sup>
- Consultation with the patient's medical care provider may be necessary for management of at-risk patients.<sup>49,67</sup>

#### *Special management considerations*

**Halitosis.** Halitosis (breath odor or bad breath) is the result of anaerobic bacterial breakdown of organic substrate. A coated tongue accounts for the majority of cases of bad breath.<sup>68,69(pp441,443)</sup> Halitosis may be caused by a variety of physiologic or pathologic processes. Physiologic factors may include food intake, smoking, or decreased salivary function. Additionally, halitosis may be associated with gastroesophageal disorders, mouth breathing<sup>70</sup>, nasal obstructions including sinusitis, foreign bodies, or tonsil stones<sup>69(443),71</sup>. Persistent halitosis typically reflects a pathologic process such as periodontal disease, pericoronitis, recurrent major aphthous ulcers infected with gram-negative anaerobic bacteria, herpetic gingivostomatitis, or necrotizing gingivitis-periodontitis.

#### Recommendations:

- Treatment of halitosis depends on its cause and involves reduction of substrates for bacteria, chemical reduction of bacterial load, and masking odor.<sup>69(pg449)</sup> Strategies for treatment include tongue scraping or cleaning prior to tooth brushing, antimicrobials such as chlorhexidine



with/without scaling and root planing, and the use of mints or chewing gum to mask odor.<sup>69(pp449,450,452)</sup>

**Respiratory diseases affecting the periodontium.** Health of the periodontium depends on saliva's mechanical cleansing and antimicrobial properties. Respiratory diseases, either directly (e.g., mouth breathing) or through side effects (e.g., xerostomia) of therapeutic agents, may alter salivary flow.<sup>72,73</sup> Nasopharyngeal obstruction from adenoid and tonsillar hypertrophy, as well as significant neuromuscular weakness with a history of snoring, can also affect periodontal health.<sup>72</sup> Depending on the individual oral/dental needs of patients with respiratory diseases, the pediatric dentist plays an important role in early diagnosis of general and oral health problems associated with respiratory diseases, care management, and establishment of a multidisciplinary approach that may include, but is not limited to, orthodontists, primary care providers, otolaryngologists, and speech pathologists.<sup>72</sup> Regular dental check-ups with oral hygiene instructions for proper home plaque biofilm control, mouth rinsing after medications, and use of fluoridated toothpaste are important preventive regimens to reduce the risk of periodontal disease and dental caries among patients with respiratory diseases.<sup>72</sup>

#### Recommendations:

- Clinicians should carefully evaluate the patient's health history and medications in order to identify respiratory conditions and medications that impact salivary flow and dental and periodontal health.
- If airway obstruction is determined to affect periodontal health, an evaluation by an otolaryngologist is recommended.
- Clinicians should consider a multidisciplinary approach, referral, and/or care coordination for patients with general and/or oral health problems associated with respiratory diseases.

#### *Oral conditions related to immunosuppressive or radiation therapies*

Patients undergoing immunosuppressive or radiation therapies may present with periodontal problems associated with treatment. Gingival bleeding, soft-tissue necrosis, salivary gland dysfunction, opportunistic infections (e.g., candidiasis, herpes simplex virus), and oral graft-versus-host disease are among the many acute and long-term complications associated with these therapies.<sup>74-77(pp69,72,73)</sup> Special attention should be given to partially-erupted molars that may be at risk for pericoronitis.<sup>76,77(pg71)</sup> When definitive periodontal therapy cannot be rendered, extraction of hopeless periodontally-involved teeth is the treatment of choice.<sup>75-77(pg71)</sup> A periodontal assessment and appropriate therapy are indicated before patients undergoing cancer treatment receive bisphosphonates.<sup>75</sup> Refer to AAPD's *Dental Management of Pediatric Patients Receiving Immunosuppressive Therapy and/or Head and Neck Radiation*.<sup>75</sup> for additional information on managing periodontal considerations in these circumstances.

#### Recommendation:

- Clinicians should work closely with the patient and parents, as well as with his multidisciplinary health care team, to ensure that any medically-necessary dental treatment is integrated, coordinated, and delivered in a timely and safe manner before, during, and after immunosuppression or radiation therapy.<sup>74</sup>

**Drug-influenced gingival enlargements.** Drug-influenced gingival enlargements have been associated with three types of medications: anticonvulsants (e.g., phenytoin, sodium valproate), calcium channel blockers (e.g., verapamil, diltiazem), and immunosuppressants (e.g., cyclosporine).<sup>21,78</sup> In most cases, the gingival enlargement is induced by the combination of the drugs (i.e., fibrotic aspect) and the plaque biofilm (i.e., inflammatory aspect).<sup>79(pp269-71)</sup> Treatment options may include: (1) possible drug discontinuation or change; (2) biofilm control by means of home oral hygiene, use of antimicrobial agents (e.g., chlorhexidine), frequent professional cleaning and SRP, removal of plaque-retentive areas (e.g., faulty restorations); and (3) surgical removal of enlarged gingiva (e.g., gingivectomy using a scalpel or laser-assisted therapy, flap surgery, or electrosurgery).<sup>35(pp293,294),78,79(pg274)</sup> and may include possible drug discontinuation or change.

Periodontal flap surgery to manage gingival enlargements is favored over gingivectomy in terms of minimizing the amount of tissue and time recurrences.<sup>80(pp734-737)</sup> However, in general, gingivectomy is indicated for small areas of gingival enlargement (i.e., up to six teeth) where there is no evidence of CAL or the need for osseous surgery, while flap surgery is indicated for larger areas (i.e., more than six teeth) with evidence of CAL or the need for osseous surgery.<sup>80(pg737)</sup> Antibiotic therapy as an adjunctive anti-microbial and anti-inflammatory agent has been proposed as another step in the management of gingival enlargements.<sup>35(pg294),78</sup>

#### Recommendations:

- Biofilm control, SRP, and timely evaluation of the initial treatment response should occur before considering surgical therapy.

#### Other periodontal conditions

##### *Oral soft-tissue and tooth-supporting structure injuries*

Orofacial trauma can result in extraoral and intraoral soft-tissue injuries such as lacerations, contusions, abrasions, and avulsions.<sup>81,82</sup> Traumatic dental injuries almost always involve the periodontal tissues which may undergo ischemia, crushing, or loss.<sup>23,36</sup> Injuries to the periodontal ligament (PDL) may range from minor lacerations with dental concussion, tearing of the fibers with subluxation, to partial or complete separation with luxation or avulsion, and loosening of the tooth can occur.<sup>83,84</sup> When foreign bodies (e.g., gravel, tooth fragment) may be embedded within the injured soft tissues, clinical inspection is supplemented by a soft-tissue radiograph.<sup>81</sup> Removal of foreign bodies is necessary to avoid tissue infection, scarring, or tattooing.<sup>85(pg626),86</sup> Cleansing, debridement, hemostasis, and

closure are the major steps in managing soft-tissue injuries with the goals to maintain tissue vascularity, enhance healing, and prevent tissue devitalization, as well as to minimize the risk of gingival recession and bone/root exposure.<sup>86</sup> Reapproximated soft-tissue wounds are sutured using the minimal amount of small-diameter sutures.<sup>85(pp642),86</sup> Because determining which wounds are tetanus prone is not possible, need for tetanus prophylaxis is based on the patient's current immunization status.<sup>87</sup> A decision for antibiotic prophylaxis is based on the severity and contamination status of the tissue injury.<sup>88</sup>

Splinting stabilizes traumatized teeth with the goals to optimize PDL reattachment and healing and to protect the teeth against further insult.<sup>89,90</sup> Characteristics of an ideal splint for mobile traumatized teeth include being passive, flexible, and non-irritating to surrounding soft tissues as well as allowing for physiological tooth mobility and proper oral hygiene.<sup>89,90</sup> Alveolar bone fractures require a more rigid splint with longer splinting time.<sup>91</sup>

The risk of PDL healing complications is very low for concussion, subluxation, and extrusive and lateral luxation injuries and significantly more for traumatic dental injuries involving multiple teeth and teeth with full root development.<sup>83,84</sup> The most common complications are "repair-related resorption (surface resorption), infection-related resorption (inflammatory resorption), ankylosis-related resorption (replacement resorption), marginal bone loss, and tooth loss".<sup>84</sup> Ankylosis-related root resorption is an expected outcome in replanted teeth, especially with an extra-alveolar dry time longer than 60 minutes or transport medium other than one capable of maximizing the vitality of the PDL cells (e.g., milk, Hanks' Balanced Salt Solution).<sup>91,92</sup>

#### Recommendations:

- Management of orofacial soft-tissue injuries should include cleansing, debridement, establishing hemostasis, and closure of wounds in a manner that maintains tissue vascularity, enhances healing, and prevents tissue devitalization.
- The clinician should determine the need for tetanus prophylaxis based on the patient's current immunization status. When immunization status is in doubt, evaluation by a physician within 48 hours is indicated.<sup>85(pp643),88,91</sup>
- A decision for antibiotic prophylaxis should be based on the severity and contamination status of the tissue injury.<sup>85(pp642,643),88</sup> Because the PDL of an avulsed tooth may have been contaminated by oral or environmental bacteria, systemic prophylactic antibiotics are recommended following tooth replantation.<sup>91</sup>
- Depending on the extent of the injury suffered by the periodontium, collaboration between the primary care dentist and a periodontologist may be needed to allow effective and successful clinical outcomes following dentoalveolar trauma.

#### *Infections of bacterial, fungal, and viral origins*

The gingiva may demonstrate a variety of lesions that are not caused by plaque biofilm and usually do not resolve after plaque biofilm removal.<sup>3</sup> Infections of bacterial (e.g., necrotizing gingivitis), fungal (e.g., candidiasis), and viral (e.g., primary herpetic gingivostomatitis, recurrent intraoral herpes simplex infection) origins are some examples of nonplaque-induced gingival lesions observed in the pediatric population.<sup>35(pp286,287)</sup> Elimination or reduction of all local and systemic risk factors that contribute to the infection initiation or progression is needed for treatment completeness, followed by close monitoring to assess treatment effectiveness, patient compliance, and risk of recurrence.

#### Recommendations:

- Initial therapy should focus on alleviating acute symptoms of pain and distress. This could include oral analgesics to control fever, malaise, and pain, as well as fluids to prevent dehydration.
- Antimicrobial therapy should be considered when an infection is not self-limiting or if there are frequent recurrences.

#### *Traumatic gingival and oral mucosa lesions*

Traumatic lesions can be accidental, iatrogenic, or self-inflicted and are physical (e.g., oral piercing, aggressive toothbrushing), chemical (e.g., dental materials, topical cocaine), or thermal (e.g., overheated foods and drinks) in nature.<sup>93,94</sup> The appearance of the lesion (e.g., acute ulcerations vs chronic gingival defects) and a detailed history are crucial in achieving a diagnosis. Self-injury behavior (SIB) has been reported among individuals with psychiatric illnesses (e.g., personality disorders, bipolar disorder, major depression, anxiety disorders, obsessive-compulsive disorder) and congenital insensitivity to pain (e.g., familial dysautonomia), as well as a variety of developmental and intellectual disabilities (e.g., autism).<sup>95</sup> Gingival picking/scratching is among the most common oral SIB.<sup>94-99</sup> Management of self-inflicted traumatic lesions may be complicated due to lack of patient's compliance. The patient's primary care provider may help rule out any medical reasons for SIB (e.g., otitis media, infection, pneumonia) or specific genetic disorders (e.g., Lesch-Nyhan syndrome) or determine comorbid psychiatric conditions. An approach that includes medical and behavioral specialists may be indicated. Periodontal plastic surgery (e.g., placing a graft to create or widen the attached keratinized tissue)<sup>100(pp683),101(pp775)</sup> may be necessary for permanent gingival defects.<sup>94,95,97</sup>

#### Recommendations:

- Management of traumatic oral lesions requires assessment of the etiologic factor, removal of the offending agent, and symptomatic therapy.
- Treatment of SIB should be individualized and may include behavior modification, pharmacotherapy, immobilization devices, oral appliances to control harmful habits, and/or psychological or psychiatric.<sup>95,98</sup>

- Re-evaluation and monitoring management approaches should occur while treating self-inflicted traumatic lesions.

### *Pericoronitis*

Pericoronitis refers to an inflammatory lesion developed when food debris and bacteria are present beneath the excess flap of soft tissue surrounding partially-erupted teeth, most frequently involving mandibular third molars.<sup>102</sup> The pericoronal flap of soft tissue may be chronic without any symptoms; however, when acute, patients may experience severe pain, mouth opening restriction, gingival abscess, cellulitis, fever, lymphadenopathy, and presence or risk for systemic complications.<sup>103</sup> A rare complication is Ludwig's angina, a life-threatening condition that occurs when infection spreads to submandibular, sublingual, and submental spaces thereby compromising the patient's airway.<sup>103</sup> The first course of treatment for acute pericoronitis is management of infection and pain.<sup>102,103</sup> Nonsteroidal anti-inflammatory drugs (NSAIDs) are the analgesics of choice since the control of inflammation helps to control acute pain.<sup>104</sup> Patient compliance for home oral hygiene is also key for treatment success.<sup>103</sup> Once acute symptoms resolve, decisions can be made regarding the need for further treatment (e.g., pericoronal tissue surgery or tooth extraction).<sup>102,103</sup>

#### Recommendations:

- Management during the acute phase should consist of<sup>102,103</sup>:
  - debridement and irrigation of the pericoronal area,
  - drainage of purulence to relieve pressure,
  - occlusion evaluation to determine the need to reduce soft tissue or adjust occlusion of opposing tooth,
  - pain control using NSAIDs,
  - antibiotics if the infection is not localized or with systemic signs and symptoms, and
  - home care plan to include oral cleaning, warm saline rinses, antiseptic agents (e.g., chlorhexidine), and sufficient fluid intake.
- After the acute phase, practitioners should<sup>102,103</sup> evaluate prognosis and likelihood that the tooth involved will either erupt without complications or continue to pose a risk for pericoronitis recurrence and decide to either remove the pericoronal flap (if not removed during the acute phase) or extract the tooth to prevent recurrence.
- Ludwig's angina requires early recognition, immediate intervention (e.g., early and aggressive antibiotic therapy, surgical drainage, nutrition, hydration), and close monitoring. Due to the threat of rapid airway compromise, emergency referral to an otolaryngologist or an oral and maxillofacial surgeon should occur without delay.<sup>105</sup>

### **Considerations for treatment coordination and/or referral of care with a periodontist**

Most pediatric patients will attain periodontal disease control with nonsurgical therapy and not require further surgical

intervention. When PPD are greater than five mm, referral to a periodontal specialist may be indicated. Periodontal surgery may improve tooth support through pocket reduction, bone augmentation, and regeneration procedures.<sup>46</sup> Other considerations for referral include: (1) extent of the disease (generalized or localized periodontal involvement); (2) presence of short-rooted teeth; (3) teeth hypermobility; (4) difficulty in SRP deep pockets and furcations; (5) possibility of damage to the developing permanent successor tooth; (6) restorability and importance of particular teeth for reconstruction; (7) lack of resolution of inflammation after thorough plaque or biofilm removal and excellent SRP; (8) presence of systemic diseases and other conditions that compromise the host response; and (9) very importantly for the pediatric population, the age of the patient.<sup>38(p563)</sup> Younger patients, both systemically healthy and compromised, with extensive CAL are more likely to have aggressive forms of periodontitis that can be rapidly destructive necessitating timely advanced therapy. Early loss of primary teeth and bone loss visible on posterior bitewing radiographs are important indicators of aggressive forms of periodontitis that require further follow-up and/or referral.<sup>106</sup> The possibility of an underlying systemic disease cannot be discarded.

The treatment for periodontitis as a manifestation of systemic conditions is dependent on the systemic disorder. Two fundamental treatment differences exist: (1) patients for whom the systemic disease and a conservative periodontal treatment approach do not represent grave danger to life; and (2) patients for whom the systemic disease (e.g., hypophosphatasia, leukocyte adhesion deficiency syndrome, neutropenia) and a conservative periodontal treatment approach may represent grave danger to life. Managing the periodontal diseases in these children, even when extractions of primary teeth at an early age is the treatment of choice, is crucial since such systemic diseases may endanger the children's lives.<sup>107-110</sup>

In terms of coordination and referral of care with a periodontist, important considerations include<sup>110,111</sup>:

- the primary care dentist will be working closely with the medical team, and all pertinent patient information needs to be available to the periodontist to determine the necessity of advanced periodontal therapies;
- the level and frequency of communication between the primary care dentist and the periodontist will be more than is required for healthy patients. Timely communication before and after each diagnostic and surgical appointment is essential; and
- the types and levels of behavioral and pharmacologic pain and anxiety control available in the periodontal office may not be ideal for the young patient. Seeing the patient together may help meet these needs.

#### *Recommendations*

- The treatment of periodontitis as a manifestation of systemic disease where a conservative periodontal treatment approach may represent grave danger to the child's

life should include communication with the pediatrician or medical specialist, as well as a periodontist, to consider the risk and benefit of conservative periodontal treatment versus tooth extractions. Extraction may be the best treatment with a continuing periodontal infection causing severe destruction of bone and developing permanent teeth and endangering the child's life.

- The treatment of periodontitis as a manifestation of systemic disease where a conservative periodontal treatment approach does not represent grave danger to the child's life should include communication with the child's pediatrician or medical specialist about the systemic condition, its diagnosis based on the oral, laboratory, and systemic findings, as well as coordination of systemic and periodontal treatments.
- Treatment should include the provision of the full scope of preventive interventions and phase I treatment, with consideration for surgical intervention if nonsurgical management fails.

### **Surgical therapy (Phase II)**

Periodontal surgical therapy is indicated when nonsurgical means cannot adequately control disease or address mucogingival and periodontal defects. Surgical modalities include esthetic, resective, regenerative, preprosthetic, and implant procedures and are utilized with the objective to stabilize the periodontal prognosis, return periodontal and dental structures lost to disease, and improve esthetics.<sup>100(pg683)</sup> During this phase, the role of the primary care dentist is to provide treatment or refer/coordinate the care with a periodontal or oral surgery specialist when the needed treatment exceeds the practitioner's scope of practice. Prior to any surgical therapy, clinicians should provide the patient an opportunity to have questions answered and obtain written informed consent to proceed with the therapy proposed. Following are some surgical therapy considerations.

#### *Pocket reduction surgery*

**Gingivectomy.** The indication for gingivectomy in the treatment of periodontal disease is to remove the soft tissue of the pocket wall in order to create visibility and access for thorough SRP. In combination with gingivoplasty (i.e., recontouring of the gingiva), gingivectomy can achieve a favorable environment for soft-tissue healing and physiological gingival contour.<sup>80(pg738),112</sup> Due to secondary wound closure, gingivectomy procedures may cause more postoperative discomfort and bleeding when compared to periodontal flap surgeries.<sup>80(pg730)</sup> In modern periodontics, gingivectomy is utilized less often for the treatment of periodontal disease.<sup>80(pg738)</sup> However, it remains beneficial in the treatment of gingival enlargements and suprabony pockets when the pocket wall is firm and fibrous.<sup>80(pg738),112</sup> Gingivectomy is not indicated in cases for which the risk of postoperative bleeding is increased, access to bone is required, the zone of keratinized tissue is narrow, or aesthetics are a concern.<sup>80(pg738)</sup>

**Flap surgery.** Periodontal flap surgery, the most widely used procedure for pocket therapy, provides great access for SRP, periodontal regeneration, and gingival and osseous resection.<sup>80(pg739)</sup> With the exception of anterior maxillary sites where compromise to the esthetics may outweigh the gains, surgical access is the preferred treatment approach for moderate and deep pockets.<sup>100(pg687)</sup>

#### *Regenerative surgery*

Periodontal regeneration aims to restore the lost periodontal tissues and their respective functions by the formation of new alveolar bone, cementum, and PDL.<sup>113-116(pg756),117</sup> Around hopeless teeth, regenerative therapy has been reported to be less costly when compared to extractions and dental implants.<sup>118</sup> Several regeneration therapies including guided tissue regeneration and bone grafts (e.g., autogenous, allogenic, xenogenic, synthetic or alloplastic) have been studied and are available to the clinician.<sup>114-116(pg760),119</sup>

Systematic and meta-analysis reviews have shown periodontal regeneration in intrabony defects results in shallower residual PPD and greater CAL gain than flap surgery alone.<sup>114,119</sup> Disadvantages of regenerative therapies include their technically-demanding surgical procedures as well as dependence on patients' compliance with home oral hygiene and professional maintenance care.<sup>114,116(pg760),119</sup>

#### *Laser therapy*

Lasers have been used successfully in several periodontal therapies such as gingivectomy/gingivoplasty, frenectomy, crown lengthening and exposure, depigmentation, periodontal regeneration, and management of excess tissue in gummy smile and pericoronitis.<sup>116(pg757),120,121</sup> Advantages associated with laser use include improved perioperative visualization from hemostasis, reduced need for sutures, wound detoxification, enhanced healing, better patient acceptance, and postoperative pain control.<sup>122-125(pg771)</sup> The greatest risk associated with lasers is unintentional tissue necrosis due to excessive temperatures.<sup>125(pg770)</sup> For periodontal disease treatment, laser-assisted new attachment procedure (LANAP) has shown to initiate regeneration and improve clinical outcomes in the nonsurgical treatment of moderate to advanced periodontitis, as either a monotherapy or as an adjunct to SRP<sup>116(pg757),120,124</sup>, due to its benefits of detoxification, calculus removal, minimally invasive access for SRP, and killing of periodontal pathogens.<sup>123-125(pg771)</sup> However, more data is needed to support the use of lasers as adjuncts to resective and regenerative therapies.<sup>116(pg756),124</sup>

#### *Extraction of periodontally-compromised teeth*

Extraction of periodontally-compromised teeth may be the best management for some patients. Important considerations include previous unsuccessful therapies, dental implants as an alternative, cost-effectiveness of periodontal procedures, as well as the patient's systemic health, compliance, and finances.<sup>114-116(pg765),118</sup> For pediatric patients, extraction of primary teeth may be indicated if the periodontal lesion approximates



the developing permanent successor, endangering the dental development.

### *Esthetic surgery*

Esthetic surgery includes a subset of periodontal procedures directed towards, but not limited to, correcting smile esthetics (e.g., esthetic crown lengthening, soft-tissue augmentation). Although some of these procedures are inappropriate for growing patients, pediatric dentists may find themselves providing guidance on these common parental concerns.<sup>101(pg776)</sup>

Esthetic crown lengthening is a surgical procedure that combines gingivectomy and osseous recontouring techniques to address complaints of a gummy smile when altered passive eruption is the primary etiologic factor for the excessive gingival display.<sup>126</sup> This procedure is best completed after the patient has completed facial growth. Success depends on a thorough diagnosis that effectively evaluates the impact of other etiologic factors such as vertical maxillary excess or microdontia.

Soft-tissue augmentation procedures, sometimes referred to as mucogingival surgery or periodontal plastic surgery<sup>127</sup>, include connective tissue grafts, primarily for the treatment of gingival recession, and free gingival grafts for areas lacking adequate keratinized gingiva. Typically, these procedures are performed on adult patients, but consideration may be given to the use on a pediatric patient undergoing orthodontic movement who is at increased risk for worsening mucogingival deformities.<sup>101(pg777)</sup>

### *Dental implants for missing teeth*

In general, conservative treatment is indicated for growing patients with missing teeth. The placement of dental implants in younger patients typically is contraindicated before age 18 for women and 21 for men due to the expected continued skeletal growth in these patients.<sup>128</sup> When indicated, implant treatment requires a carefully coordinated and multidisciplinary team approach. Important considerations include:

- the number of missing teeth along with soft and hard tissue anatomy,
- growth and development,
- systemic conditions and psychological and behavioral maturity<sup>129</sup>, and
- alternative therapies such as orthodontic and prosthetic treatments.

Assessment of growth and development is key to successful outcomes for endosseous dental implants in pediatric patients. Early placement of implants in the growing patient can result in rotation of the dental implant and infraocclusion as the adjacent teeth continue to erupt and the jaw grows.<sup>129</sup> Since patients vary considerably in their growth patterns, individual patients may have periods of rapid and slower growth.<sup>130</sup> Thus, chronological age is not a good indicator of completion of growth. In contrast, skeletal maturation, assessed by cephalometric analysis or hand-wrist radiographs, is preferred,<sup>131</sup> especially for definitive placement of endosseous implants

within the aesthetic zone. However, limited studies have suggested the use of extra-narrow implants may be used as temporary restorations in late adolescence when the patient is older than 15 years.<sup>132</sup>

### *Recommendations*

- If PPD inhibits subgingival access or anatomic/morphologic defects require correction, the clinician should inform the patient of the need for and benefits/risks of periodontal surgical therapy, as well as treatment alternatives.
- Extraction of periodontally-compromised teeth may be the best management for some patients.
- Clinicians should consider referral to a specialist when the surgical interventions are beyond their scope of practice.
- Determination for advisability and timing of implant placement must be based on the specific circumstances of the individual patient. The patient's stage of growth and development is critical to treatment success.

### **Maintenance phase**

The long-term success of periodontal therapy outcomes is highly associated with the quality of recall maintenance.<sup>39,133</sup> Following are some considerations of the maintenance therapy phase:

- determination of recall procedures (i.e., prophylaxis, periodontal maintenance).
- determination of recall interval based on risk factors and history of disease.
- use of antimicrobial adjuncts during maintenance.
- individualized home care reinforcement.
- decision to when re-enter phase I or phase II therapy.

A classic study assessing the efficacy of a maintenance care program demonstrated that patients placed on a three-month recall maintained excellent oral hygiene parameters and stable periodontal attachment levels for two to six years following periodontal therapy, while the nonrecall control group demonstrated significant periodontal attachment loss.<sup>134</sup> A 30-year outcome report of this same study population further demonstrated that patients placed on an individualized maintenance program with a three- to 12-month recall interval maintained stable periodontal conditions for 30 years.<sup>135</sup> A review assessing predefined periodontal recall intervals conducive to periodontal health and stability concluded that evidence supports a two-to-four-month recall interval for patients affected by moderate to advanced periodontal disease.<sup>133</sup> Moreover, evidence supports a maintenance therapy program with at least 12-month interval recalls for patients who are periodontally healthy, are periodontally stable, or have mild forms of periodontitis.<sup>133</sup>

*Recommendations*

- Clinicians should educate their patients and caregivers about the importance of supportive periodontal therapy to prevent disease relapse and provide individualized periodontal supportive care when needed.
- Every two- to four-months and at least every 12-months recall intervals are recommended for patients with higher and lower periodontal disease risk, respectively.

**References**

1. American Academy of Pediatric Dentistry. Risk assessment and management of periodontal and pathologies in pediatric dental patients. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2022:466-84.
2. American Academy of Pediatric Dentistry. Classification of periodontal disease in infants, children, adolescents, and individuals with special health care needs. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2019:387-401.
3. Caton JG, Armitage G, Berglundh T, et al. A new classification scheme for periodontal and peri-implant diseases and conditions—Introduction and key changes from the 1999 classification. *J Periodontol* 2018;89(Suppl 1): S1-S8.
4. Stenberg WV. Periodontal problems in children and adolescents. In: Nowak AJ, Christensen JR, Mabry TR, Townsend JA, Wells MH, eds. *Pediatric Dentistry Infancy Through Adolescence*. 6th ed. Philadelphia, Pa.: Elsevier; 2019:371.
5. Alrayyes S, Hart TC. Periodontal disease in children. *Dis Mon* 2011;57(4):184-91.
6. Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: Framework and proposal of a new classification and case definition. *J Periodontol* 2018; 89(Suppl 1):S159-S172.
7. White JM, Elangovan S, Lee C-T. Periodontal risk assessment. In: Newman MG, Klokkevold PR, Elangova S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:537.
8. Douglass CW. Risk assessment and management of periodontal disease. *J Am Dent Assoc* 2006;137(Suppl): 27S-32S.
9. Chapple ILC. Risk assessment in periodontal care: The principles. In: Chapple ILC, Papapanou P, eds. *Risk Assessment in Oral Health: A Concise Guide for Clinical application*. Switzerland, AG: Springer Nature; 2020: 77-88.
10. Bouchard P, Carra MC, Boillot A, Mora F, Rangé H. Risk factors in periodontology: A conceptual framework. *J Clin Periodontol* 2017;44(2):125-31.
11. Lang NP, Suvan JE, Tonetti MS. Risk factor assessment tools for the prevention of periodontitis progression a systematic review. *J Clin Periodontol* 2015;42(Suppl 16): S59-70.
12. Mullins JM, Even JB, White JM. Periodontal management by risk assessment: A pragmatic approach. *J Evid Based Dent Pract* 2016;16(Suppl):91-8.
13. Petsos H, Arendt S, Eickholz P, Nickles K, Dannewitz B. Comparison of two different periodontal risk assessment methods with regard to their agreement: Periodontal risk assessment versus periodontal risk calculator. *J Clin Periodontol* 2020;47(8):921-32.
14. Sai Sujai GV, Triveni VS, Barath S, Harikishan G. Periodontal risk calculator versus periodontal risk assessment. *J Pharm Bioallied Sci* 2015;7(Suppl 2):S656-9.
15. Trombelli L, Minenna L, Toselli L, et al. Prognostic value of a simplified method for periodontal risk assessment during supportive periodontal therapy. *J Clin Periodontol* 2017;44(1):51-7.
16. Kwok V, Caton JG. Commentary: Prognosis revisited: A system for assigning periodontal prognosis. *J Periodontol* 2007;78(11):2063-71.
17. Elangovan S, Do JH, Takei HH, Carranza FA, Newman MG. Periodontal treatment plan. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:553-4.
18. Bimstein E, Eidelman E. Morphological changes in the attached and keratinized gingiva and gingival sulcus in the mixed dentition period. A 5-year longitudinal study. *J Clin Periodontol* 1988;15(3):175-9.
19. Bimstein E, Huja PE, Ebersole JL. The potential lifespan impact of gingivitis and periodontitis in children. *J Clin Pediatr Dent* 2013;38(2):95-9.
20. Cole E, Ray-Chaudhuri A, Vaidyanathan M, Johnson J, Sood S. Simplified basic periodontal examination (BPE) in children and adolescents: A guide for general dental practitioners. *Dent Update* 2014;41(4):328-37.
21. Murakami S, Mealey BL, Mariotti A, Chapple ILC. Dental plaque-induced gingival conditions. *J Periodontol* 2018;89(Suppl 1):S17-S27.
22. Sjödin B, Matsson L. Marginal bone loss in the primary dentition. A survey of 7-9-year-old children in Sweden. *J Clin Periodontol* 1994;21(5):313-9.
23. Drummond BK, Brosnan MG, Leichter JW. Management of periodontal health in children: Pediatric dentistry and periodontology interface. *Periodontol* 2000 2017;74(1): 158-67.
24. Bimstein E, Soskolne AW. A radiographic study of interproximal alveolar bone crest between the primary molars in children. *ASDC J Dent Child* 1988;55(5):348-50.
25. Al Jamal G, Al-Batayneh OB, Hamamy D. The alveolar bone height of the primary and first permanent molars in healthy 6- to 9-year-old Jordanian children. *Int J Paediatr Dent* 2011;21(2):151-9.
26. Bimstein E, Treasure ET, Williams SM, Dever JG. Alveolar bone loss in 5-year-old New Zealand children: Its prevalence and relationship to caries prevalence, socio-economic status and ethnic origin. *J Clin Periodontol* 1994;21(7): 447-50.



27. Bimstein E. Radiographic description of the distribution of aggressive periodontitis in primary teeth. *J Clin Pediatr Dent* 2018;42(2):91-4.
28. Tugnait A, Clerehugh V, Hirschmann PN. The usefulness of radiographs in diagnosis and management of periodontal diseases: A review. *J Dent* 2000;28(4):219-26.
29. Needleman HL, Ku TC, Nelson L, Allred E, Seow WK. Alveolar bone height of primary and first permanent molars in healthy seven- to nine-year-old children. *ASDC J Dent Child* 1997;64(3):188-96.
30. Bimstein E, Matsson L. Growth and development considerations in the diagnosis of gingivitis and periodontitis in children. *Pediatr Dent* 1999;21(3):186-91.
31. Chang CP, Barker JC, Hoeft KS, Guerra C, Chung LH, Burke NJ. Importance of content and format of oral health instruction to low-income Mexican immigrant parents: A qualitative study. *Pediatr Dent* 2018;40(1):30-6.
32. Renz ANPJ, Newton JT. Changing the behavior of patients with periodontitis. *Periodontol 2000* 2009;51:252-68.
33. Gorbunkova A, Pagni G, Brizhak A, Farronato G, Rasperini G. Impact of orthodontic treatment on periodontal tissues: A narrative review of multidisciplinary literature. *Int J Dent* 2016;2016:4723589.
34. Kanellis MJ, Owais AI, Warren JJ, et al. Managing caries in the primary dentition with silver nitrate: Lessons learned from a clinical trial. *J Calif Dent Assoc* 2018;46(1):37-44.
35. Silva DR. Periodontal health and disease in childhood and adolescents. In: Newman MG, Takei HH, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:286-94.
36. Yu CY, Abbott PV. Responses of the pulp, periradicular, and soft tissues following trauma to the permanent teeth. *Aust Dent J* 2016;61(Suppl 1):39-58.
37. Clerehugh V, Tugnait A. Diagnosis and management of periodontal diseases in children and adolescents. *Periodontol 2000* 2001;26:146-68.
38. Elangovan S, Takei HH. Nonsurgical phase of periodontal therapy. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:560-3.
39. Graziani F, Karapetsa D, Alonso B, Herrera D. Non-surgical and surgical treatment of periodontitis: How many options for one disease? *Periodontol 2000* 2017;75(1):152-88.
40. Jahn CA, Mancinelli-Lyle D. Plaque biofilm control for the periodontal patient. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:611.
41. Suvan J, Leira Y, Moreno Sancho FM, et al. Subgingival instrumentation for treatment of periodontitis. A systematic review. *J Clin Periodontol* 2020;47(Suppl 22):155-75. Available at: "<https://onlinelibrary.wiley.com/doi/10.1111/jcpe.13245>". Accessed March 5, 2024.
42. Feres M, Figueiredo LC, Soares GM, Faveri M. Systemic antibiotics in the treatment of periodontitis. *Periodontol 2000* 2015;67(1):131-86.
43. Herrera D, Matesanz P, Martín C, Oud V, Feres M, Teughels W. Adjunctive effect of locally delivered anti-microbials in periodontitis therapy: A systematic review and meta-analysis. *J Clin Periodontol* 2020;47(Suppl 22):239-56.
44. Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nat Rev Dis Primers* 2017;3:17038.
45. Pretzl B, Sälzer S, Ehmke B, et al. Administration of systemic antibiotics during non-surgical periodontal therapy—A consensus report. *Clin Oral Investig* 2019;23(7):3073-85.
46. Teughels W, Feres M, Oud V, Martín C, Matesanz P, Herrera D. Adjunctive effect of systemic antimicrobials in periodontitis therapy: A systematic review and meta-analysis. *J Clin Periodontol* 2020;47(Suppl 22):257-81.
47. Drisko CL. Periodontal debridement: Still the treatment of choice. *J Evid Based Dent Pract* 2014;14(Suppl):33-41.e1.
48. Dar-Odeh N, Fadel HT, Abu-Hammad S, Abdeljawad R, Abu-Hammad OA. Antibiotic prescribing for orofacial infections in the paediatric outpatient: A review. *Antibiotics (Basel)* 2018;7(2):38.
49. Albandar JM, Susin C, Hughes FJ. Manifestations of systemic diseases and conditions that affect the periodontal attachment apparatus: Case definitions and diagnostic considerations. *J Periodontol* 2018;89(Suppl 1):S183-S203.
50. Giannetti L, Roberto Apponi R, Dello Diago AM, Jafferany M, Goldust M, Sadoughifar R. Papillon-Lefèvre syndrome: Oral aspects and treatment. *Dermatol Ther* 2020;33(3):e13336.
51. Moghaddasi M, Ghassemi M, Shekari Yazdi M, Habibi SAH, Mohebi N, Goodarzi A. The first case report of Haim Munk disease with neurological manifestations and literature review. *Clin Case Rep* 2021;9(9):e04802.
52. Keestra JA, Grosjean I, Coucke W, Quirynen M, Teughels WJ. Non-surgical periodontal therapy with systemic antibiotics in patients with untreated chronic periodontitis: A systematic review and meta-analysis. *Periodontol Res* 2015;50(3):294-314.
53. Merchant SN, Vovk A, Kalash D, et al. Localized aggressive periodontitis treatment response in primary and permanent dentitions. *J Periodontol* 2014;85(12):1722-9.
54. Miller KA, Branco-de-Almeida LS, Wolf S, et al. Long-term clinical response to treatment and maintenance of localized aggressive periodontitis: A cohort study. *J Clin Periodontol* 2017;44(2):158-68.
55. Montenegro SCL, Retamal-Valdes B, Bueno-Silva B, et al. Do patients with aggressive and chronic periodontitis exhibit specific differences in the subgingival microbial composition? A systematic review. *J Periodontol* 2020;91(11):1503-20.

*References continued on the next page.*

56. Silva MP, Feres M, Sirotto TA, et al. Clinical and microbiological benefits of metronidazole alone or with amoxicillin as adjuncts in the treatment of chronic periodontitis: A randomized placebo-controlled clinical trial. *J Clin Periodontol* 2011;38(9):828-37.
57. Nibali L, Koidou VP, Hamborg T, Donos N. Empirical or microbiologically guided systemic antimicrobials as adjuncts to non-surgical periodontal therapy? A systematic review. *J Clin Periodontol* 2019;46(10):999-1012.
58. Rabelo CC, Feres M, Gonçalves C, et al. Systemic antibiotics in the treatment of aggressive periodontitis. A systematic review and a Bayesian Network meta-analysis. *J Clin Periodontol* 2015;42(7):647-57.
59. Sgolastra F, Petrucci A, Gatto R, Monaco A. Effectiveness of systemic amoxicillin/metronidazole as an adjunctive therapy to full-mouth scaling and root planing in the treatment of aggressive periodontitis: A systematic review and meta-analysis. *J Periodontol* 2012;83(6):731-43.
60. Sgolastra F, Severino M, Petrucci A, Gatto R, Monaco A. Effectiveness of metronidazole as an adjunct to scaling and root planing in the treatment of chronic periodontitis: A systematic review and meta-analysis. *J Periodontal Res* 2014;49(1):10-9.
61. Haas AN, de Castro GD, Moreno T, et al. Azithromycin as an adjunctive treatment of aggressive periodontitis: 12-months randomized clinical trial. *J Clin Periodontol* 2008;35(8):696-704.
62. Zhang Z, Zheng Y, Bian X. Clinical effect of azithromycin as an adjunct to non-surgical treatment of chronic periodontitis: A meta-analysis of randomized controlled clinical trials. *J Periodontal Res* 2016;51(3):275-83.
63. Araújo L, Demoly P. Macrolides allergy. *Curr Pharm Des* 2008;14(27):2840-62.
64. Bartold PM, du Bois AH, Gannon S, Haynes DR, Hirsch RS. Antibacterial and immunomodulatory properties of azithromycin treatment implications for periodontitis. *Inflammopharmacology* 2013;21(4):321-38.
65. Zeng L, Xu P, Choonara I, et al. Safety of azithromycin in pediatrics: A systematic review and meta-analysis. *Eur J Clin Pharmacol* 2020;76(12):1709-21.
66. American Academy of Pediatric Dentistry. Useful medications for oral conditions. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry 2024:642-9.
67. Jepsen S, Caton JG, Albandar JM, et al. Periodontal manifestations of systemic diseases and developmental and acquired conditions: Consensus report of workgroup 3 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Periodontol* 2018;89(Suppl 1):S237-S248.
68. Quirynen M, Dadamio J, Van den Velde S, et al. Characteristics of 2000 patients who visited a halitosis clinic. *J Clin Periodontol* 2009;36(11):970-5.
69. Teughels W, Dadamio J, Robben J, Dekeyser C, Quirynen. Halitosis (breath malodor). In: Newman MG, Klokkevold PR, Elangova S, Hernandez-Kapila Y, Carranza FA, eds. Newman and Carranza's Clinical Periodontology and Implantology. 14th ed. Philadelphia, Pa.: Elsevier; 2023: 441-52.
70. Motta LJ, Bachiega JC, Guedes CC, Laranja LT, Bussadori SK. Association between halitosis and mouth breathing in children. *Clinics* 2011;66(6):939-42. Available at: "https://www.sciencedirect.com/science/article/pii/S1807593222007517?ref=pdf\_download&fr=RR-2&rr=89df8f8a2902f7f4". Accessed July 4, 2024.
71. Nguyen BK, Quaraishi HA. Tonsillectomy and adenoidectomy—Pediatric Clinics of North America. *Pediatr Clin N Am* 2022;69(2):247-59.
72. Ballikaya E, Dogan BG, Onay O, Tekcicek MU. Oral health status of children with mouth breathing due to adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol* 2018;113:11-5.
73. Moraschini V, Calasans-Maia JA, Calasans-Maia MD. Association between asthma and periodontal disease: A systematic review and meta-analysis. *J Periodontol* 2018; 89(4):440-55.
74. Epstein JB, Thariat J, Bensadou R, et al. Oral complications of cancer and cancer therapy: From cancer treatment to survivorship. *CA Cancer J Clin* 2012;62(6):400-22.
75. American Academy of Pediatric Dentistry. Dental management of pediatric patients receiving immunosuppressive therapy and/or head and neck radiation. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2024:545-53.
76. Hong CH, da Fonseca M. Considerations in the pediatric population with cancer. *Dent Clin North Am* 2008;52(1):155-8.
77. da Fonseca M. Oral and dental care of local and systemic diseases. In: Nowak AJ, Christensen JR, Mabry TR, Townsend JA, Wells MH, eds. *Pediatric Dentistry Infancy Through Adolescence*. 6th ed. Philadelphia, Pa.: Elsevier; 2019:69-73.
78. Mawardi H, Alsubhi A, Salem N, et al. Management of medication-induced gingival hyperplasia: A systematic review. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2021;131(1):62-72.
79. Kantarci A, Marchesan JT, Divaris K. Gingival enlargement and management. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. Newman and Carranza's Clinical Periodontology and Implantology. 14th ed. Philadelphia, Pa.: Elsevier; 2023: 269-74.
80. Newman MG, Do JH, Takei HH, Whang M, Shin K. Periodontal surgical therapy. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. Newman and Carranza's Clinical Periodontology and Implantology. 14th ed. Philadelphia, Pa.: Elsevier; 2023: 730-9.
81. Bourguignon C, Cohenca N, Lauridsen E, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 1. Fractures and luxations. *Dent Traumatol* 2020;36(4):314-30.

82. Levin L, Day PF, Hicks L, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: General introduction. *Dent Traumatol* 2020;36(4):309-13.
83. Hermann NV, Lauridsen E, Ahrensburg SS, Gerds TA, Andreasen JO. Periodontal healing complications following concussion and subluxation injuries in the permanent dentition: A longitudinal cohort study. *Dent Traumatol* 2012;28(5):386-93.
84. Hermann NV, Lauridsen E, Ahrensburg SS, Gerds TA, Andreasen JO. Periodontal healing complications following extrusive and lateral luxation in the permanent dentition: A longitudinal cohort study. *Dent Traumatol* 2012;28(5):394-402.
85. Andersson L, Andreasen JO. Soft tissue injuries. In: Andreasen JO, Andreasen FM, Andersson L, eds. *Textbook and Color Atlas of Traumatic Injuries to the Teeth*. 5th ed. Copenhagen, Denmark: Wiley-Blackwell; 2018: 626-43.
86. Elias H, Baur DA. Management of trauma to supporting dental structures. *Dent Clin North Am* 2009;53(4): 675-89.
87. Rhee P, Nunley MK, Demetriades D, Velmahos G, Doucet JJ. Tetanus and trauma: A review and recommendations. *J Trauma* 2005;58(5):1082-8.
88. Day PF, Duggal M, Nazzari H. Interventions for treating traumatised permanent front teeth: Avulsed (knocked out) and replanted. *Cochrane Database Syst Rev* 2019;2 (2):CD006542.
89. Goswami M, Eranhikkal A. Management of traumatic dental injuries using different types of splints: A case series. *Int J Clin Pediatr Dent* 2020;13(2):199-202.
90. Sobczak-Zagalska H, Emerich K. Best splinting methods in case of dental injury: A literature review. *J Clin Pediatr Dent* 2020;44(2):71-8.
91. Fouad AF, Abbott PV, Tsilingaridis G, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 2. Avulsion of permanent teeth. *Dent Traumatol* 2020;36(4):331-42.
92. Khinda VIS, Kaur G, Brar GS, Kallar S, Khurana H. Clinical and practical implications of storage media used for tooth avulsion. *Int J Clin Pediatr Dent* 2017;10(2): 158-65.
93. Holmstrup P, Plemons J, Meyle J. Non-plaque-induced gingival diseases. *J Periodontol* 2018;89(Suppl 1): S28-S45.
94. Rawal SY, Claman LJ, Kalmar JR, Tatakis DN. Traumatic lesions of the gingiva: A case series. *J Periodontol* 2004; 75(5):762-9.
95. Romer M, Dougherty NJ. Oral self-injurious behaviors in patients with developmental disabilities. *Dent Clin North Am* 2009;53(2):339-50.
96. Dilsiz A, Aydin T. Self-inflicted gingival injury due to habitual fingernail scratching: A case report with a 1-year follow up. *Eur J Dent* 2009;3(2):150-4.
97. Krejci CB. Self-inflicted gingival injury due to habitual fingernail biting. *J Periodontol* 2000;71(6):1029-31.
98. Malaga EG, Aguilera EMM, Eaton C, Ameeraly P. Management of self-harm injuries in the maxillofacial region: A report of 2 cases and review of the literature. *Oral Maxillofac Surg* 2016;74(6):1198.e1-9.
99. Medina AC, Sogbe R, Gómez-Rey AM, Mata M. Factitious oral lesions in an autistic paediatric patient. *Int J Paediatr Dent* 2003;13(2):130-7.
100. Elangovan S, Takei HH. Surgical phase of periodontal therapy. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:683-7.
101. Takei HH, Scheyer T, Azzi RR, Allen EP, Han TJ. Periodontal plastic and esthetic surgery. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:775-7.
102. Klokkevold PR, Carranza FA. Acute gingival infections and management. In: Newman MG, Takei HH, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:249.
103. Schmidt J, Kunderova M, Pilbauerova N, Kapitan M. A review of evidence-based recommendations for pericoronitis management and a systematic review of antibiotic prescribing for pericoronitis among dentists: Inappropriate pericoronitis treatment is a critical factor of antibiotic overuse in dentistry. *Int J Environ Res Public Health* 2021;18(13):6796.
104. Moussa N, Ogle OE. Acute pain management. *Oral Maxillofac Surg Clin North Am* 2022;34(1):35-47.
105. Bridwell R, Gottlieb M, Koyfman A, Long B. Diagnosis and management of Ludwig's angina: An evidence-based review. *Am J Emerg Med* 2021;41:1-5.
106. Devi A, Narwal A, Bharti A, Kumar V. Premature loss of primary teeth with gingival erythema: An alert to dentist. *J Oral Maxillofac Pathol* 2015;19(2):271.
107. Delcourt-Debruyne EM, Boutigny HR, Hildebrand HF. Features of severe periodontal disease in a teenager with Chédiak-Higashi syndrome. *J Periodontol* 2000;71(5): 816-24.
108. Lozano ML, Rivera J, Sánchez-Guiu I, Vicente V. Towards the targeted management of Chédiak-Higashi syndrome. *Orphanet J Rare Dis* 2014;9:132.
109. Thumbigere Math V, Rebouças P, Giovani PA, et al. Periodontitis in Chédiak-Higashi Syndrome: An altered immunoinflammatory response. *JDR Clin Trans Res* 2018;3(1):35-46.
110. Kraut R. Implants for children. In: Babbush CA, Hahn JA, Krauser JT, Rosenlicht JL, eds. *Dental Implants-E-Book: The Art and Science*. 2nd ed. Maryland Heights, Mo.: Saunders Elsevier; 2010:389-402.
111. Kraut RA. Dental implants for children: Creating smiles for children without teeth. *Pract Periodontics Aesthet Dent* 1996;8(9):909-13.

*References continued on the next page.*

112. Deas DE, Moritz AJ, Sagun RS Jr, Gruwell SF, Powell CA. Scaling and root planing vs. conservative surgery in the treatment of chronic periodontitis. *Periodontol* 2000 2016;71(1):128-39.
113. Reynolds MA, Kao RT, Camargo PM, et al. Periodontal regeneration – intrabony defects: A consensus report from the AAP Regeneration Workshop. *J Periodontol* 2015; 86(2 Suppl):S105-7.
114. Aimetti M, Fratini A, Manavella V, et al. Pocket resolution in regenerative treatment of intrabony defects with papilla preservation techniques: A systematic review and meta-analysis of randomized clinical trials. *J Clin Periodontol* 2021;48(6):843-58.
115. Larsson L, Decker AM, Nibali L, Pilipchuk SP, Berglundh T, Giannobile WV. Regenerative medicine for periodontal and peri-implant diseases. *J Dent Res* 2016;95(3): 255-66.
116. Kao RT, Clark DR, Takei HH, Lin GH. Periodontal regeneration. In: Newman MG, Takei HH, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023: 756-65.
117. Myneni S, Bassir SH. Clinical practice guideline for treatment of periodontitis. In: Newman MG, Klokkevold PR, Elangovan S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023: 566.
118. Cortellini P, Stalpers G, Mollo A, Tonetti MS. Periodontal regeneration versus extraction and dental implant or prosthetic replacement of teeth severely compromised by attachment loss to the apex: A randomized controlled clinical trial reporting 10-year outcomes, survival analysis and mean cumulative cost of recurrence. *J Clin Periodontol* 2020;47(6):768-76.
119. Stavropoulos A, Bertl K, Spineli LM, Sculean A, Cortellini P, Tonetti M. Medium- and long-term clinical benefits of periodontal regenerative/reconstructive procedures in intrabony defects: Systematic review and network meta-analysis of randomized controlled clinical studies. *J Clin Periodontol* 2021;48(3):410-30.
120. Aoki A, Mizutani K, Schwarz F, et al. Periodontal and peri-implant wound healing following laser therapy. *Periodontol* 2000 2015;68(1):217-69.
121. Mossaad AM, Abdelrahman MA, Kotb AM, Alolayan AB, Elsayed SA. Gummy smile management using diode laser gingivectomy versus botulinum toxin injection: A prospective study. *Ann Maxillofac Surg* 2021;11(1):70-4.
122. Protásio ACR, Galvão EL, Falci SGM. Laser techniques or scalpel incision for labial frenectomy: A meta-analysis. *J Maxillofac Oral Surg* 2019;18(4):490-9.
123. Behdin S, Monje A, Lin GH, Edwards B, Othman A, Wang HL. Effectiveness of laser application for periodontal surgical therapy: Systematic review and meta-analysis. *J Periodontol* 2015;86(12):1352-63.
124. Jha A, Gupta V, Adinarayan R. LANAP, periodontics and beyond: A review. *J Lasers Med Sci* 2018;9(2):76-81.
125. Kao RT, Lin GH, John S, Klokkevold PR, Cobb CM. Lasers in periodontal and peri-implant therapy. In: Newman MG, Klokkevold PR, Elangova S, Hernandez-Kapila Y, Carranza FA, eds. *Newman and Carranza's Clinical Periodontology and Implantology*. 14th ed. Philadelphia, Pa.: Elsevier; 2023:770-1.
126. Garber DA, Salama MA. The aesthetic smile: Diagnosis and treatment. *Periodontology* 2000 1996;11(1):18-28.
127. Wennström JL. Mucogingival therapy. *Ann Periodontol* 1996;1(1):671-701.
128. Op Heij DG, Opdebeeck H, van Steenberghe D, Kokich VG, Belser U, Quirynen M. Facial development, continuous tooth eruption, and mesial drift as compromising factors for implant placement. *Int J Oral Maxillofac Implants* 2006;21(6):867078.
129. Bohner L, Hanisch M, Kleinheinz J, Jung S. Dental implants in growing patients: A systematic review. *Br J Oral Maxillofac Surg* 2019;57(5):397-406.
130. Gross EI, Nowak AJ. The dynamics of change. In: Nowak AJ, Christensen JR, Mabry TR, Townsend JA, Wells MH, eds. *Pediatric Dentistry Infancy Through Adolescence*. 6th ed. Philadelphia, Pa.: Elsevier; 2019:181.
131. Kamatham R, Avisa P, Vinnakota DN, Nuvvula S. Adverse effects of implants in children and adolescents: A systematic review. *J Clin Pediatr Dent* 2019;43(2):69-77.
132. Lambert F, Botilde G, Lecloux G, Rompen E. Effectiveness of temporary implants in teenage patients: A prospective clinical trial. *Clin Oral Implants Res* 2017; 28(9):1152-57.
133. Trombelli L, Simonelli A, Franceschetti G, Maietti E, Farina R. What periodontal recall interval is supported by evidence? *Periodontol* 2000 2020;84(1):124-33.
134. Axelsson P, Lindhe J. The significance of maintenance care in the treatment of periodontal disease. *J Clin Periodontol* 1981;8(4):281-94.
135. Axelsson P, Nyström B, Lindhe J. The long-term effect of a plaque control program on tooth mortality, caries and periodontal disease in adults. Results after 30 years of maintenance. *J Clin Periodontol* 2004;31(9):749-57.