

# Antibiotic Prophylaxis for Dental Patients at Risk for Infection

## Latest Revision

2019

### Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that numerous medical conditions predispose patients to bacteremia-induced infections. Because it is not possible to predict when a susceptible patient will develop an infection, prophylactic antibiotics are recommended when these patients undergo procedures that are at risk for producing bacteremia. These recommendations are intended to help practitioners make decisions regarding antibiotic prophylaxis for dental patients at risk.

### Methods

Recommendations on antibiotic prophylaxis for dental patients at risk for infection were developed by the Clinical Affairs Committee and adopted in 1990.<sup>1</sup> This document by the Council of Clinical Affairs is a revision of the previous version, last revised in 2014<sup>2</sup>, and based on a review of current dental and medical literature pertaining to post-procedural bacteremia-induced infections. This revision included database searches using key terms: infective endocarditis (IE), bacteremia, antibiotic prophylaxis, and dental infection. Articles were evaluated by title and/or abstract and relevance to dental care for children, adolescents, and those with special health care needs. Thirty-five citations were chosen from this method and from references within selected articles. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians. In addition, *Prevention of infective endocarditis: Guidelines from the American Heart Association*,<sup>3</sup> *Infective Endocarditis in Childhood: 2015 Update: A Scientific Statement From the American Heart Association*,<sup>4</sup> and the American Dental Association (ADA) report *The Use of Prophylactic Antibiotics Prior to Dental Procedures in Patients with Prosthetic Joints*<sup>5</sup> were reviewed.

### Background

Bacteremia (bacteria in the bloodstream) is anticipated following invasive dental procedures and can lead to complications in an immunodeficient patient.<sup>6,7</sup> High risk cardiac disease, immunosuppression, and immunodeficiencies may compromise one's ability to fight simple infection. The rationale for antibiotic prophylaxis is to reduce or eliminate transient bacteremia caused by invasive dental procedures.<sup>8</sup>

**How to Cite:** American Academy of Pediatric Dentistry. Antibiotic prophylaxis for dental patients at risk for infection. *The Reference Manual of Pediatric Dentistry*. Chicago, Ill.: American Academy of Pediatric Dentistry; 2020:447-52.

Antibiotic usage may result in the development of resistant organisms.<sup>3,6,7,9-11</sup> Utilization of antibiotic prophylaxis for patients at risk does not provide absolute prevention of infection. Post-procedural symptoms of acute infection (e.g., fever, malaise, weakness, lethargy) may indicate antibiotic failure and need for further medical evaluation.

The decision to use antibiotic prophylaxis should be made on an individual basis. Some medical conditions that may predispose patients to post-procedural infections are discussed below. This is not intended to be an exhaustive list; rather, the categorization should help practitioners identify children who may be at increased risk. If a patient reports a syndrome or medical condition with which the practitioner is not familiar, it is appropriate to contact the child's physician to determine susceptibility to bacteremia-induced infections.

To date, the evidence base supporting the efficacy and use of antibiotic prophylaxis is limited, especially in the pediatric population. Many of the indications are based on consensus.<sup>4,12-14</sup> The conservative use of antibiotics is indicated to minimize the risk of developing resistance to current antibiotic regimens.<sup>3,9,15,16</sup> Given the increasing number of organisms that have developed resistance to current antibiotic regimens, as well as the potential for an adverse anaphylactic reaction to the drug administered, it is best to be judicious in the use of antibiotics for the prevention of IE<sup>3</sup> or other distant-site infections.<sup>9,11,17</sup>

### Recommendations

Antibiotic prophylaxis is recommended with certain dental procedures,<sup>3,4,6,8,18</sup> but this should be directed against the most likely infecting organism. When procedures involve infected tissues or are performed on a patient with a compromised host response, additional doses or a prescribed pre- and post-operative regimen of antibiotics may be necessary. Emphasis should be placed on the prevention of disease, establishment of good oral health care habits, and routine oral health assessments

#### ABBREVIATIONS

**AAPD:** American Academy of Pediatric Dentistry. **ADA:** American Dental Association. **AHA:** American Heart Association. **CIED:** Cardiovascular implantable electronic device. **GI:** Gastrointestinal. **GU:** Genitourinary. **IE:** Infective endocarditis.

through a dental home. This may prevent the frequent need for the use of antibiotic therapy and, thus, decrease the risks of resistance and adverse events relation to use of antibiotics.<sup>8,19,20</sup>

### Patients with cardiac conditions

Infective endocarditis is an example of an uncommon but life-threatening complication resulting from bacteremia. The incidence of pediatric admissions due to infective endocarditis was between 0.05 and 0.12 cases per 1000 admissions in a multicenter study of U.S. children's hospitals from 2003-2010.<sup>4</sup> Only a limited number of bacterial species have been implicated in resultant postoperative infections; Viridans group streptococci, *Staphylococcus aureus* and *Enterococcus* species are the main microorganisms implicated in IE.<sup>3,4</sup> Enterococcal and other organisms such as *Haemophilus* species, *Aggregatibacter* species, *Cardiobacterium hominis*, *Eikenella corrodens*, and *Kingella* species are less common.<sup>4</sup> Routine daily activities such as toothbrushing, flossing, and chewing contribute more to the incidence of bacteremia when compared to dental procedures.<sup>4</sup> Thus, focus has shifted from antibiotic prophylaxis to an emphasis on oral hygiene and the prevention of oral diseases.<sup>4,8,13,14,18,20</sup>

In 2007, the American Heart Association (AHA) revised its guidelines for the prevention of IE and reducing the risk for producing resistant strains of bacteria.<sup>3</sup> The significant reasons for the revision include<sup>3</sup>:

- "IE is much more likely to result from frequent exposure to random bacteremias associated with daily activities than from bacteremia caused by a dental, [gastrointestinal] GI tract, or [genitourinary] GU tract procedure."<sup>3</sup> Daily activities would include toothbrushing, flossing, chewing, using toothpicks, using water irrigation devices, and other activities.
- "Prophylaxis may prevent an exceedingly small number of cases of IE if any, in individuals who undergo a dental, GI tract, or GU tract procedure.
- The risk of antibiotic-associated adverse events exceeds the benefit, if any, from prophylactic antibiotic therapy.
- Maintenance of optimal oral health and hygiene may reduce the incidence of bacteremia from daily activities and is more important than prophylactic antibiotics for a dental procedure to reduce the risk of IE."<sup>3</sup>

The AHA guidelines focus on antibiotic prophylaxis prior to certain dental procedures for patients in the highest risk group (See Table 1).<sup>3,4,6</sup> Globally, there is still a lack of consensus with regards to the benefit of antibiotic prophylaxis for prevention of infective endocarditis. Since the change in recommendations, the rate and incidence of IE have been low.<sup>4</sup>

Children with cyanosis with specific periodontal concerns may have an increased risk of IE, which makes optimum oral hygiene very important.<sup>3,4,22</sup> At-risk patients with poor oral hygiene and gingival bleeding after routine activities (e.g., toothbrushing) have shown an increased incidence of bacteremia as a measure for risk of IE.<sup>3,22,23</sup> The focus should

be on maintaining good oral hygiene, routine dental examinations, infection control to reduce bacteremia, and discouraging tattooing or piercing rather than relying on antibiotic prophylaxis for patients at risk.<sup>13,14,18-20,23</sup> These patients and their parents need to be educated and motivated to maintain personal oral hygiene through daily plaque removal, including flossing.<sup>3</sup> There is a shift in the emphasis on improved access to dental care and oral health in patients with underlying cardiac conditions at high risk for IE and less focus on a dental procedure and antibiotic coverage.<sup>4</sup> Professional prevention strategies should be based upon the individual's assessed risk for caries and periodontal disease.

In addition to those diagnoses listed in the AHA guidelines, patients with a reported history of injection drug use may be considered at risk for developing IE in the absence of cardiac anomalies.<sup>22</sup> Patients also should be discouraged from tattooing and piercing.<sup>13,14,24</sup> Consultation with the patient's physician may be necessary to determine susceptibility to bacteremia-induced infections.

Antibiotics are recommended for all dental procedures that involve manipulation of gingival tissue or the periapical region of teeth or perforation of the oral mucosa for cardiac patients with the highest risk<sup>3</sup> (see Tables 1 and 2). Specific antibiotic regimens can be found in Table 3. Practitioners and patients/parents can review the entire AHA guidelines in the AHA Circulation archives<sup>3</sup> (available at "<http://circ.ahajournals.org/cgi/content/full/116/15/1736>") for additional background information as well as discussion of special circumstances (e.g., patients already receiving antibiotic therapy, patients on anti-coagulant therapy).

### Patients with compromised immunity

Non-cardiac patients with a compromised immune system may be at risk for complications of bacteremia and distant site infection following invasive dental procedures. Existing evidence does not support the extensive use of antibiotic prophylaxis; prophylaxis should be limited to immunocompromised patients and those at high risk.<sup>19</sup> Consultation with the patient's physician is recommended for management of patients with a compromised immune system. Although there is not enough data to support its use, high risk patients who should be considered for use of prophylaxis includes, but is not limited to, those with<sup>13,14,25</sup>:

1. Immunosuppression\* secondary to:
  - a. human immunodeficiency virus (HIV);
  - b. severe combined immunodeficiency (SCIDS);
  - c. neutropenia;
  - d. cancer chemotherapy; or
  - e. hematopoietic stem cell or solid organ transplantation.
2. History of head and neck radiotherapy.
3. Autoimmune disease (e.g., juvenile arthritis, systemic lupus erythematosus).

\* Discussion of antibiotic prophylaxis for patients receiving immunosuppressive therapy and/or radiation therapy appears in a separate AAPD document.<sup>26</sup>

**Table 1. CARDIAC CONDITIONS ASSOCIATED WITH THE HIGHEST RISK OF ADVERSE OUTCOME FROM ENDOCARDITIS FOR WHICH PROPHYLAXIS WITH DENTAL PROCEDURES IS REASONABLE**

Prosthetic cardiac valve or prosthetic material used for cardiac valve repair
Previous infective endocarditis
Congenital heart disease (CHD)* <ul style="list-style-type: none"> <li>Unrepaired cyanotic CHD, including palliative shunts and conduits</li> <li>Completely repaired congenital heart defect with prosthetic material or device, whether placed by surgery or by catheter intervention, during the first six months after the procedure †</li> <li>Repaired CHD with residual defects at the site or adjacent to the site of a prosthetic patch or prosthetic device (which inhibit endothelialization)</li> </ul>
Cardiac transplantation recipients who develop cardiac valvulopathy

\* Except for the conditions listed above, antibiotic prophylaxis is no longer recommended for any other form of CHD.

† Prophylaxis is reasonable because endothelialization of prosthetic material occurs within six months after the procedure.

Reprinted with permission. *Circulation*. 2007;116:1736-1754. ©2007, American Heart Association, Inc.<sup>2</sup>

**Table 2. DENTAL PROCEDURES FOR WHICH ENDOCARDITIS PROPHYLAXIS IS REASONABLE FOR PATIENTS IN TABLE 1**

All dental procedures that involve manipulation of gingival tissue or the periapical region of teeth or perforation of the oral mucosa\*\*

\* The following procedures and events do not need prophylaxis: routine anesthetic injections through non-infected tissue, taking dental radiographs, placement of removable prosthodontic or orthodontic appliances, adjustment of orthodontic appliances, placement of orthodontic brackets, shedding of deciduous teeth, and bleeding from trauma to the lips or oral mucosa.

Reprinted with permission. *Circulation*. 2007;116:1736-1754. ©2007, American Heart Association, Inc.<sup>2</sup>

**Table 3. REGIMENS FOR A DENTAL PROCEDURE**

Regimen: Single Dose 30 to 60 min Before Procedure			
Situation	Agent	Adults	Children
Oral	Amoxicillin	2 g	50 mg/kg
Unable to take oral medication	Ampicillin OR	2 g IM or IV	50 mg/kg IM or IV
	Cefazolin or ceftriaxone	1 g IM or IV	50 mg/kg IM or IV
Allergic to penicillins or ampicillin—oral	Cephalexin*† OR	2 g	50 mg/kg
	Clindamycin OR	600 mg	20 mg/kg
	Azithromycin or clarithromycin	500 mg	15 mg/kg
Allergic to penicillin or ampicillin and unable to take oral medication	Cefazolin or ceftriaxone† OR	1 g IM or IV	50 mg/kg IM or IV
	Clindamycin	600 mg IM or IV	20 mg/kg IM or IV

IM indicates intramuscular; IV, intravenous.

\* Or other first- or second-generation oral cephalosporin in equivalent adult or pediatric dosage.

† Cephalosporins should not be used in an individual with a history of anaphylaxis, angioedema, or urticaria with penicillins or ampicillin.

Reprinted with permission. *Circulation*. 2007;116:1736-1754. ©2007, American Heart Association, Inc.<sup>2</sup>

**Table 4. MANAGEMENT OF PATIENTS WITH PROSTHETIC JOINTS UNDERGOING DENTAL PROCEDURES<sup>5</sup>****Clinical Recommendation:**

In general, for patients with prosthetic joint implants, prophylactic antibiotics are *not* recommended prior to dental procedures to prevent prosthetic joint infection.

For patients with a history of complications associated with their joint replacement surgery who are undergoing dental procedures that include gingival manipulation or mucosal incision, prophylactic antibiotics should only be considered after consultation with the patient and orthopedic surgeon.\* To assess a patient's medical status, a complete health history is always recommended when making final decisions regarding the need for antibiotic prophylaxis.

**Clinical Reasoning for the Recommendation:**

- There is evidence that dental procedures are not associated with prosthetic joint implant infections.
- There is evidence that antibiotics provided before oral care do not prevent prosthetic joint implant infections.
- There are potential harms of antibiotics including risk for anaphylaxis, antibiotic resistance, and opportunistic infections like *Clostridium difficile*.
- The benefits of antibiotic prophylaxis may not exceed the harms for most patients.
- The individual patient's circumstances and preferences should be considered when deciding whether to prescribe prophylactic antibiotics prior to dental procedures.

Copyright © 2015 American Dental Association. All rights reserved. This page may be used, copied, and distributed for non-commercial purposes without obtaining prior approval from the ADA. Any other use, copying, or distribution, whether in printed or electronic format, is strictly prohibited without the prior written consent of the ADA.

**ADA. Center for Evidence-Based Dentistry™**

\* In cases where antibiotics are deemed necessary, it is most appropriate that the orthopedic surgeon recommend the appropriate antibiotic regimen and when reasonable write the prescription.

Sollecito T, Abt E, Lockhart P, et al. The use of prophylactic antibiotics prior to dental procedures in patients with prosthetic joints. Evidence-based clinical practice guideline for dental practitioners — a report of the American Dental Association Council on Scientific Affairs. JADA. 2015;146(1):11–16.

Reprinted with permission. Copyright ©2015, American Dental Association. J Am Dent Assoc 2015;146(1):11-16.e8.

Publisher by Elsevier Inc. All right reserved. "www.ada.org".

4. Sickle cell anemia.<sup>27</sup>
5. Asplenicism or status post splenectomy.
6. Chronic high dose steroid usage.
7. Uncontrolled diabetes mellitus.
8. Bisphosphonate therapy.<sup>28,29</sup>
9. Hemodialysis.

**Patients with shunts, indwelling vascular catheters, or medical devices**

The AHA recommends that antibiotic prophylaxis for non-valvular devices, including indwelling vascular catheters (e.g., central lines) and cardiovascular implantable electronic devices (CIED), is indicated only at the time of placement of these devices in order to prevent surgical site infection.<sup>23,25</sup> The AHA found no convincing evidence that microorganisms associated with dental procedures cause infection of CIED and nonvalvular devices at any time after implantation.<sup>23,25</sup> The infections occurring after device implantation most often are caused by *Staphylococcus aureus* and coagulase negative staphylococci or other microorganisms that are non-oral in origin but are associated with surgical implantation or other active infections.<sup>23,29,30</sup> Consultation with the child's physician is recommended for management of patients with nonvalvular devices.

Ventriculoatrial (VA), ventriculocardiac (VC), or ventriculovenous (VV) shunts for hydrocephalus are at risk of bacteremia-induced infections due to their vascular access.<sup>25,31</sup> In contrast, ventriculoperitoneal (VP) shunts do not involve any vascular structures and, consequently, do not require

antibiotic prophylaxis.<sup>25,31</sup> Consultation with the child's physician is recommended for management of patients with vascular shunts.

**Patients with prosthetic joints**

For patients with a history of total joint arthroplasty, deep hematogenous infections can lead to life threatening complications such as a loss of the prosthetic joint or even increased morbidity and mortality.<sup>32,33</sup> Given the increasing risk of developing antibiotic resistance and adverse reactions, antibiotic prophylaxis prior to dental procedures is not recommended in the prevention of prosthetic joint infections.<sup>5</sup>

Consultation with the child's physician may be necessary for management of at-risk patients as well as patients with other implanted devices (e.g., Harrington rods, external fixation devices).<sup>25,32-35</sup>

**References**

1. American Academy of Pediatric Dentistry. Antibiotic chemoprophylaxis for pediatric dental patients. Boston, Mass.: American Academy of Pediatric Dentistry; 1990.
2. American Academy of Pediatric Dentistry. Antibiotic prophylaxis for dental patients at risk for infection. *Pediatr Dent* 2014;36(special issue):287-92.
3. Wilson W, Taubert KA, Gevitz M, et al. Prevention of infective endocarditis: Guidelines from the American Heart Association—A Guideline From the American Heart Association Rheumatic Fever, Endocarditis and Kawasaki

- Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation* 2007;116(15):1736-54. E-published April 19, 2007. Available at: "<http://circ.ahajournals.org/cgi/content/full/116/15/1736>". Accessed July 2, 2019. Erratum in *Circulation* 2007;116(15):e376-e7.
4. Baltimore RS, Gweitz M, Baddour LM, et al. Infective endocarditis in childhood: 2015 update: A scientific statement from the American Heart Association. *Circulation* 2015;132(15):1487-515.
  5. Sollecito TP, Abt E, Lockhart PB, et al. The use of prophylactic antibiotics prior to dental procedures in patients with prosthetic joints: Evidence-based clinical practice guideline for dental practitioners—A report of the American Dental Association Council on Scientific Affairs. *J Am Dent Assoc* 2015;146(1):11-6.
  6. Lockhart PB, Brennan MT, Kent ML, Norton JH, Weinrib DA. Impact of amoxicillin prophylaxis on the incidence, nature, and duration of bacteremia in children after intubation and dental procedures. *Circulation* 2004;109(23):2878-84.
  7. Roberts GJ, Jaffrey EC, Spract DA, Petrie A, Greville C, Wilson M. Duration, prevalence and intensity of bacteremia after dental extractions in children. *Heart* 2006;92(9):1274-7.
  8. Daly CG. Antibiotic prophylaxis for dental procedures. *Aust Prescr* 2017;40(5):184-8.
  9. Fluent MT, Jacobsen PL, Hicks LA. Considerations for responsible antibiotic use in dentistry. *J Am Dent Assoc* 2016;147(8):683-6.
  10. Dajani AS, Taubert KA, Wilson W, et al. Prevention of bacterial endocarditis: Recommendations by the American Heart Association. *JAMA* 1997;227(22):1794-801.
  11. Centers for Disease Control and Prevention. Antibiotic/Antimicrobial resistance. About antimicrobial resistance: A brief overview. Available at: "<https://www.cdc.gov/drugresistance/about.html>". Accessed March 26, 2019. Archived by WebCite® at: "<http://www.webcitation.org/779R5H5EJ>"
  12. Glenny AM, Oliver R, Roberts GJ, Hooper L, Worthington HV. Antibiotics for the prophylaxis of bacterial endocarditis in dentistry. *Cochrane Database System Rev* 2013;4(10):CD003813. Available at: "<https://doi-org.proxy.library.adelaide.edu.au/10.1002/14651858.CD003813pub4>".
  13. Cahill TJ, Dayer M, Prendergast B, Thornhill M. Do patients at risk of infective endocarditis need antibiotics before dental procedures? *BMJ* 2017;358:j3942.
  14. Cahill TJ, Harrison JL, Jewell P, et al. Antibiotic prophylaxis for infective endocarditis: A systematic review and meta-analysis. *Heart* 2017;103(12):937-44.
  15. Watters W, Rethman MP, Hanson NB, et al. Prevention of orthopaedic implant infection in patients undergoing dental procedures. *J Am Acad Orthop Surg* 2013;21(3):180-9.
  16. de Sa DD, Tleyieh IM, Anavekar NS, et al. Epidemiological trends of infective endocarditis: A population-based study in Olmsted County, Minnesota. *Mayo Clin Proc* 2010;85(5):422-6. Erratum in *Mayo Clin Proc* 2010;85(8):722.
  17. Thornhill MH, Dayer MJ, Prendergast B, Baddour LM, Jones S, Lockhart PB. Incidence and nature of adverse reactions to antibiotics used as endocarditis prophylaxis. *J Antimicrob Chemother* 2015;70(8):2382-8.
  18. National Heart Foundation of New Zealand Advisory Group. Guideline for the prevention of infective endocarditis associated with dental and other medical interventions. Auckland: National Heart Foundation of New Zealand. Available at: "<http://www.ttophs.govt.nz/vdb/document/312>". Accessed December 15, 2018. (Archived by WebCite® at: "<http://www.webcitation.org/74gYDlgr>")
  19. Habib G, Hoen B, Tornos P, et al. Guidelines on the prevention, diagnosis, and treatment of infective endocarditis (new version 2009): The Task Force on the Prevention, Diagnosis, and Treatment of Infective Endocarditis of the European Society of Cardiology (ESC). *Eur Heart J* 2009;30(9):2369-413.
  20. National Institute for Health and Care Excellence. Context In: Prophylaxis against infective endocarditis: Antimicrobial prophylaxis against infective endocarditis in adults and children undergoing interventional procedures. London: NICE. Available at: "<https://www.nice.org.uk/guidance/cg64/chapter/Recommendations>". Accessed December 15, 2018. (Archived by WebCite® at: "<http://www.webcitation.org/74gYehSug>")
  21. Lockhart PB, Brennan MT, Sasser HC, Fox PC, Paster BJ, Bahrani-Mougeot FK. Bacteremia associated with tooth-brushing and dental extraction. *Circulation* 2008;117(24):3118-25.
  22. Gewitz MH, Taubert KA. Infective endocarditis and prevention. In: Moss AJ, Adams FH, Allen HD, eds. *Moss and Adam's Heart Disease in Infants, Children, and Adolescents: Including the Fetus and Young Adult*, 8th ed. Philadelphia, Pa.: Lippincott Williams & Wilkins; 2016:1441-53.
  23. Baddour LM, Epstein AE, Erickson CC, et al. Update on cardiovascular implantable electronic device infections and their management. *Circulation* 2010;121(3):458-77.
  24. Lick SD, Edozie SN, Woodside KJ, Conti VR. Streptococcus viridans endocarditis from tongue piercing. *J Emerg Med* 2005;29(1):57-9.
  25. Lockhart PB, Loven B, Brennan MT, Fox PC. The evidence base for the efficiency of antibiotic prophylaxis in dental practice. *J Am Dent Assoc* 2007;138(4):458-74.

*References continued on the next page.*

26. American Academy of Pediatric Dentistry. Dental management of pediatric patients receiving immunosuppressive therapy and/or radiation therapy. *Pediatr Dent* 2018;40(6):392-400.
27. Tate AR, Norris CK, Minniti CP. Antibiotic prophylaxis for children with sickle cell disease: A survey of pediatric dentistry residency program directors and pediatric hematologists. *Pediatr Dent* 2006;28(3):332-5.
28. Montefusco V, Gay F, Spina F, et al. Antibiotic prophylaxis before dental procedures may reduce the incidence of osteonecrosis of the jaw in patients with multiple myeloma treated with bisphosphonates. *Leuk Lymphoma* 2008;49(11):2156-62.
29. Rogers SN, Hung J, Barber AJ, Lowe D. A survey of consultant members of the British Association of Oral and Maxillofacial Surgeons regarding bisphosphonate-induced osteonecrosis of the jaws. *Br J Oral Maxillofac Surg* 2009;47(8):598-601.
30. Hong CHL, Allred R, Napenas JJ, Brennan MT, Baddour LM, Lockhart PB. Antibiotic prophylaxis for dental procedures to prevent indwelling venous catheter-related infections. *Am J Med* 2010;123(12):1128-33.
31. Baddour LM, Bettman MA, Bolger AF, Bolger A, Ferrieri P. Nonvalvular cardiovascular device-related infections. *Circulation* 2003;108(16):2015-31.
32. Aminoshariae A, Kulild J. Premedication of patients undergoing dental procedures causing bacteremia after total joint arthroplasty. *J Endod* 2010;36(6):974-7.
33. Rethman MP, Watters W 3rd, Abt E, et al. The American Academy of Orthopedic Surgeons and the American Dental Association clinical practice guideline on the prevention of orthopaedic implant infection in patients undergoing dental procedures. *J Bone Joint Surg* 2013;95(8):745-7.
34. Berbari EF, Osmon DR, Carr A, et al. Dental procedures as risk factors for prosthetic hip or knee infection: A hospital-based prospective case-control study. *Clin Infect Dis* 2010;50(1):8-16. Erratum in *Clin Infect Dis* 2010;50(6):944.
35. Little JW, Jacobson JJ, Lockhart PB, American Academy of Oral Medicine. The dental treatment of patients with joint replacements: A position paper from the American Academy of Oral Medicine. *J Am Dent Assoc* 2010;141(6):667-71.