

Oral and Dental Aspects of Child Abuse and Neglect: Clinical Report

CLINICAL REPORT Guidance for the Clinician in Rendering Pediatric Care

September 2024



How to Cite: Tate AR, Fisher-Owens SA, Spiller L, et al; American Academy of Pediatrics, Section on Oral Health, Council on Child Abuse and Neglect Oral and Dental Aspects of Child Abuse and Neglect: Clinical Report. *Pediatrics* 2024;154(3):e2024068024.

Authors

Anupama Rao Tate, DMD, MPH,^a Susan A. Fisher-Owens, MD, MPH, FAAP,^b Lora Spiller, MD, FAAP,^c Jillian Muhlbauer, DMD,^d James L. Lukefahr, MD, FAAP,^e Section on Oral Health, Council on Child Abuse and Neglect

Abstract

In all 50 states, the District of Columbia, and the Commonwealth of Puerto Rico, pediatricians, dental professionals, and other physicians are mandated to report suspected cases of abuse and neglect to social service or law enforcement agencies. The purpose of this clinical report is to review the oral aspects of abuse and dental neglect in children and the role of pediatricians, dental professionals, and other physicians in evaluating such conditions. This clinical report addresses recommendations on the evaluation of bite marks, as well as perioral and intra-oral injuries, infections, and diseases that may raise suspicion for child abuse or neglect. Some physicians may have received less education pertaining to oral health, dental injury, and oral disease. These physicians may not detect the mouth and gum findings possibly related to abuse or neglect as readily as abuse injuries involving other areas of the body. Therefore, pediatricians, dental professionals, and other physicians are encouraged to collaborate to increase the prevention, detection, and treatment of these conditions in children.

^aDivision of Oral Health, Children's National Hospital, and Department of Pediatrics, George Washington Medical School, Washington, District of Columbia; ^bDepartments of Pediatrics and Preventive and Restorative Dental Sciences, University of California, San Francisco, California; ^cDivision of Child Abuse Pediatrics, Department of Pediatrics, University of Texas Health San Antonio, San Antonio, Texas; and ^dPrivate practice, Arnold, Maryland.

All authors were responsible for reviewing and updating of one or more sections. Drs Tate and Lukefahr were responsible for conceptualizing, writing, and revising the manuscript and considering input from all reviewers and the board of directors; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Clinical reports from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, clinical reports from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this report does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

DOI: <https://doi.org/10.1542/peds.2024-068024>

Address correspondence to Anupama Rao Tate, DMD, MPH. E-mail: anutate@gmail.com

BACKGROUND

Child abuse and neglect affect at least 1 out of 7 children in the United States each year.¹ Maltreatment incidents may have been higher than observed during and after the global severe acute respiratory syndrome coronavirus 2 pandemic because of factors such as increased family stressors, isolation away from community support systems, and underreporting.²

Children may be exposed to multiple kinds of maltreatment that manifest in the mouth. Abuse can include physical and sexual abuse and may be evidenced by bite marks, dental neglect, and medical child abuse (MCA). Bullying and the human trafficking of children also occur and can have serious long-term effects. Abuse and neglect may be the presenting problem, noticed during a physical examination, or children or adolescents may disclose information about these experiences. It is important for pediatricians, dental professionals, and other physicians to be alert to and knowledgeable about signs and symptoms of child abuse and neglect and to know how to respond. Because of varying resources within different communities, not all physician or dental professionals of a certain job specification may be available everywhere, and thus, job roles may sometimes overlap.

All pediatricians, dental professionals, and other physicians in the United States have a legal requirement to report suspected child abuse and neglect to child protection authorities, although the specific requirements and procedures vary among jurisdictions. Racial, ethnic, and economic disparities in the reporting of child maltreatment to authorities have long been recognized³ and have been under closer scrutiny in recent years. Children with disabilities are a vulnerable population at increased risk of child abuse and neglect, and therefore, special attention should be paid to reduce this risk in these children and youth.⁴ The Joint Commission now requires health care organizations to use written criteria to identify patients who may have experienced physical assault, sexual assault, sexual molestation, domestic abuse, or elder or child abuse and neglect.⁵ A recognized bias among health care professionals is the failure to report suspected abuse when the family

is of similar background to the professional.⁶ Recognition of implicit bias and its influence on diagnosing and reporting child maltreatment is also an important consideration.⁷⁻⁹

PHYSICAL ABUSE

According to the Fourth National Incidence Study of Child Abuse and Neglect, in 2006 (the most recent year evaluated) at least 323 000 US children experienced physical abuse.¹⁰ Identifying and reporting suspected abuse can be challenging for pediatricians, dental professionals, and other physicians but has the potential to interrupt the cycle of abuse, secure the child's safety, and in some cases, save the child's life.¹¹ Missed opportunities to recognize and report physical abuse remain a significant problem; in one study,¹² 27.5% of abused infants had a previous minor but inflicted injury that had been recognized by a parent or physician but usually not reported to authorities. The majority of these "sentinel injuries" involved bruising to the face, forehead, or ear. The mnemonic **TEN-4** (Torso, Ear, or Neck in children less than or equal to 4 years of age, and anywhere in an infant under 4 months of age) has been widely adopted to assist in identifying bruises concerning for abuse.¹³ An expanded mnemonic with improved sensitivity for identifying abusive bruise injuries has since been introduced: **TEN-4-FACESp** (Torso, Ear, Neck, Frenulum, Angle of jaw, Cheeks, Eyelids, Subconjunctivae in children younger than 4 years, bruising anywhere in infants younger than 5 months, or patterned bruising).¹⁴

Craniofacial, head, face, and neck injuries occur in more than half of child abuse cases.^{12,15-18} All suspected individuals who have experienced abuse or neglect, including children in state custody or foster care, need to be examined carefully at some point during the evaluation for signs of oral trauma, caries, gingivitis, and other oral health problems. These conditions are more prevalent in children who are maltreated than in the general pediatric population.¹⁹

Because of their exposed location on the body and anatomic complexity, the head and face are more vulnerable to serious injury than are other body regions.²⁰ A review of 40 cases of child homicides resulting from family violence noted that 36 (90%) had orofacial injuries.²¹ Recent research has found that children younger than 4 years with orofacial (especially cheek, ear, jaw, and frenum) and torso bruising are at risk for having been physically abused and are at risk for future, more serious abuse.¹⁴ Of children up to age 5 years with oral injury who underwent skeletal imaging, 25% had occult fractures; 30% of 37 infants with oral injury who underwent neuroimaging had central nervous system injuries.²² Oral injuries may be inflicted with instruments such as eating utensils or a bottle during forced feedings, hands, fingers, scalding liquids, or caustic substances. Physical abuse may result in contusions; burns or lacerations of the tongue, lips, buccal mucosa, palate (soft and hard), gingiva, alveolar mucosa, or frenum; fractured, displaced, or avulsed teeth; or facial bone and jaw fractures. Naidoo²³ cited the lips as the most common site for inflicted oral injuries (54%) followed by the oral mucosa, teeth, gingiva, and tongue. Lacerations to the oral frena in premobile infants are often the result of physical abuse and are frequently associated with other findings of serious physical abuse.^{22,24} Trauma to the teeth may result in pulpal necrosis, leaving the teeth gray and discolored.²⁵ Gags applied to the mouth may result in bruises, lichenification, or scarring at the corners of the mouth.²⁶ In some cases, serious injuries of the oral cavity, including posterior pharyngeal injuries and retropharyngeal abscesses, have been inflicted by caregivers who fabricate illness in a child²⁷ to simulate hemoptysis or other symptoms requiring medical care. All findings in cases in which there is reasonable suspicion of abuse or neglect, regardless of mechanism, need to be reported for further investigation. Unintentional or accidental injuries to the mouth are common and may be distinguished from abuse by judging whether the history (including the timing and mechanism of the injury) is consistent with the characteristics of the injury and the child's developmental capabilities. Multiple injuries, injuries in different stages of healing, or a discrepant history ought to arouse suspicion for abuse. In conjunction with reporting, consultation with or referral to a knowledgeable

dentist or child abuse pediatrician may be helpful. The clinical report from the American Academy of Pediatrics (AAP) titled "The Evaluation of Suspected Child Physical Abuse" provides additional guidance.¹¹

SEXUAL ABUSE

The oral cavity is a frequent site of sexual abuse in children, and acquisition of sexually transmitted infections (STIs) via this route can occur.²⁸ Testing for STIs within the oral cavity may be considered when:

- The child gives a history of oral-genital contact;
- The child has experienced penetration or has evidence of recent or healed penetrative injury to the genitals, anus, or oropharynx;
- The child has been abused by a stranger;
- The child has been abused by an assailant known to be infected with or at high risk for STIs;
- The child has a sibling, other relative, or household contact with an STI;
- The child lives in an area with a high rate of STIs in the community;
- The child has signs or symptoms of an STI (eg, vaginal discharge, genital itching or odor, urinary symptoms, or genital lesions);
- The child or the parent of a prepubertal child requests STI testing; and
- The child is unable to verbalize details of the assault.²⁹

Culture or nucleic acid amplification tests (NAATs) may be used to test for *Neisseria gonorrhoeae* and *Chlamydia trachomatis* infection. NAATs are preferred to culture by most clinicians because of their superior sensitivity. Only Clinical Laboratory Improvement Amendments-approved, US Food and Drug Administration-cleared NAAT assays should be used.²⁹ Consultation with an expert is recommended for positive STI test results in infants and prepubertal children, because vertical and nonsexual transmission of certain pathogens can occur (eg, persistence of perinatally acquired *Chlamydia*, postnatal HIV exposure through infected human milk, and others).^{30,31}

In one series, STIs were detected in 22.9% of adolescents evaluated for sexual abuse and in 11.4% of preadolescents.³² Oropharyngeal STIs are detected less frequently than anogenital infections; they commonly do not present with pain, discharge, erythema, or other visible signs of injury or infection.^{33,34} When an STI is confirmed in a child or adolescent being evaluated for sexual abuse, testing of other sites, including the oropharynx, is indicated regardless of what contact the child disclosed may be indicated. In one study, NAATs for oropharyngeal *Chlamydia* and gonorrhea were positive in 6% of adolescent patients, despite more than half of these patients providing no history of genital-oral contact.³⁴ Oral infection with *Trichomonas vaginalis* has never been described. Anogenital and oropharyngeal human papillomavirus infections may be transmitted sexually, vertically from birthing parent to infant during delivery, or horizontally through nonsexual contact from a child or caregiver's hand to the genitals or mouth.^{35,36} In the setting of sexual abuse, national practice guidelines from the AAP and Centers for Disease Control and Prevention support postexposure vaccination against human papillomavirus for children 9 years and older if they have not already started the vaccine series.^{33,37}

Children who present acutely with a recent history of sexual abuse need to receive specialized forensic evidence collection for semen and other foreign materials resulting from the assault. Specialized hospitals and child protection clinics equipped with protocols and experienced personnel are best suited for collecting such specimens and maintaining a chain of evidence necessary for investigations. In many cases, oral swab specimens are collected irrespective of whether the child specifically discloses oral-genital contact. Typically, the buccal mucosa and tongue are swabbed with a sterile, cotton-tipped applicator; the swab can be air dried and packaged appropriately for laboratory analysis.

MEDICAL CHILD ABUSE

MCA, also known as "Munchausen syndrome by proxy" and "caregiver fabricated-illness," is a form of child maltreatment in which a caregiver

subjects a child to unnecessary and potentially harmful medical care because of the caregiver's exaggeration, falsification, or induction of an illness or medical condition.³⁸ These children are subjected to unnecessary medical interventions, diagnostic tests, repeated examinations, surgeries, and medications. There is no typical presentation of MCA, although commonly, the fabricated illness involves multiple organ systems. Apnea, seizures, feeding problems, bleeding, and rashes are frequently reported symptoms. One recent study found that gastrointestinal complaints in children who had experienced MCA led to 6 times more types of feeding complications compared with controls who were also being evaluated for gastrostomy tube placement. In this study, the majority of patients experienced resolution of their nonoral feeding issues after MCA intervention (ie, removal from potential offender).³⁹

MCA can also present with oral manifestations. One review described 4 cases of MCA involving the oral cavity in children ranging from 9 months to 6 years of age.⁴⁰ In one case, recurrent mucosal bleeding and ulcerations were initially treated as Stevens Johnson syndrome and then cicatricial pemphigoid. This child died of poisoning and her oral manifestations were found to be caused by forced ingestion of toxic cleaning products. Another case presented with mucosal desquamation, drooling, and dysphagia and was eventually discovered to be caused by the caregiver placing an acid-based powder in the infant's food and toys. In all 4 cases, and in most reports in the literature, the perpetrator was the mother.⁴¹

Making a diagnosis of MCA can be difficult because of false information provided and/or induction of illness by the caregiver. Numerous medical professionals from various specialties and institutions are often involved in treating these patients. Diagnosing MCA may require a thorough review of all available medical records, which is often voluminous and can be time consuming, but is needed to discover concerning patterns and make the correct diagnosis.⁴²

BITE MARKS ON THE SKIN

Acute or healed bite marks on the skin may indicate abuse. Dentists trained as forensic odontologists can assist pediatricians, other dental professionals, and physicians in the detection and evaluation of bite marks related to physical and sexual abuse.⁴³ Bite marks on the skin need to be suspected when ecchymoses, abrasions, or lacerations are found in an elliptical, horseshoe-shaped, or ovoid pattern.⁴³ Frequently, bite marks are found on the cheeks, back, sides, arms, buttocks, and genitalia.⁴⁴ Bites produced by dogs and other carnivorous animals tend to tear flesh, whereas human bites compress flesh and can cause abrasions, contusions, and lacerations, but rarely avulsions of tissue.⁴⁵ An intercanine distance (ie, the linear distance between the central point of the cuspid tips) measuring more than 3.0 cm is suspicious for an adult human bite.⁴⁵

The pattern, size, contour, color, and evolution of a bite mark ideally can be evaluated by a forensic odontologist. If a specialist is unavailable, a child abuse pediatrician or pediatric dental professional experienced in identifying the patterns of child abuse injuries may examine and document the bite mark characteristics photographically with an identification tag and scale marker (eg, ruler) in the photograph. Further information is available on the American Board of Forensic Odontology (ABFO) Website to locate an expert and review the standards and guidelines for evaluating bite marks (www.abfo.org).

Bite marks found on human skin are challenging to interpret and differentiate because of similarities in dentition, distortion on skin, and time elapsed between the injury and the analysis.⁴³ Recent investigations have led to questions about the scientific validity and overapplication of forensic patterned evidence (bite mark analysis in particular) and its role in legal proceedings.⁴⁶ In response, the ABFO has adopted more conservative guidelines and no longer endorses positive identification of a suspect dentition to a bitemark.⁴⁶ More high-quality research is needed regarding bite mark analysis and its use in identifying potential perpetrators.

BULLYING

Because children with dental abnormalities are at increased risk for bullying, advocating for antibullying prevention programs in schools and other community settings by pediatricians, dental professionals, and other physicians may be beneficial. Thirty percent of children in the sixth through 10th grades report having been bullied and/or having bullied others.⁴⁷ Investigators from numerous countries and cultures have found that children with orofacial or dental abnormalities (including malocclusion) are frequently subjected to bullying,⁴⁸⁻⁵⁰ and as a result, may suffer serious psychological consequences, including depression and suicidal ideation.⁵¹⁻⁵³ Children who reported physical abuse, intimate partner violence, forced sex, and bullying were found to also report poor oral health.⁵⁴ Some children and adolescents are at high risk for being bullied for reasons besides orofacial abnormalities but are likely to have interactions with pediatricians or dentists. These include LGBTQ+ youth,⁵⁵ overweight children and adolescents,⁵⁶ children from racially minoritized groups,⁵⁷ and children with special health care needs.⁵⁸ There are antibullying prevention programs in schools and other community settings, as well as AAP resources including HealthyChildren.org, which offers caregivers educational resources on bullying among many other topics (<https://healthychildren.org/English/safety-prevention/at-play/Pages/Bullying-Its-Not-Ok.aspx>).

HUMAN SEX TRAFFICKING

Human trafficking is a major global child health issue with substantial medical, psychological, and dental ramifications. The term "human trafficking" is often confused with "human smuggling"; however, human trafficking requires exploitation and does not require any physical transportation from one location to another to meet the definition.⁵⁹ This clinical report focuses on human sex trafficking, defined by the US Department of State as "a commercial sex act that is induced by force, fraud, or coercion, or in which the person induced to perform such an act has not attained 18 years of age."⁵⁹ Precise numbers of children experiencing human or sex trafficking are difficult to obtain because of the complicated nature of these definitions, as well as underreporting and failure to recognize trafficking. The average age of children who are exploited for sex is 12 years, and children as young as 6 years are targeted.⁶⁰ Children who are or have been in foster care,⁶¹ are experiencing homelessness, identify as LGBTQ+,⁶² run away from home,⁶³ or are incarcerated in juvenile detention facilities⁶⁴ are more likely to be trafficked. Advances in technology, with the Internet and mobile devices, have facilitated perpetrators' ability to traffic children, often without requirement of the child to leave their own home.

Trafficking survivors often experience significant physical and sexual violence and may have observable injuries that are poorly explained. Survivors also are at higher risk for STIs, unsafe abortions, malnutrition, and untreated medical conditions. Posttraumatic stress disorder, anxiety, depression, suicidal ideation, self-harm, and substance use concerns are common to people who have experienced human trafficking.⁶⁴ Dental problems also rank high in this list: For women and adolescents who were trafficked in Europe, 58% reported tooth pain.⁶⁵ In the United States, more than half (54.3%) of women and adolescents who were trafficked reported dental problems, most commonly tooth loss (42.9%).⁶⁶ Children who have experienced trafficking may present to dental offices for care. It has been reported that child trafficking survivors have twice the risk for dental problems, because they "often suffer from inadequate nutrition leading to retarded growth and poorly formed teeth, as well as dental caries, infections, and tooth loss."⁶⁷ For older children, dental problems may trace back to their situation of origin, with limited access to or poor quality of care.

Trafficked children experience unique challenges in accessing medical and dental care. However, studies have shown that a large percentage of these youth do present for health care during the time they are being trafficked.⁶⁸ One study found that 26.5% of trafficked individuals were seen by a dentist during the time they were being trafficked, despite the fact that few were identified by clinicians.⁶⁹ Validated screening tools exist for health care settings but are still underutilized.⁷⁰

DENTAL NEGLECT

The American Academy of Pediatric Dentistry recognizes the influence of social factors on children's oral health including access to care, dental disease, behaviors, and oral health inequalities.⁷¹ Dental neglect, as defined by the American Academy of Pediatric Dentistry, is the "willful failure of parent or guardian, despite adequate access to care, to seek and follow through with treatment necessary to ensure a level of oral health essential for adequate function and freedom from pain and infection."⁷² Dental caries, periodontal diseases, and other oral conditions can lead to pain, infection, loss of function, and systemic infection if left untreated.⁷² These undesirable outcomes can adversely affect learning, communication, nutrition, and other activities necessary for normal growth and development.^{18,73} Dental neglect is the most frequent type of abuse seen by dentists and can be challenging to identify because it is chronic and insidious in nature.^{74,75} Some children who present for dental care have severe early childhood caries (formerly termed "infant bottle" or "nursing" caries). It is important to differentiate caregivers with adequate knowledge but willful failure to seek care from caregivers with barriers to care or those without knowledge or awareness of their child's need for dental care when determining the need to report such cases to child protective services. Pediatricians can play an important role in encouraging families to seek dental care and explaining benefits of completing needed dental treatment.⁷⁶ Several factors are considered necessary for the diagnosis of neglect⁷⁷ :

- a child is harmed or at risk for harm because of lack of dental health care;
- the recommended dental care offers significant net benefit to the child;
- the anticipated benefit of the dental treatment is significantly greater than its morbidity, so parents would likely choose treatment over nontreatment;
- access to health/dental care is available but not used; and
- the parent understands the dental advice given.

Failure to seek or obtain proper dental care may result from factors such as family isolation (physical or cultural), lack of finances, transportation difficulty, ignorance, or lack of perceived value of oral health.⁷⁸⁻⁸⁰ The point at which to consider a parent negligent and begin intervention occurs after the parent has been properly alerted by a pediatrician, dental professional, or other physician about the nature and extent of the child's condition, the specific treatment needed, and the mechanism of accessing that treatment.⁷⁸ Many families face challenges in accessing dental care or insurance for their children, and dental services may not be readily available. A child's social, emotional, and medical ability to undergo treatment, as well as cultural and religious differences, are important to consider.⁸⁰

Pediatricians, dental professionals, and other physicians can make a reasonable effort to help caregivers understand the explanation of the dental disease and its implications, the risks and benefits of dental treatment, and the use of the appropriate analgesic and anesthetic during the procedure to ensure the child's comfort. Assistance can be provided to the family in finding financial aid, transportation, or public facilities for needed services. If, despite these efforts, the parent fails to obtain therapy, the case warrants reporting to the appropriate child protective services agency.⁷⁸

CONCLUSIONS

Physical or sexual abuse may also result in oral or dental injuries or conditions. Pediatricians, dental professionals, and other physicians can document suspicious injuries, obtain laboratory evidence, photo documentation, and/or consult with experts when appropriate. The role of forensic bite mark analysis in legal proceedings has been brought into question because of lack of scientific validity, and as a result, the ABFO no longer endorses positive identification of a suspect dentition to a bitemark.

Dental neglect has been recognized for years to be a significant child health issue. Pediatricians, dental professionals, and other physicians can be alert to children and youth at particular risk for orofacial and dental neglect and its psychosocial consequences, such as LGBTQ+ youth who are gender

nonconforming, children who are overweight or obese, youth who are subjected to human trafficking, and children with special health care needs.⁵⁵⁻⁵⁷

Medical and dental collaboration to prevent child maltreatment is important. The National Maternal and Child Oral Health Resource Center (Special Care | Module 3: Oral Health Supervision [mchoralhealth.org]) has resources for health care providers seeking information on this issue.⁸¹ Pediatric dentists and oral and maxillofacial surgeons, whose advanced education programs include a mandated child abuse curriculum, can provide valuable information and assistance to other health care providers about oral and dental aspects of child abuse and neglect. Physician members of multidisciplinary child abuse and neglect teams are encouraged to identify such dental providers in their communities to serve as consultants for these teams. In addition, physicians with experience or expertise in child abuse and dental neglect can make themselves available to dentists and dental organizations as consultants and educators. Such collaborative efforts will strengthen the ability to prevent and detect child abuse and neglect and enhance the ability to care for and protect children.

RECOMMENDATIONS

1. Pediatricians, dental professionals, and other physicians must report injuries that are concerning for abuse or neglect to child protective services in accordance with local or state legal requirements. For more information about requirements in your state, contact the AAP state advocacy team (e-mail: stgov@aap.org).
2. In cases of sexual abuse that may involve the mouth, even without overt signs, pediatricians, dental professionals, and other physicians should know how to collect a history to elicit this information and to perform oral STI testing using appropriate tests to support forensic investigations. When sexual abuse is suspected or diagnosed in a child, the case needs to be reported to child protective services and/or law enforcement agencies for investigation.⁸²⁻⁸⁵ A multidisciplinary child abuse evaluation for the child and family is preferred when available.
3. Consultation with a child abuse pediatrician may be helpful when a pediatrician, dental professional, or other physician suspects MCA.
4. Because ABFO guidelines no longer endorse positive identification of a suspect dentition to a bitemark, pediatricians, dental professionals, and other physicians should consult with a forensic odontologist, pediatric dentist experienced in forensic issues, or child abuse pediatrician for help in evaluating bite marks. To locate an expert and review the standards and guidelines for evaluating bite marks, go to www.abfo.org.
5. Because children with dental abnormalities are at increased risk for bullying, pediatricians, dental professionals, and other physicians should ask their patients about bullying and advocate for antibullying prevention programs in schools and other community settings.
6. Because children who have experienced human trafficking may visit the health care setting (dental and medical), pediatricians, dental professionals, and other physicians are encouraged to be aware of human trafficking screening tools and the risk factors for human trafficking to identify children at risk (of all genders) and to safely report and connect patients to resources. They can also advocate for antitrafficking efforts.
7. If parents fail to obtain necessary treatment of a child's oral or dental disease after barriers to care have been addressed, the case must be reported to the appropriate child protective services agency as concerning for dental neglect.
8. Pediatricians, dental professionals, and other physicians are encouraged to consult with colleagues on challenging cases and collaborate with them to provide education and support toward improved identification of child maltreatment.
9. Pediatricians, dental professionals, and other physicians should recognize and acknowledge implicit bias and reduce its influence on diagnosing and reporting child maltreatment.

LEAD AUTHORS

Anupama Rao Tate, DMD, MPH
Susan A. Fisher-Owens, MD, MPH, FAAP Lora Spiller, MD, FAAP
Jillian Muhlbauer, DMD
James L. Lukefahr, MD, FAAP

SECTION ON ORAL HEALTH EXECUTIVE COMMITTEE, 2022–2023

Jeffrey M. Karp, DMD, MS, Chairperson Eric L. Beyer, MD, IBCLC, FABM, FAAP
Chevon Brooks, MD, FAAP
Lauren M. Feldman, DMD
C. Eve Kimball, MD, FAAP Anupama Rao Tate, DMD, MPH

PAST SECTION ON ORAL HEALTH EXECUTIVE COMMITTEE MEMBERS

Karen Sokal-Gutierrez, MD, MPH, FAAP
John H. Unkel, DDS, MD, MPA, FAAP

SECTION ON ORAL HEALTH LIAISONS

Matt Zaborowski, MPH, CPH, American Dental Association
Matt Crespín, MPH, RDH, American Dental Hygienists' Association
Amr Moursi, DDS, PhD, American Academy of Pediatric Dentistry

SECTION ON ORAL HEALTH STAFF

Kera Beskin, MPH, MBA

COUNCIL ON CHILD ABUSE AND NEGLECT, 2022–2023

Suzanne Breen Haney, MD, MS, FAAP, Chairperson
Antoinette Laskey, MD, MPH, MBA, FAAP, Chairperson-elect
Andrea Asnes, MD, MSW, FAAP
Verena Brown, MD, FAAP
Rebecca G. Girardet, MD, FAAP

Nancy Heavilin, MD, FAAP
Amanda Bird Hoffert Gilmartin, MD, FAAP
Natalie Kissoon, MD, FAAP
Bethany Anne Mohr, MD, FAAP
Patricia Morgan, MD, FAAP
Shalon Marie Nienow, MD, FAAP
Norell Rosado, MD, FAAP

COUNCIL ON CHILD ABUSE AND NEGLECT LIAISONS

Rachael Keefe, MD, MPH, FAAP, Council on Foster Care, Adoption, and Kinship Care
Brooks Keeshin, MD, FAAP, American Academy of Child and Adolescent Psychiatry
Jennifer Matjasko, PhD, Centers for Disease Control and Prevention
Serena Wagoner, DO, Section on Pediatric Trainees
Elaine Stedt, MSW, ACSW, Administration for Children, Youth and Families, Office on Child Abuse and Neglect

STAFF

Tammy Piazza Hurley
Jeff Hudson, MA

ABBREVIATIONS

AAP: American Academy of Pediatrics.
ABFO: American Board of Forensic Odontology.
MCA: medical child abuse.
NAAT: nucleic acid amplification test.
STI: sexually transmitted infection.

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2024 by the American Academy of Pediatrics.

FUNDING: No external funding.

FINANCIAL/CONFLICT OF INTEREST DISCLOSURES: Dr Fischer-Owens has disclosed a financial relationship with Colgate, as a DSMB member. Disclosures are reviewed and mitigated through a Conflict of Interest process approved by the AAP Board of Directors.

REFERENCES

- Centers for Disease Control and Prevention. Fast facts: Preventing child abuse and neglect. Available at: "https://www.cdc.gov/violenceprevention/childabuseandneglect/fastfact.html#:~:text=At%20least%201%20in%207,neglect%20in%20the%20United%20States". Accessed November 29, 2023.
- Substance Abuse and Mental Health Services Administration. Intimate partner violence and child abuse considerations during COVID-19. Available at: "https://www.samhsa.gov/sites/default/files/social-distancing-domestic-violence.pdf". Accessed November 29, 2023.
- Lane WG, Rubin DM, Monteith R, Christian CW. Racial differences in the evaluation of pediatric fractures for physical abuse. *JAMA*. 2002; 288(13):1603-1609.
- Legano LA, Desch LW, Messner SA, Idzerda S, Flaherty EG. Council on Child Abuse and Neglect; Council on Children With Disabilities. Maltreatment of children with disabilities. *Pediatrics*. 2021;147(5): e2021050920.
- The Joint Commission; Quick Safety. Addressing intimate partner violence and helping to protect patients. Available at: "https://www.jointcommission.org/-/media/tjc/newsletters/quick-safety-63-addressing-ipv-final3-1-10-22.pdf". Accessed November 29, 2023.
- Jenny C, Hymel KP, Ritzen A, Reinert SE, Hay TC. Analysis of missed cases of abusive head trauma. *JAMA*. 1999;281(7):621-626.
- Wakeman SE, Bryant A, Harrison N. Redefining child protection: Addressing the harms of structural racism and punitive approaches for birthing people, dyads, and families affected by substance use. *Obstet Gynecol*. 2022;140(2):167-173.
- Feely M, Bosk EA. That which is essential has been made invisible: The need to bring a structural risk perspective to reduce racial disproportionality in child welfare. *Race Soc Probl*. 2021;13(1):49-62.
- Berkman E, Brown E, Scott M, Adiele A. Racism in child welfare: Ethical considerations of harm. *Bioethics*. 2022;36(3):298-304.
- Sedlak AJ, Mettenberg J, Basena M, et al. Fourth National Incidence Study of Child Abuse and Neglect (NIS-4): Report to Congress, Executive Summary. US Department of Health and Human Services, Administration for Children and Families; 2010.
- Christian CW. Committee on Child Abuse and Neglect, American Academy of Pediatrics. The evaluation of suspected child physical abuse. *Pediatrics*. 2015;135(5):e1337-e1354.
- Sheets LK, Leach ME, Koszewski IJ, Lessmeier AM, Nugent M, Simpson P. Sentinel injuries in infants evaluated for child physical abuse. *Pediatrics*. 2013;131(4):701-707.
- Pierce MC, Kaczor K, Aldridge S, O'Flynn J, Lorenz DJ. Bruising characteristics discriminating physical child abuse from accidental trauma. *Pediatrics*. 2010;125(1):67-74.

14. Pierce MC, Kaczor K, Lorenz DJ, et al. Validation of a clinical decision rule to predict abuse in young children based on bruising characteristics. *JAMA Netw Open*. 2021;4(4):e215832.
15. Cairns AM, Mok JY, Welbury RR. Injuries to the head, face, mouth, and neck in physically abused children in a community setting. *Int J Paediatr Dent*. 2005;15(5):310-318.
16. Crouse CD, Faust RA. Child abuse and the otolaryngologist: Part II. *Otolaryngol Head Neck Surg*. 2003;128(3):311-317.
17. Cavalcanti AL. Prevalence and characteristics of injuries to the head and orofacial region in physically abused children and adolescents—A retrospective study in a city of the Northeast of Brazil. *Dent Traumatol*. 2010;26(2):149-153.
18. da Fonseca MA, Feigal RJ, ten Bense RW. Dental aspects of 1248 cases of child maltreatment on file at a major county hospital. *Pediatr Dent*. 1992;14(3):152-157.
19. Valencia-Rojas N, Lawrence HP, Goodman D. Prevalence of early childhood caries in a population of children with history of maltreatment. *J Public Health Dent*. 2008;68(2):94-101.
20. Sarkar R, Ozanne-Smith J, Bassed R. Systematic review of the patterns of orofacial injuries in physically abused children and adolescents. *Trauma Violence Abuse*. 2021;22(1):136-146.
21. Sarkar R, Bassed R, Dipnall JF, Ozanne-Smith J. Orofacial injuries in child family homicide: A population study. *Forensic Sci Med Pathol*. 2021;17(4):553-564.
22. Dorfman MV, Metz JB, Feldman KW, Farris R, Lindberg DM. ExSTRA Investigators. Oral injuries and occult harm in children evaluated for abuse. *Arch Dis Child*. 2018;103(8):747-752.
23. Naidoo S. A profile of the orofacial injuries in child physical abuse at a children's hospital. *Child Abuse Negl*. 2000;24(4):521-534.
24. Maguire S, Hunter B, Hunter L, Sibert JR, Mann M, Kemp AM. Welsh Child Protection Systematic Review Group. Diagnosing abuse: A systematic review of torn frenum and other intraoral injuries. *Arch Dis Child*. 2007;92(12):1113-1117.
25. Keels MA. Section on Oral Health, American Academy of Pediatrics. Management of dental trauma in a primary care setting. *Pediatrics*. 2014;133(2):e466-e476.
26. Griest KJ. Supporting evidence in physical child abuse. In: Griest KJ, ed. *Pediatric Homicide: Medical Investigation*. CRC Press; 2010:144-146.
27. Levin AV. Otorhinolaryngologic manifestations. In: Levin AV, Sheridan MS, eds. *Munchausen Syndrome by Proxy: Issues in Diagnosis and Treatment*. Lexington Books; 1995:219-230.
28. Woods CRW. Gonococcal infections in neonates and young children. *Semin Pediatr Infect Dis*. 2005;16(4):258-270.
29. Workowski KA, Bachmann LH, Chan PA, et al. Centers for Disease Control and Prevention. Sexually transmitted infections treatment guidelines, 2021. *MMWR Recomm Rep*. 2021;70(4):1-187.
30. De Jong AR. Sexually transmitted diseases in sexually abused children. *Sex Transm Dis* 1986;13(3):123-126.
31. Everett VD, Ingram DL, Flick LAR, Russell TA, Tropez-Sims ST, McFadden AY. A comparison of sexually transmitted diseases (STDs) found in a total of 696 boys and 2973 girls evaluated for sexual abuse. *Pediatr Res* 1998;43(4 Pt 2):91A.
32. Gavril AR, Kellogg ND, Nair P. Value of follow-up examinations of children and adolescents evaluated for sexual abuse and assault. *Pediatrics*. 2012;129(2):282-289.
33. Nelson JD, Mohs E, Dajani AS, Plotkin SA. Gonorrhea in preschool- and school-aged children. Report of the prepubertal gonorrhea cooperative study group. *JAMA*. 1976;236(12):1359-1364.
34. Kellogg ND, Melville JD, Lukefahr JL, Nienow SM, Russell EL. Genital and extragenital gonorrhea and chlamydia in children and adolescents evaluated for sexual abuse. *Pediatr Emerg Care*. 2018;34(11):761-766.
35. Stevens-Simon C, Nelligan D, Breese P, Jenny C, Douglas JM Jr. The prevalence of genital human papillomavirus infections in abused and nonabused preadolescent girls. *Pediatrics*. 2000;106(4):645-649.
36. Bussen S, S Dutterlin M, Schmidt U, Bussen D. Anogenital warts in childhood—Always a marker for sexual abuse? *Geburtshilfe Frauenheilkd* 2012;72(1):43-48.
37. Crawford-Jakubiak JE, Alderman EM, Leventhal JM. Committee on Child Abuse and Neglect; Committee on Adolescence. Care of the adolescent after an acute sexual assault. *Pediatrics*. 2017;139(3):e20164243.
38. Roesler TA, Jenny C. *Medical Child Abuse: Beyond Munchausen by Proxy*. American Academy of Pediatrics; 2008.
39. Feldman KW, Ambartsumyan L, Goldin A, et al. Gastrointestinal problems and nonoral feeding in children evaluated for medical child abuse (fabricated and induced illness). *Child Abuse Rev*. 2022;31(3):1-10.
40. Olczak-Kowalczyk D, Wolska-Kusnierz B, Bernatowska E. Fabricated or induced illness in the oral cavity in children. A systematic review and personal experience. *Cent Eur J Immunol*. 2015;40(1):109-114.
41. Yates G, Bass C. The perpetrators of medical child abuse (Munchausen syndrome by proxy)—A systematic review of 796 cases. *Child Abuse Negl*. 2017;72:45-53.
42. Flaherty EG, Macmillan HL. Committee On Child Abuse And Neglect. Caregiver-fabricated illness in a child: A manifestation of child maltreatment. *Pediatrics*. 2013;132(3):590-597.
43. Silva RHA, de Oliveira Musse J, Melani RFH, Oliveira RN. Human bite mark identification and DNA technology in forensic dentistry. *Braz J Oral Sci*. 2006;5(19):1193-1197.
44. Jessee S, Deinaud S. *Dentalcare.com*. Child abuse and neglect: Implications for the dental professional. Available at: “https://i-med.sk/sites/default/files/viaco/child_abuse_and_neglect_implications_for_the_dental_professional.pdf”. Accessed November 29, 2023.
45. Wagner GN. Bitemark identification in child abuse cases. *Pediatr Dent*. 1986;8(1 Spec No):96-100.
46. Barsley RE, Bernstein ML, Brumit PC, et al. Epidermis and enamel: Insights into gnawing criticisms of human bitemark evidence. *Am J Forensic Med Pathol*. 2018;39(2):87-97.
47. Committee on Injury, Violence, and Poison Prevention. Policy statement—role of the pediatrician in youth violence prevention. *Pediatrics*. 2009;124(1):393-402.
48. Seehra J, Fleming PS, Newton T, DiBiase AT. Bullying in orthodontic patients and its relationship to malocclusion, self-esteem and oral health-related quality of life. *J Orthod*. 2011;38(4):247-256, quiz 294.
49. Al-Bitar ZB, Al-Omari IK, Sonbol HN, Al-Ahmad HT, Cunningham SJ. Bullying among Jordanian schoolchildren, its effects on school performance, and the contribution of general physical and dentofacial features. *Am J Orthod Dentofacial Orthop*. 2013;144(6):872-878.
50. Islam MI, Chadwick V, Esgin T, Martiniuk A. Bullied because of their teeth: Evidence from a longitudinal study on the impact of oral health on bullying victimization among Australian Indigenous children. *Int J Environ Res Public Health*. 2022;19(9):4995.
51. Hunt O, Burden D, Hepper P, Stevenson M, Johnston C. Self-reports of psychosocial functioning among children and young adults with cleft lip and palate. *Cleft Palate Craniofac J*. 2006;43(5):598-605.
52. Al-Omari IK, Al-Bitar ZB, Sonbol HN, Al-Ahmad HT, Cunningham SJ, Al-Omiri M. Impact of bullying due to dentofacial features on oral health-related quality of life. *Am J Orthod Dentofacial Orthop*. 2014;146(6):734-739.
53. Seehra J, Newton JT, DiBiase AT. Bullying in schoolchildren—its relationship to dental appearance and psychosocial implications: An update for GDPs. *Br Dent J*. 2011;210(9):411-415.
54. Kvist T, Annerbäck EM, Sahlqvist L, Flodmark O, Dahlidof G. Association between adolescents' self-perceived oral health and self-reported experiences of abuse. *Eur J Oral Sci*. 2013;121(6):594-599.
55. Earnshaw VA, Reisner SL, Juvonen J, Hatzembuehler ML, Perrotti J, Schuster MA. LGBTQ bullying: Translating research to action in pediatricians. *Pediatrics*. 2017;140(4):e20170432.

References continued on the next page.

56. Lumeng JC, Forrest P, Appugliese DP, Kaciroti N, Corwyn RF, Bradley RH. Weight status as a predictor of being bullied in third through sixth grades. *Pediatrics*. 2010;125(6):e1301-e1307.
57. Gage NA, Katsiyannis A, Rose C, Adams SE. Disproportionate bullying victimization and perpetration by disability status, race, and gender: A national analysis. *Adv Neurodev Disord*. 2021;5(3):256-268.
58. Van Cleave J, Davis MM. Bullying and peer victimization among children with special health care needs. *Pediatrics*. 2006;118(4):e1212-e1219.
59. US Department of State. Trafficking in persons report. Available at: "<https://www.state.gov/wp-content/uploads/2021/09/TIPR-GPA-upload-07222021.pdf>". Accessed May 18, 2022.
60. Smith LA, Vardaman SH, Snow MA. Shared Hope International. The national report on domestic minor sex trafficking: America's prostituted children. Available at: "http://sharedhope.org/wp-content/uploads/2012/09/SHI_National_Report_on_DMST_2009.pdf". Accessed May 23, 2016.
61. US Department of Education. Human trafficking of children in the United States: A fact sheet for schools. Available at: "<http://www2.ed.gov/about/offices/list/osdfs/factsheet.pdf>". Accessed May 18, 2016.
62. Swaner R, Labriola M, Rempel M, Walker A, Spadafore J. Youth Involvement in the Sex Trade: A National Study. Center for Court Innovation; 2016.
63. Middleton JS, Gattis MN, Frey LM, Roe-Sepowitz D. Youth experiences survey (YES): Exploring the scope and complexity of sex trafficking in a sample of youth experiencing homelessness. *J Soc Serv Res*. 2018;44:141-157.
64. Greenbaum J, Crawford-Jakubiak JE. Committee on Child Abuse and Neglect. Child sex trafficking and commercial sexual exploitation: Health care needs of victims. *Pediatrics*. 2015;135(3):566-574.
65. Zimmerman C, Hossain M, Yun K, et al. The health of trafficked women: A survey of women entering posttrafficking services in Europe. *Am J Public Health*. 2008;98(1):55-59.
66. Lederer LJ, Wetzel CA. The health consequences of sex trafficking and their implications for identifying victims in healthcare facilities. *Ann Health Law*. 2014;23(1):61-91.
67. Crane PA, Moreno M. Human trafficking: What is the role of the health care provider? *J Appl Res Child*. 2011;2(1):7.
68. Hornor G, Sherfield J. Commercial sexual exploitation of children: Health care use and case characteristics. *J Pediatr Health Care*. 2018;32(3):250-262.
69. Chisolm-Straker M, Baldwin S, Gaigb e-Togbe B, Ndukwe N, Johnson PN, Richardson LD. Health care and human trafficking: We are seeing the unseen. *J Health Care Poor Underserved*. 2016;27(3):1220-1233.
70. Greenbaum VJ, Dodd M, McCracken C. A short screening tool to identify victims of child sex trafficking in the health care setting. *Pediatr Emerg Care*. 2018;34(1):33-37.
71. American Academy of Pediatric Dentistry. Policy on social determinants of children's oral health and health disparities. In: *The Reference Manual of Pediatric Dentistry*. American Academy of Pediatric Dentistry; 2022:29-33.
72. American Academy of Pediatric Dentistry. Definition of dental neglect. In: *The Reference Manual of Pediatric Dentistry*. American Academy of Pediatric Dentistry; 2021:16.
73. Sanger RG, Bross DC. *Clinical Management of Child Abuse and Neglect: A Guide for the Dental Profession*. Quintessence Publishing Company; 1984.
74. Kilpatrick NM, Scott J, Robinson S. Child protection: A survey of experience and knowledge within the dental profession of New South Wales, Australia. *Int J Paediatr Dent*. 1999;9(3):153-159.
75. Kiran K, Kamala BK. Child abuse and the role of a dental professional—The Indian scenario. *Child Abuse Negl*. 2011;35(3):157-158.
76. Krol DM, Whelan K. Section on Oral Health. Maintaining and improving the oral health of young children. *Pediatrics*. 2023;151(1):e2022060417.
77. Farrell M, Erickson MH, Labella BE. Child neglect. In: Klika JB, Conte JR, eds. *The APSAC Handbook on Child Maltreatment*, 4th ed. Sage Publications; 2017:127-144.
78. California Society of Pediatric Dentists. Dental neglect: When to report. *California Pediatrician*. 1989;(Fall):31-32.
79. Krug EG, Dahlberg LL, Mercy JA, Zwi AB, Lozano R. World Health Organization. World report on violence and health. Available at: "www.who.int/violence_injury_prevention/violence/world_report/en/". Accessed May 23, 2016.
80. Bhatia SK, Maguire SA, Chadwick BL, et al. Characteristics of child dental neglect: A systematic review. *J Dent*. 2014;42(3):229-239.
81. National Maternal and Child Health Resource Center. Oral health supervision module 3.4: Child abuse and neglect. Available at: "<https://www.mchoralhealth.org/SpecialCare/3-supervision/3-4.php>". Accessed June 7, 2024.
82. Mouden LD, Bross DC. Legal issues affecting dentistry's role in preventing child abuse and neglect. *J Am Dent Assoc*. 1995;126(8):1173-1180.
83. Katner DR, Brown CE. Mandatory reporting of oral injuries indicating possible child abuse. *J Am Dent Assoc*. 2012;143(10):1087-1092.
84. Blain SM, Skerber N, Shanel-Hogan KA. Oral and dental aspects of child abuse and neglect. In: Peterson MS, Durfee M, eds. *Child Abuse and Neglect: Guidelines for Identification, Assessment, and Case Management*. Volcano Press; 2003:58-60.
85. Brown R, Hines L. Maxillofacial, Neck, and Dental Manifestations of Child Abuse. In: Laskey A, Sirotnak A, eds. *Child Abuse: Medical Diagnosis and Management*. American Academy of Pediatrics; 2020:167-197.