

Management of Medical Emergencies

For all emergencies

1. Discontinue dental treatment
2. Call for assistance /someone to bring oxygen and emergency kit
3. Position patient: ensure open and unobstructed airway
4. Monitor vital signs
5. Be prepared to support respiration, support circulation, provide CPR, and call for emergency medical services

Condition	Signs and symptoms	Treatment	Drug dosage	Drug delivery
Allergic reaction (mild or delayed)	Hives; itching; edema; erythema—skin, mucosa conjunctiva	<ol style="list-style-type: none"> 1. Discontinue all sources of allergy-causing substances 2. Administer diphenhydramine 	Diphenhydramine: follow manufacturer's instructions based on child's age/weight	Oral
Allergic reaction (sudden onset): anaphylaxis	Urticaria-itching, flushing, hives; rhinitis; wheezing/difficulty breathing; bronchospasm; laryngeal edema; weak pulse; marked fall in blood pressure; loss of consciousness	<p>This is a true, life-threatening emergency</p> <ol style="list-style-type: none"> 1. Call for emergency medical services 2. Administer epinephrine 3. Administer oxygen 4. Monitor vital signs 5. Transport to emergency medical facility by advanced medical responders 	Epinephrine (1 mg/mL): 0.01 mg/kg every 5 minutes until recovery or until help arrives ¹	IM or SubQ (Auto injector available)
Acute asthmatic attack	Shortness of breath; wheezing; coughing; tightness in chest; cyanosis; tachycardia	<ol style="list-style-type: none"> 1. Sit patient upright or in a comfortable position 2. Administer oxygen 3. Administer bronchodilator 4. If bronchodilator is ineffective, administer epinephrine 5. Call for emergency medical services with transportation for advanced care if indicated 	<ol style="list-style-type: none"> 1. Albuterol (patient's or emergency kit inhaler) 2. Epinephrine (1 mg/mL): 0.01 mg/kg every 15 minutes as needed¹ 	Inhale IM or SubQ
Local anesthetic toxicity	Light-headedness; changes in vision and/or speech; metallic taste; changes in mental status; confusion, agitation; tinnitus; tremor; seizure; tachypnea; bradycardia; unconsciousness; cardiac arrest	<ol style="list-style-type: none"> 1. Assess and support airway, breathing, and circulation (CPR if warranted) 2. Administer oxygen 3. Monitor vital signs 4. Call for emergency medical services with transportation for advanced care if indicated 	Supplemental oxygen	Mask
Local anesthetic reaction: vasoconstrictor	Anxiety; tachycardia/palpitations; restlessness; headache; tachypnea; chest pain; cardiac arrest	<ol style="list-style-type: none"> 1. Reassure patient 2. Assess and support airway, breathing, and circulation (CPR if warranted) 3. Administer oxygen 4. Monitor vital signs 5. Call for emergency medical services with transportation for advanced care if indicated 	Supplemental oxygen	Mask
Overdose: benzodiazepine	Somnolence; confusion; diminished reflexes; respiratory depression; apnea; respiratory arrest; cardiac arrest	<ol style="list-style-type: none"> 1. Assess and support airway, breathing, and circulation (CPR if warranted) 2. Administer oxygen 3. Monitor vital signs 4. If severe respiratory depression, establish IV access and reverse with flumazenil 5. Monitor recovery (for at least 2 hours after the last dose of flumazenil) and call for emergency medical services with transportation for advanced care if indicated 	Flumazenil 0.01 mg/kg (maximum: 0.2 mg); may repeat at 1 minute intervals not to exceed a cumulative dose of 0.05 mg/kg or 1 mg, whichever is less ¹	IV (if IV access is not available, may be given IM)

Abbreviations in table: CPR=cardiopulmonary resuscitation; IM=intramuscular; IN=intranasal; IV=intravenous; kg=kilogram; mg=milligram; mL=milliliter; SubQ=subcutaneous.

For all emergencies

1. Discontinue dental treatment
2. Call for assistance / someone to bring oxygen and emergency kit
3. Position patient: ensure open and unobstructed airway
4. Monitor vital signs
5. Be prepared to support respiration, support circulation, provide CPR, and call for emergency medical services

Condition	Signs and symptoms	Treatment	Drug dosage	Drug delivery
Overdose: narcotic	Decreased responsiveness; respiratory depression; respiratory arrest; cardiac arrest	<ol style="list-style-type: none"> 1. Assess and support airway, breathing, and circulation (CPR if warranted) 2. Administer oxygen 3. Monitor vital signs 4. If severe respiratory depression, reverse with naxolone 5. Monitor recovery (for at least 2 hours after the last dose of naxolone) and call for emergency medical services with transportation for advanced care if indicated 	Naxolone 0.1 mg/kg up to 2 mg. ^{1,2} May be repeated to maintain reversal.	IV, IM, or SubQ
Seizure	Warning aura: disorientation, blinking, or blank stare; uncontrolled muscle movements; muscle rigidity; unconsciousness; postictal phase—sleepiness, confusion, amnesia, slow recovery	<ol style="list-style-type: none"> 1. Recline and position to prevent injury 2. Ensure open airway and adequate ventilation 3. Monitor vital signs 4. If status is epilepticus, give either diazepam OR midazolam and call for emergency medical services with transportation for advanced care if indicated 	Diazepam (5 mg/mL): 0.15-0.2 mg/kg per dose; maximum 10 mg per dose. May repeat dose once. ¹ OR Midazolam: 0.2 mg/kg (maximum 10 mg) ¹	IV IM, IN, IV
Syncope (fainting)	Feeling of warmth; skin pale and moist; pulse rapid initially then gets slow and weak; dizziness; hypotension; cold extremities; unconsciousness	<ol style="list-style-type: none"> 1. Recline, feet up 2. Loosen clothing that may be binding 3. Ammonia inhaler 4. Administer oxygen 5. Cold towel on back of neck 6. Monitor recovery 	Ammonia in vials	Inhale

Abbreviations in table: CPR=cardiopulmonary resuscitation; IM=intramuscular; IN=intranasal; IV=intravenous; kg=kilogram; mg=milligram; mL=milliliter; SubQ=subcutaneous.

Reference

1. Sheno RP, Timm N, AAP Committee on Drugs, AAP Committee on Pediatric Emergency Medicine. Drugs used to treat pediatric emergencies. *Pediatrics* 2020;145(1):e20193450. Available at: "https://publications.aap.org/pediatrics/article/145/1/e20193450/36970/Drugs-Used-to-Treat-Pediatric-Emergencies". Accessed August 24, 2022.

DISCLAIMER: This information is not intended to be a comprehensive list of all medications that may be used in all emergencies. Drug information is constantly changing and is often subject to interpretation. While care has been taken to ensure the accuracy of the information presented, the AAPD is not responsible for the continued currency of the information, errors, omissions, or the resulting consequences. Decisions about drug therapy must be based upon the independent judgment of the clinician, changing drug information, and evolving healthcare practices.